

Some Cases of the Mumford–Tate Conjecture and Shimura Varieties

Adrian Vasiu, Binghamton University

December 18, 2007

Final version, to appear in Indiana Univ. Math. J.

Dedicated to Jean-Pierre Serre, on his 81th anniversary

ABSTRACT. We prove the Mumford–Tate conjecture for those abelian varieties over number fields whose extensions to $\mathbb{C}$ have attached adjoint Shimura varieties that are products of simple, adjoint Shimura varieties of certain Shimura types. In particular, we prove the conjecture for the orthogonal case (i.e., for the B_n and $D_n^{\mathbb{R}}$ Shimura types). As a main tool, we construct embeddings of Shimura varieties (whose adjoints are) of prescribed abelian type into unitary Shimura varieties of PEL type. These constructions implicitly classify the adjoints of Shimura varieties of PEL type.

Key words: abelian and Shimura varieties, reductive and homology groups, p -adic Galois representations, Frobenius tori, and Hodge cycles.

MSC 2000: Primary 11G10, 11G15, 11G18, 11R32, 14G35, and 14G40.

Contents

1. Introduction	1
2. Some complements on Shimura pairs	11
3. Basic techniques and the proof of Theorem 1.3.1	17
4. Injective maps into unitary Shimura pairs of PEL type	22
5. Applications to Frobenius tori	34
6. Non-special A_n types	42
7. The proof of the Main Theorem	48
References	63

1. Introduction

If K is a field, let $\overline{K}$ be an algebraic closure of K . If M is a free module of finite rank over a commutative ring R with unit, let $M^* := \text{Hom}_R(M, R)$ and let $\mathbf{GL}_M$ be

the group scheme over R of linear automorphisms of M . If $*_R$ or $*$ is either an object or a morphism of the category of $\mathrm{Spec}(R)$ -schemes, let $*_U$ be its pull back via an affine morphism $m : \mathrm{Spec}(U) \rightarrow \mathrm{Spec}(R)$. If there exist several ways to take m , we mention the homomorphism $R \rightarrow U$ that defines m but often we do not add it to the notation $*_U$.

1.1. The conjecture. Let E be a number field. We fix an embedding $i_E : E \hookrightarrow \mathbb{C}$. To fix the ideas, we will take $\overline{E} = \overline{\mathbb{Q}}$ to be the algebraic closures of $E = i_E(E)$ and $\mathbb{Q}$ in $\mathbb{C}$. Let A be an abelian variety over E . For the simplicity of notations, let

$$L_A := H_1(A(\mathbb{C}), \mathbb{Z})$$

be the first homology group of the complex manifold $A(\mathbb{C})$ with coefficients in $\mathbb{Z}$, let

$$\mu_A : \mathbb{G}_{m, \mathbb{C}} \rightarrow \mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{C}}$$

be the *Hodge cocharacter*, and let H_A be the *Mumford–Tate group* of $A_{\mathbb{C}}$. We recall that if $L_A \otimes_{\mathbb{Z}} \mathbb{C} = F^{-1,0} \oplus F^{0,-1}$ is the classical Hodge decomposition, then $\beta \in \mathbb{G}_{m, \mathbb{C}}(\mathbb{C})$ acts through μ_A trivially on $F^{0,-1}$ and as the multiplication with β on $F^{-1,0}$. We also recall that H_A is a *reductive group* over $\mathbb{Q}$ and that H_A is the smallest subgroup of $\mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}}$ with the property that μ_A factors through $H_{A, \mathbb{C}}$, cf. [De3, Props. 3.4 and 3.6].

Let $p \in \mathbb{N}$ be a prime. Let $T_p(A)$ be the *Tate-module* of A . As a $\mathbb{Z}_p$ -module (resp. as a $\mathrm{Gal}(E)$ -module) we identify it canonically with $L_A \otimes_{\mathbb{Z}} \mathbb{Z}_p$ (resp. $\mathrm{Hom}_{\mathbb{Z}_p}(H_{\acute{e}t}^1(A_{\overline{E}}, \mathbb{Z}_p), \mathbb{Z}_p)$). Let $G_{\mathbb{Q}_p}$ be the *identity component of the algebraic envelope* of the p -adic Galois representation

$$\rho : \mathrm{Gal}(E) \rightarrow \mathbf{GL}_{T_p(A) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p}(\mathbb{Q}_p) = \mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p}(\mathbb{Q}_p),$$

cf. [Bog] and [Se3, Sect. 135]. Let E^{conn} be the smallest finite field extension of E that is contained in $\overline{\mathbb{Q}}$ and that has the property that $\rho(\mathrm{Gal}(E^{\mathrm{conn}}))$ is a compact, open subgroup of $G_{\mathbb{Q}_p}(\mathbb{Q}_p)$ (see [Bog]). It is known that the field E^{conn} does not depend on the prime p (see [Se3, p. 15]). The Mumford–Tate conjecture says (see [Mu1] and [Se2, Sect. 9]):

1.1.1. Conjecture. *As subgroups of $\mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p}$ we have $G_{\mathbb{Q}_p} = H_{A, \mathbb{Q}_p}$.*

The main goal of the paper is to prove the Conjecture in many cases. We first recall fundamental results that pertain to the Conjecture and that play a key role in the paper.

1.1.2. Theorem (Pyatetskii-Shapiro–Deligne–Borovoi). *As subgroups of $\mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p}$, $G_{\mathbb{Q}_p}$ is a subgroup of $H_{A, \mathbb{Q}_p}$ (for instance, cf. [De3, Prop. 2.9 and Thm. 2.11]).*

1.1.3. Theorem (Faltings). *The group $G_{\mathbb{Q}_p}$ is reductive (cf. [Fa]).*

1.1.4. Theorem (Faltings). *The centralizers of $G_{\mathbb{Q}_p}$ and $H_{A, \mathbb{Q}_p}$ in $\mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p}$ coincide; more precisely the centralizers of $G_{\mathbb{Q}_p}(\mathbb{Q}_p)$ and $H_{A, \mathbb{Q}_p}(\mathbb{Q}_p)$ in $\mathrm{End}(L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p)$ are equal to the semisimple $\mathbb{Q}_p$ -subalgebra $\mathrm{End}(A_{\overline{E}}) \otimes_{\mathbb{Z}} \mathbb{Q}_p$ of $\mathrm{End}(L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p)$ (cf. [Fa]).*

Let v be a prime of E such that A has good reduction A_v with respect to it. Let $l \in \mathbb{N}$ be the prime divided by v . Let T_v be the *Frobenius torus* of the abelian variety A_v (see [Ch]); it is a torus over $\mathbb{Q}$ that depends only on the isogeny class of A_v . If $p \neq l$, then $T_{v, \mathbb{Q}_p}$ is naturally a torus of $G_{\mathbb{Q}_p}$, uniquely determined up to $\mathrm{Im}(\rho)$ -conjugation.

1.1.5. Theorem (Serre). *Let $\mathfrak{P}$ be the set of primes v of E which are of good reduction for A , which are relatively prime to p , and for which $T_{v, \mathbb{Q}_p}$ is a maximal torus of $G_{\mathbb{Q}_p}$. If $E = E^{\text{conn}}$, then the set $\mathfrak{P}$ has Dirichlet density 1 (cf. [Ch, Cor. 3.8]). Thus the rank of $G_{\mathbb{Q}_p}$ does not depend on p (cf. [Ch, Thm. 3.10]).*

1.1.6. Theorem (Larsen–Pink–Tankeev). *If the Mumford–Tate conjecture for the abelian variety A is true for a prime p , then it is true for all rational primes (cf. either [LP, Thm. 4.3] or [Ta1, Lem. 3.4]).*

The proof of [Ta1, Lem. 3.4] relies on Theorems 1.1.2 to 1.1.5 and on the following Lemma.

1.1.7. Lemma (Zarhin). *Let $\mathcal{V}$ be a finite dimensional vector space over a field of characteristic 0. Let $\mathfrak{g}_1 \subseteq \mathfrak{g}_2 \subseteq \text{End}(\mathcal{V})$ be Lie algebras of reductive subgroups of $\mathbf{GL}_{\mathcal{V}}$. We assume that the centralizers of $\mathfrak{g}_1$ and $\mathfrak{g}_2$ in $\text{End}(\mathcal{V})$ coincide and that the ranks of $\mathfrak{g}_1$ and $\mathfrak{g}_2$ are equal. Then $\mathfrak{g}_1 = \mathfrak{g}_2$ (cf. [Za1, Sect. 5, Key Lemma]).*

1.1.8. Theorem (Pink). *The group $G_{\overline{\mathbb{Q}_p}}$ is generated by cocharacters of it with the property that the resulting representations of $\mathbb{G}_{m, \overline{\mathbb{Q}_p}}$ on $L_A \otimes_{\mathbb{Z}} \overline{\mathbb{Q}_p}$ are associated to the trivial and the identity characters of $\mathbb{G}_{m, \overline{\mathbb{Q}_p}}$, the multiplicities being equal (cf. [Pi, Thms. 3.18 and 5.10]).*

1.1.9. Corollary (Pink). *Every simple factor $\mathcal{F}_1$ of $G_{\overline{\mathbb{Q}_p}}^{\text{ad}}$ is of classical Lie type and the highest weight of each non-trivial, simple $\text{Lie}(\mathcal{F}_1)$ -submodule of $L_A \otimes_{\mathbb{Z}} \overline{\mathbb{Q}_p}$, is a minuscule weight (cf. [Pi, Cor. 5.11]; see also [Ta2] for a slightly less general result).*

1.2. Conventions, notations, and preliminary organization. A reductive group H over a field K is assumed to be connected. Let $Z(H)$, H^{der} , H^{ad} , and H^{ab} denote the center, the derived group, the adjoint group, and the abelianization (respectively) of H . We have $Z^{\text{ab}}(H) = H/H^{\text{der}}$ and $H^{\text{ad}} = H/Z(H)$. Let $Z^0(H)$ be the maximal subtorus of $Z(H)$. If K is a finite, separable extension of another field $\tilde{K}$, let $\text{Res}_{K/\tilde{K}} H$ be the reductive group over $\tilde{K}$ obtained from H through the Weil restriction of scalars (see [BLR, Ch. 7, Subsect. 7.6] and [Va2, Subsect. 2.3]). For each commutative $\tilde{K}$ -algebra $\tilde{R}$, we have a functorial group identity $\text{Res}_{K/\tilde{K}} H(\tilde{R}) = H(\tilde{R} \otimes_{\tilde{K}} K)$. Let $\mathbb{S} := \text{Res}_{\mathbb{C}/\mathbb{R}} \mathbb{G}_{m, \mathbb{R}}$.

Always n is a natural number. If M and R and as before Subsection 1.1 and if ψ is a perfect alternating form on the free R -module M , then $\mathbf{Sp}(M, \psi)$ and $\mathbf{GSp}(M, \psi)$ are viewed as reductive group schemes over R .

A *Shimura pair* (H, X) consists of a reductive group H over $\mathbb{Q}$ and an $H(\mathbb{R})$ -conjugacy class X of homomorphisms $\mathbb{S} \rightarrow H_{\mathbb{R}}$ that satisfy Deligne’s axioms of [De2, Subsect. 2.1]: the Hodge $\mathbb{Q}$ -structure on $\text{Lie}(H)$ defined by any $x \in X$ is of type $\{(-1, 1), (0, 0), (1, -1)\}$, $\text{Ad} \circ x(i)$ is a Cartan involution of $\text{Lie}(H_{\mathbb{R}}^{\text{ad}})$, and no simple factor of H^{ad} becomes compact over $\mathbb{R}$. Here $\text{Ad} : H_{\mathbb{R}} \rightarrow \mathbf{GL}_{\text{Lie}(H_{\mathbb{R}}^{\text{ad}})}$ is the adjoint representation. These axioms imply that X has a natural structure of a hermitian symmetric domain, cf. [De2, Cor. 1.1.17]. For generalities on (the types of) the Shimura variety $\text{Sh}(H, X)$ defined by (H, X) we refer to [De1], [De2], [Mi2], [Mi3], and [Va1, Subsects. 2.2 to 2.5]. Shimura pairs of the form $(\mathbf{GSp}(W, \psi), S)$ define *Siegel modular varieties*. The adjoint (resp. the toric part) of a

Shimura pair (H, X) is denoted by $(H^{\text{ad}}, X^{\text{ad}})$ (resp. $(H^{\text{ab}}, X^{\text{ab}})$), cf. [Va1, Subsubsection 2.4.1].

We say (H, X) (or $\text{Sh}(H, X)$) is *unitary* if the group H^{ad} is non-trivial and each simple factor of $H_{\mathbb{Q}}^{\text{ad}}$ is of some A_n Lie type.

1.2.1. On techniques. In [Va1] we developed some techniques to study Shimura varieties and local deformations of abelian varieties endowed with de Rham tensors. With this paper we begin to exploit them to achieve progress in the proof of Conjecture 1.1.1. We mostly use techniques from reductive groups (see Sections 3, 6, and 7) and from Shimura varieties (see Sections 2, 4, 5, 6, and 7). See Subsection 1.3 for what the techniques of Sections 3 to 7 can achieve in connection to Conjecture 1.1.1.

For more details on how this work is structured see Subsections 1.5 and 1.6.

1.3. Our results on Conjecture 1.1.1. The group of homotheties of $L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p$ is a torus of $G_{\mathbb{Q}_p}$ (see [Bog]). The following theorem refines this result.

1.3.1. Theorem. *The two tori $Z^0(H_{A, \mathbb{Q}_p})$ and $Z^0(G_{\mathbb{Q}_p})$ of $\mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p}$ coincide i.e., we have $Z^0(H_{A, \mathbb{Q}_p}) = Z^0(G_{\mathbb{Q}_p})$.*

Let (H_A, X_A) be the Shimura pair attached to $A_{\mathbb{C}}$. Thus X_A is the $H_A(\mathbb{R})$ -conjugacy class of the monomorphism $h_A : \mathbb{S} \hookrightarrow H_{A, \mathbb{R}}$ that defines the classical Hodge $\mathbb{Z}$ -structure on L_A . We refer to $(H_A^{\text{ad}}, X_A^{\text{ad}})$ as the *adjoint Shimura pair attached to $A_{\mathbb{C}}$* and we write it

$$(H_A^{\text{ad}}, X_A^{\text{ad}}) = \prod_{j \in J} (H_j, X_j)$$

as a finite product of simple, adjoint Shimura pairs indexed by a set J which is assumed to be disjoint from $\mathbb{Z}$ (in order not to create confusion with homology groups).

Each H_j is a simple group over $\mathbb{Q}$. The simple factors of $H_{j, \overline{\mathbb{Q}}}$ have the same Lie type $\mathfrak{L}_j$ which is (cf. [Sa1], [Sa2], or [De2]) a classical Lie type. If $\mathfrak{L}_j$ is the A_n , B_n , or C_n Lie type, then (H_j, X_j) is said to be of A_n , B_n , or C_n type (respectively). If $n \geq 4$ and $\mathfrak{L}_j = D_n$, then (H_j, X_j) is of either $D_n^{\mathbb{H}}$ or $D_n^{\mathbb{R}}$ type (cf. [De2, Table 2.3.8]; see Definition 2.2.1 (a) for the $D_n^{\mathbb{H}}$ and $D_n^{\mathbb{R}}$ types). See Subsection 2.1 for the classical groups $\mathbf{SO}^*(2n)_{\mathbb{R}}$ and $\mathbf{SO}(2, 2n - 2)_{\mathbb{R}}$ over $\mathbb{R}$. If (H_j, X_j) is of $D_n^{\mathbb{H}}$ (resp. of $D_n^{\mathbb{R}}$) type, then all simple, non-compact factors of $H_{j, \mathbb{R}}$ are isomorphic to $\mathbf{SO}^*(2n)_{\mathbb{R}}^{\text{ad}}$ (resp. to $\mathbf{SO}(2, 2n - 2)_{\mathbb{R}}^{\text{ad}}$). The converse of the last sentence holds provided we have $n \geq 5$.

We say (H_j, X_j) is of *non-inner type* if H_j is isomorphic to $\text{Res}_{F_j/\mathbb{Q}} \tilde{H}_j$, where F_j is a number field and where $\tilde{H}_j$ is an absolutely simple, adjoint group over F_j which is not an inner form of a split group (see Definition 2.3 (a)).

1.3.2. Theorem. *We assume that $p \neq l$ and that v is unramified over l . There exists a number $N(A) \in \mathbb{N}$ that depends only on A (and not on p or l) and such that the following statements holds: if $l \geq N(A)$ and $j \in J$, then all images of $T_{v, \overline{\mathbb{Q}_p}}$ in simple factors of $H_{j, \overline{\mathbb{Q}_p}}$ have equal ranks that do not depend on the prime $p \neq l$.*

If the abelian variety A_v is ordinary, then Theorem 1.3.2 follows easily from [No, Thm. 2.2]. The following Corollary complements Theorem 1.1.5.

1.3.3. Corollary. *The rank of the subgroup of $H_{A, \mathbb{Q}_p}^{\text{ad}}$ whose extension to $\overline{\mathbb{Q}_p}$ is the product of the images of $G_{\overline{\mathbb{Q}_p}}$ in the simple factors of $H_{A, \mathbb{Q}_p}^{\text{ad}}$, does not depend on p .*

1.3.4. Main Theorem. *Conjecture 1.1.1 holds if for each element $j \in J$ the adjoint Shimura pair (H_j, X_j) is of one of the following types:*

- (a) *non-special A_n type;*
- (b) *B_n type;*
- (c) *C_n type, with n odd;*
- (d) *$D_n^{\mathbb{H}}$ type, with $n \geq 5$ odd and such that $2n \notin \{(2^{m+1}) \mid m \in \mathbb{N}\}$;*
- (d') *non-inner $D_{2n}^{\mathbb{H}}$ type, with n an odd prime;*
- (e) *$D_n^{\mathbb{R}}$ type, with $n \geq 4$.*

The types listed in Theorem 1.3.4 and the non-inner $D_4^{\mathbb{H}}$ type are the only types for which we can show that for each $j \in J$ the group $G_{\overline{\mathbb{Q}_p}}$ surjects onto a simple factor of $H_{j, \overline{\mathbb{Q}_p}}$ (see Theorem 7.1). The non-special A_n types are introduced in Definition 6.1 (c). Here we list some examples, cf. Examples 6.2.4.

1.3.5. Examples. The simple, adjoint Shimura pair (H_j, X_j) is of non-special A_n type if any one of the following four conditions is satisfied:

- (i) the group $H_{j, \mathbb{R}}$ has two simple factors $\mathbf{SU}(a_i, n+1-a_i)_{\mathbb{R}}^{\text{ad}}$, $i = \overline{1, 2}$, such that $1 \leq a_1 < a_2 \leq \frac{n+1}{2}$ and either $\text{g.c.d.}(a_1, n+1) = 1$ or $\text{g.c.d.}(a_2, n+1) = 1$;
- (ii) the group $H_{j, \mathbb{R}}$ has a simple factor $\mathbf{SU}(a, n+1-a)_{\mathbb{R}}^{\text{ad}}$, with $\text{g.c.d.}(a, n+1) = 1$ and with the pair $(a, n+1-a) \notin \{(\binom{r}{s}, \binom{r}{s-1}) \mid s, r \in \mathbb{N}, 2 \leq s \leq r-1\}$;
- (iii) the natural number $n+1$ is an integral power of 2 and the group $H_{j, \mathbb{R}}$ has as a normal factor a product $\mathbf{SU}(\frac{n+1}{2}, \frac{n+1}{2})_{\mathbb{R}}^{\text{ad}} \times_{\mathbb{R}} \mathbf{SU}(b, n+1-b)_{\mathbb{R}}^{\text{ad}}$, where $b \in \mathbb{N}$ is an odd number different from $\frac{n+1}{2}$;
- (iv) the Shimura pair (H_j, X_j) is of A_n type, where $n+1$ is either 4 or a prime.

1.3.6. Theorem. *We assume that there exists no element $j \in J$ such that (H_j, X_j) is of $D_4^{\mathbb{H}}$ type. Then no simple factor of $\text{Lie}(G_{\overline{\mathbb{Q}_p}}^{\text{ad}})$ surjects onto two or more distinct simple factors of $\text{Lie}(H_{A, \mathbb{Q}_p}^{\text{ad}})$.*

1.3.7. Theorem (the independence property for abelian varieties). *We assume that there exists no element $j \in J$ such that (H_j, X_j) is of $D_4^{\mathbb{H}}$ type and that A is a product of abelian varieties over E for which the Mumford–Tate conjecture holds. Then Conjecture 1.1.1 holds (i.e., the Mumford–Tate conjecture holds also for A).*

Theorems 1.3.1 and 1.3.2 are used in the proof of Theorem 1.3.4. Theorem 1.3.6 is a consequence of part of the proof of Theorem 1.3.4. Theorem 1.3.7 is an application of Theorem 1.3.6.

1.4. Earlier results that pertain to Conjecture 1.1.1. Previous works of Serre, Chi, Tankeev, Pink, Larsen, Zarhin, and few others proved Conjecture 1.1.1 in many cases.

The cases involve Shimura pairs (H_A, X_A) whose adjoints are simple of A_n , B_n , C_n , and $D_n^{\mathbb{H}}$ types (with n arbitrary large) and are worked out under the assumption that either the rank of $\text{End}(A_{\overline{\mathbb{Q}}})$ is small (like 1 or 4) or $\dim(A) \in \{qr | q \text{ is a prime, } r \in \{1, \dots, 9\}\}$ (see [Ch], [Ta1], [Ta2], [LP], [Pi, p. 216 and Thms. 5.14 and 5.15], and [Za2]). To our knowledge, except [Pi, Thm. 5.14] and [Za2], all previous results that pertain to Conjecture 1.1.1 are encompassed by Theorem 1.3.4. The result [Pi, Thm. 5.14] proves Conjecture 1.1.1 when $\text{End}(A_{\overline{\mathbb{Q}}}) = \mathbb{Z}$ and $2 \dim(A) \notin \{m^k, \binom{2k}{k} | m \in \mathbb{N}, k \in 1 + 2\mathbb{N}\}$. Results of [Za2] prove Conjecture 1.1.1 when $\text{End}(A_{\overline{\mathbb{Q}}}) = \mathbb{Z}$ and A is the Jacobian of certain hyperelliptic curves. In both places [Pi, Thm. 5.14] and [Za2], the group H_A is a $\mathbf{GSp}_{2 \dim(A), \mathbb{Q}}$ group and therefore the adjoint Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is simple of $C_{\dim(A)}$ type.

We recall from [Pi, Def. 4.1] that a pair $(\mathcal{G}_0, \rho_0)$ defined by an irreducible, faithful, finite dimensional representation $\rho_0 : \mathcal{G}_0 \hookrightarrow \mathbf{GL}_{\mathcal{W}_0}$ over $\overline{\mathbb{Q}_p}$, is called an *irreducible weak* (resp. an *irreducible strong*) *Mumford–Tate pair of weight $\{0, 1\}$* if $\mathcal{G}_0$ is a reductive group over $\overline{\mathbb{Q}_p}$ that is generated by cocharacters of it which act (resp. is generated by the $\mathcal{G}_0(\overline{\mathbb{Q}_p})$ -conjugates of one cocharacter of it which acts) on $\mathcal{W}_0$ via the trivial and the identity characters of $\mathbb{G}_{m, \overline{\mathbb{Q}_p}}$. We say that the pair $(\mathcal{G}_0, \rho_0)$ is non-trivial, if $\dim_{\overline{\mathbb{Q}_p}}(\mathcal{W}_0) \geq 2$.

Irreducible weak Mumford–Tate pairs of weight $\{0, 1\}$ are naturally attached to the p -adic Galois representation ρ of Subsection 1.1 as follows. We take $\mathcal{W}_0$ to be a simple $H_{A, \overline{\mathbb{Q}_p}}$ -submodule of $L_A \otimes_{\mathbb{Z}} \overline{\mathbb{Q}_p}$. Let $\mathcal{G}_0$ and $\mathcal{H}_0$ be the images of $G_{\overline{\mathbb{Q}_p}}$ and $H_{A, \overline{\mathbb{Q}_p}}$ (respectively) in $\mathbf{GL}_{\mathcal{W}_0}$. The natural faithful representation $\rho_0 : \mathcal{G}_0 \hookrightarrow \mathbf{GL}_{\mathcal{W}_0}$ is irreducible, cf. Theorem 1.1.4. Theorem 1.1.8 implies that the pair $(\mathcal{G}_0, \rho_0)$ is an irreducible weak Mumford–Tate pair of weight $\{0, 1\}$. From the very definition of H_A , one gets that the pair $(\mathcal{H}_0, \mathcal{H}_0 \hookrightarrow \mathbf{GL}_{\mathcal{W}_0})$ is also an irreducible weak Mumford–Tate pair of weight $\{0, 1\}$.

The previous works mentioned above rely on the classification of (non-trivial) irreducible weak and strong Mumford–Tate pairs of weight $\{0, 1\}$ over $\overline{\mathbb{Q}_p}$ (see [Pi]; see also [Sa2] and [Se1]) and on the results 1.1.2 to 1.1.9.

1.4.1. What the results 1.3.4 to 1.3.7 bring new. To our knowledge, Theorem 1.3.4 is the very first result on Conjecture 1.1.1 which is stated in terms of Shimura types (and thus which is stated independently on the structures of either $\text{End}(A_{\overline{\mathbb{Q}}})$ or H^{ab}). If H_A^{ad} is an absolutely simple, adjoint group, then only the $D_n^{\mathbb{R}}$ type case of Theorem 1.3.4 is completely new. The A_n and $D_n^{\mathbb{R}}$ types cases of Theorem 1.3.4 are new general instances for the validity of Conjecture 1.1.1. For instance, the $D_n^{\mathbb{R}}$ type case of Theorem 1.3.4 was not known before even when $\text{End}(A_{\overline{\mathbb{Q}}}) = \mathbb{Z}$. Also, the “applicability” of Theorem 1.3.4 (a) goes significantly beyond the numerical tests of Subsection 6.2 (see Remark 7.5.1 (a)). See Subsection 7.5 for the simplest cases not settled by Theorem 1.3.4 (they contain the case when $\dim(A) = 4$ and H_A is a $\mathbf{GSp}_{8, \mathbb{Q}}$ group). See Corollary 7.4 for a variant of Theorem 1.3.4 (d’) that handles to some extent the non-inner $D_4^{\mathbb{H}}$ types. Theorem 1.3.7 is a particular case of a more general result on abelian motives over E (see Theorem 7.6); we do not know any non-trivial analogue of Theorems 1.3.7 and 7.6 in the literature.

The proofs of the results 1.3.1 to 1.3.7 rely on:

- results of *arithmetic nature* (like the results 1.1.2 to 1.1.5, 1.1.8, and 1.1.9 and the main results of [Wi] and [Zi]);

- results of *group theoretical nature* (like the results 1.1.6 and 1.1.7, the classification of irreducible weak and strong Mumford–Tate pairs of weight $\{0, 1\}$, and the Satake–Deligne results on symplectic embeddings to be recalled in Subsection 1.5 below).

Our new ideas that put together all these results are presented below in Subsections 1.5 and 1.6. The main contribution of these new ideas is that they allow us: (i) to state Theorem 1.3.4 only in terms of Shimura types and (ii) to prove Theorems 1.3.6 and 1.3.7.

1.5. On injective maps into unitary Shimura pairs of PEL type. Let λ_A be a polarization of A . Let ψ_A be the non-degenerate alternating form on $W_A := L_A \otimes_{\mathbb{Z}} \mathbb{Q}$ induced by λ_A . We get naturally an injective map

$$f_A : (H_A, X_A) \hookrightarrow (\mathbf{GSp}(W_A, \psi_A), S_A)$$

of Shimura pairs, where S_A is the $\mathbf{GSp}(W_A, \psi_A)(\mathbb{R})$ -conjugacy class of $f_{A, \mathbb{R}} \circ h_A$. As a hermitian symmetric domain, S_A is two disjoint copies of the Siegel domain of genus $\dim(A)$. The inclusion $X_A \hookrightarrow S_A$ is a totally geodesic and holomorphic embedding between hermitian symmetric domains. Such embeddings were classified in [Sa1].

Satake and Deligne used rational variants of [Sa1] to show that each simple, adjoint Shimura pair of A_n , B_n , C_n , $D_n^{\mathbb{H}}$, or $D_n^{\mathbb{R}}$ type is isomorphic to the adjoint Shimura pair attached to some complex abelian variety (cf. [Sa2, Part III] and [De2, Prop. 2.3.10]). The goal of Section 4 is to present unitary variants of [De2, Subsubsects. 2.3.10 to 2.3.13]. Starting from a simple, adjoint Shimura pair (G, X) of A_n , B_n , C_n , $D_n^{\mathbb{H}}$, or $D_n^{\mathbb{R}}$ type, we construct explicitly in Subsections 4.1 to 4.8 “good” injective maps

$$f_2 : (G_2, X_2) \hookrightarrow (G_4, X_4) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$$

of Shimura pairs, where $(G_2^{\text{ad}}, X_2^{\text{ad}}) = (G, X)$, where $(G_4^{\text{ad}}, X_4^{\text{ad}})$ is a simple, adjoint Shimura pair of A_{n_4} type for some positive integer n_4 that depends only on the type of (G, X) , and where G_4 is the subgroup of $\mathbf{GSp}(W_1, \psi_1)$ that fixes a semisimple $\mathbb{Q}$ -subalgebra of $\text{End}(W_1)$. If (G, X) is of A_n type, then we can take $G_4 = G_2$ (cf. Proposition 4.1) and thus we have $n_4 = n$. If (G, X) is not of A_n type, see Theorem 4.8 for the extra “good” properties enjoyed by the injective maps f_2 of Shimura pairs.

These constructions play key roles in the proofs of the Theorems 1.3.2, 1.3.4, 1.3.6, and 1.3.7 (see Subsection 1.6 below). We also use these constructions to classify in Corollary 4.10 all Shimura pairs that are adjoints of Shimura pairs of PEL type (their definition is recalled in Subsection 2.4; for instance, the Shimura pair (G_4, X_4) is of PEL type).

The constructions rely on a modification of the proof of [De2, Prop. 2.3.10] (see proof of Proposition 4.1) and on a natural *twisting process* of certain homomorphisms between semisimple groups over number fields (see Subsection 4.5). The embeddings of hermitian symmetric domains of classical Lie type into Siegel domains constructed in [Sa1], are obtained by first achieving embeddings into hermitian symmetric domains associated to unitary groups over $\mathbb{R}$. This idea is not used in [De2] and thus the twisting process of Subsection 4.5 is the main new ingredient needed to get constructions that are explicit rational versions of [Sa1] and that are more practical than the ones of [Sa2].

1.6. On the proofs of the results 1.3.1 to 1.3.7. The proof of Theorem 1.3.1 relies on an elementary Lemma 3.3 and appeals to canonical split cocharacters of filtered F -isocrystals over finite fields defined in [Wi, p. 512] (see Subsection 3.4). Lemma 3.3 is due in essence to Bogomolov and Serre, cf. [Bog] and [Se3, pp. 31, 43, and 44]. Though Theorem 1.3.1 was known to Serre (and to others), we could not trace it in the literature.

We call an abelian variety B over E *adjoint-isogenous* to A if the adjoints of the Mumford–Tate groups of $A_{\mathbb{C}}$, $B_{\mathbb{C}}$, and $A_{\mathbb{C}} \times_{\mathbb{C}} B_{\mathbb{C}}$ are the same (cf. Definition 5.1). Due to Theorems 1.3.1, 1.1.2, 1.1.3, and 1.1.4, to prove Conjecture 1.1.1 one only has to show that $G_{\mathbb{Q}_p}^{\text{ad}} = H_{A, \mathbb{Q}_p}^{\text{ad}}$. But to check that we have $G_{\mathbb{Q}_p}^{\text{ad}} = H_{A, \mathbb{Q}_p}^{\text{ad}}$, we can replace A by any other abelian variety B over E that is adjoint-isogenous to A (cf. the shifting process expressed by Proposition 5.4.1). In other words, (as the prime p is arbitrary) we have:

1.6.1. Proposition. *We assume that A and B are adjoint-isogenous abelian varieties over a number field. Then the Mumford–Tate conjecture holds for A if and only if it holds for B .*

1.6.2. Example. We assume that the adjoint group H_A^{ad} is absolutely simple and that the simple, adjoint Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of $D_n^{\mathbb{R}}$ type with $n \geq 5$. Let H_A^{sc} be the simply connected semisimple group cover of H_A^{ad} (or of H_A^{der}). Let K be a number field such that the group $H_{A, K}^{\text{sc}}$ is split (i.e., it is a $\mathbf{Spin}_{2n, K}$ group). Let $H_{A, K}^{\text{sc}} \hookrightarrow \mathbf{GL}_{2^n, K}$ be the spin representation (i.e., the direct sum of the two half spin representations of $H_{A, K}^{\text{sc}}$). We have a natural sequence of monomorphisms

$$H_A^{\text{sc}} \hookrightarrow \text{Res}_{K/\mathbb{Q}} H_{A, K}^{\text{sc}} \hookrightarrow \text{Res}_{K/\mathbb{Q}} \mathbf{GL}_{2^n, K} \hookrightarrow \mathbf{GL}_{2^n[K:\mathbb{Q}], \mathbb{Q}} \hookrightarrow \mathbf{GSp}_{2^{n+1}[K:\mathbb{Q}], \mathbb{Q}}$$

between reductive groups over $\mathbb{Q}$. We write $\mathbf{GSp}_{2^{n+1}[K:\mathbb{Q}], \mathbb{Q}} = \mathbf{GSp}(W_B, \tilde{\psi}_B)$, where W_B is a rational vector space of dimension $2^{n+1}[K:\mathbb{Q}]$ and where $\tilde{\psi}_B$ is a non-degenerate alternating form on W_B . Let H_B be the reductive subgroup of $\mathbf{GL}_{W_B}$ generated by the subgroups H_A^{sc} and $Z(\mathbf{GSp}(W_B, \psi_B))$; we have natural identifications $H_B^{\text{der}} = H_A^{\text{sc}}$ and $H_B^{\text{ad}} = H_A^{\text{ad}}$. The centralizer of H_B in $\mathbf{GL}_{W_B \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_p}}$ is isomorphic to $\mathbf{GL}_{[K:\mathbb{Q}], \overline{\mathbb{Q}_p}}^2$ (each copy of $\mathbf{GL}_{[K:\mathbb{Q}], \overline{\mathbb{Q}_p}}$ corresponds to one of the two half spin representations).

Let $h_A^{\text{ad}} = h_B^{\text{ad}} : \mathbb{S} \rightarrow H_{A, \mathbb{R}}^{\text{ad}} = H_{B, \mathbb{R}}^{\text{ad}}$ be the composite of the monomorphism $h_A : \mathbb{S} \hookrightarrow H_{A, \mathbb{R}}$ with the natural epimorphism $H_{A, \mathbb{R}} \twoheadrightarrow H_{A, \mathbb{R}}^{\text{ad}}$. It is easy to see that there exists a unique monomorphism $h_B : \mathbb{S} \hookrightarrow H_{B, \mathbb{R}}$ which lifts h_B^{ad} and which defines a Hodge $\mathbb{Q}$ -structure on W_B of type $\{(-1, 0), (0, -1)\}$, cf. either the proof of [De2, Prop. 2.3.10] or [Pi, Table 4.2]. Let ψ_B be a non-degenerate alternating form on W_B that is fixed by $H_B^{\text{der}} = H_A^{\text{sc}}$ and such that $2\pi i \psi_B$ is a polarization of this Hodge $\mathbb{Q}$ -structure on W_B , cf. [De2, Cor. 2.3.3]. Let X_B be the $H_B(\mathbb{R})$ -conjugacy class of h_B and let S_B be the $\mathbf{GSp}(W_B, \psi_B)(\mathbb{R})$ -conjugacy class of the composite of h_B with the monomorphism $H_{B, \mathbb{R}} \hookrightarrow \mathbf{GSp}(W_B \otimes_{\mathbb{Q}} \mathbb{R}, \psi_B)$. By replacing E with a finite field extension of it, there exists a principally polarized abelian variety (B, λ_B) over E such that the notations match (i.e., the analogues of h_A and f_A for (B, λ_B) are precisely h_B and $f_B : (H_B, X_B) \hookrightarrow (\mathbf{GSp}(W_B, \psi_B), S_B)$) and such that the abelian varieties A and B are adjoint-isogenous.

Let $\tilde{G}_{\mathbb{Q}_p}$ be the identity component of the algebraic envelope of the p -adic Galois representation attached to B ; it is a subgroup of $H_{B, \mathbb{Q}_p}$. We show that the assumption

that $\tilde{G}_{\mathbb{Q}_p} \neq H_{B, \mathbb{Q}_p}$ leads to a contradiction. Due to Theorem 1.1.8, this assumption implies that $\tilde{G}_{\mathbb{Q}_p}^{\text{der}}$ is isogenous to a product of two simple, adjoint groups of B_{n_1} and B_{n-n_1-1} Lie type (respectively), where $n_1 \in \{0, \dots, n-1\}$ (cf. [Pi, Prop. 4.3]; here the B_0 Lie type corresponds to the trivial group). Moreover, loc. cit. implies that the representation $\text{Lie}(\tilde{G}_{\mathbb{Q}_p}^{\text{der}}) \hookrightarrow \text{End}(W_B \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_p})$ is a direct sum of $2[K : \mathbb{Q}]$ irreducible representations that are isomorphic to the tensor product of the spin representations of the two simple factors of $\text{Lie}(\tilde{G}_{\mathbb{Q}_p}^{\text{der}})$ of B_{n_1} and B_{n-n_1-1} Lie type (in other words, the restriction to $\tilde{G}_{\mathbb{Q}_p}^{\text{der}}$ of either one of the two half spin representations of $H_{B, \mathbb{Q}_p}$ is the same irreducible representation of $\tilde{G}_{\mathbb{Q}_p}^{\text{der}}$).

Thus the centralizer of $\tilde{G}_{\mathbb{Q}_p}$ in $\mathbf{GL}_{W_B \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_p}}$ is isomorphic to $\mathbf{GL}_{2[K : \mathbb{Q}], \overline{\mathbb{Q}_p}}$ and therefore it is distinct from the centralizer of $H_{B, \mathbb{Q}_p}$ in $\mathbf{GL}_{W_B \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_p}}$. This contradicts Theorem 1.1.4.

Therefore we have $\tilde{G}_{\mathbb{Q}_p} = H_{B, \mathbb{Q}_p}$. Thus the Mumford–Tate conjecture holds for B (cf. Theorem 1.1.6) and thus also for A (cf. Proposition 1.6.1). This holds even if $n = 4$, provided we define the two half spin representations in such away that h_B does exist. If either n is odd or n is even and the simple, adjoint Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of non-inner $D_n^{\mathbb{R}}$ type, then the representation $H_{A, \mathbb{Q}_p}^{\text{der}} \hookrightarrow \mathbf{GL}_{W_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_p}}$ is a direct sum of spin representations and therefore the above replacement of A by B is not truly necessary.

1.6.3. The role of the shifting process. The main role of the shifting process (i.e., of Proposition 5.4.1) is to replace the “potentially complicated” injective map f_A of Shimura pairs by a “simpler” one that is a “direct sum” (i.e., is a Hodge quasi product in the sense of Subsection 2.4) of composite injective maps $f_2 : (G_2, X_2) \hookrightarrow (G_4, X_4) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ of Shimura pairs that are as in Subsection 1.5. See Proposition 5.10 for three advantages one gets through such replacements; they are due to Deligne. We emphasize that:

(i) If B is an abelian variety over E which is adjoint-isogenous to A , then in general B is unrelated to A in any usual way (like via isogenies, Weil restrictions of scalars, products of abelian subvarieties of A , etc.). For instance, B could be the product between A and any other abelian variety over E which has complex multiplication over $\overline{E}$.

(ii) To perform the replacement of f_A by a “direct sum” of composite injective maps f_2 of Shimura pairs, we often have to pass to a finite field extension of E .

(iii) If the adjoint group H_A^{ad} is $\mathbb{Q}$ -simple, then by replacing f_A with an f_2 , the Mumford–Tate group H_A gets replaced by a reductive, normal subgroup of G_2 which contains G_2^{der} (more precisely, the injective map $f_A : (H_A, X_A) \hookrightarrow (\mathbf{GSp}(W_A, \psi_A), S_A)$ of Shimura pairs gets replaced by a composite injective map

$$(H_A, X_A) \hookrightarrow (G_2, X_2) \hookrightarrow (G_4, X_4) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1) = (\mathbf{GSp}(W_A, \psi_A), S_A)$$

of Shimura pairs for which we have $H_A^{\text{der}} = G_2^{\text{der}}$ and which is defined by an f_2 as in Subsection 1.5).

1.6.4. On Theorem 1.3.2. The proof of Theorem 1.3.2 goes as follows (see Subsection 5.8). Using the shifting process, we can assume that H_A^{ad} is a simple $\mathbb{Q}$ -group and we can replace f_A by an injective map f_2 of Shimura pairs as in Subsection 1.5. Using this and

the main result of [Zi], we get that for $l \gg 0$ and up to a $G_4(\mathbb{Q}_p)$ -conjugation of the torus $T_{v, \mathbb{Q}_p}$, T_v is naturally isomorphic to a torus T'_4 of an inner form G'_4 of G_4 that is locally isomorphic. But from the constructions of f_2 we get that each simple factor of $\text{Lie}(G_{4, \mathbb{Q}_p}^{\text{ad}})$ contains the non-zero image of a unique simple factor of $\text{Lie}(G_{2, \mathbb{Q}_p}^{\text{ad}})$ (see property 5.7 (a)). Thus we get Theorem 1.3.2 by considering the ranks of the images of $T'_{4, \mathbb{Q}_p}$ in simple factors of $G_{4, \mathbb{Q}_p}^{\text{ad}} = G_{4, \mathbb{Q}_p}'^{\text{ad}}$. The passage from Theorem 1.3.2 to Corollary 1.3.3 is standard (see Subsection 5.8).

1.6.5. On Theorems 1.3.4, 1.3.6, and 1.3.7. The proof of Theorem 1.3.4 involves two main steps (see Theorems 7.1 and 7.2) and it ends in Subsection 7.3. Theorem 7.1 shows that if for each element $j \in J$ the simple, adjoint Shimura pair (H_j, X_j) is of either non-inner $D_4^{\mathbb{H}}$ type or a type listed in Theorem 1.3.4, then for each $j \in J$ the group $G_{\mathbb{Q}_p}$ surjects onto a simple factor of $H_{j, \mathbb{Q}_p}$. The greatest part of the proof of Theorem 7.1 is a standard application of the classification of irreducible weak and strong Mumford–Tate pairs of weight $\{0, 1\}$ and of the advantages offered by the shifting process.

Theorem 7.2 shows that under some hypotheses (like the ones of Theorem 1.3.4), we have $G_{\mathbb{Q}_p}^{\text{ad}} = H_{A, \mathbb{Q}_p}^{\text{ad}}$. Its proof is technical and lengthy. To explain the main point of the proofs of Theorems 7.2, 1.3.6, and 1.3.7, we will assume in this and the next paragraph that there exists no element $j \in J$ such that (H_j, X_j) is of $D_4^{\mathbb{H}}$ type and that (based on the shifting process) the injective map f_A has been replaced by a “direct sum” of injective map f_2 as in Subsection 1.5. Even more, to ease the notations we will also assume that the adjoint group H_A^{ad} is $\mathbb{Q}$ -simple and thus that f_A has been replaced (as in (iii)) by a single injective map f_2 . The torus $Z^0(H_A)$ is a subtorus of $Z^0(G_2)$ (cf. (iii)) and its dimension is very much controlled by $G_{\mathbb{Q}_p}$. If there exists a simple factor of $\text{Lie}(G_{\mathbb{Q}_p}^{\text{ad}})$ that surjects onto two or more distinct simple factors of $\text{Lie}(H_{A, \mathbb{Q}_p}^{\text{ad}})$, then directly from the construction of the injective map f_2 of Shimura pairs we get that the centralizers of $G_{\mathbb{Q}_p}^{\text{der}}$ and $H_{A, \mathbb{Q}_p}^{\text{der}}$ in $\mathbf{GL}_{T_p(A) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p}$ are distinct and thus one hopes to reach a contradiction with Theorem 1.1.4. But in order to reach the desired contradiction, one has to check that the subtorus $Z^0(H_A)$ of $Z^0(G_2)$ is small enough (so that the centralizers of $G_{\mathbb{Q}_p}$ and $H_{A, \mathbb{Q}_p}$ in $\mathbf{GL}_{T_p(A) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p}$ are distinct). Thus the main philosophy of the proof of Theorem 7.2 is:

(iv) the smaller the subgroup $G_{\mathbb{Q}_p}$ of $H_{A, \mathbb{Q}_p} \leq G_{2, \mathbb{Q}_p}$ is known to be (in terms of several simple factors of $\text{Lie}(G_{\mathbb{Q}_p}^{\text{ad}})$ surjecting onto two or more distinct simple factors of $\text{Lie}(H_{A, \mathbb{Q}_p}^{\text{ad}})$), the smaller subtori of $Z^0(G_2)$ we get which contain $Z^0(H_A)$ (and thus the better upper bounds we get for $\dim(Z^0(H_A))$).

Thus if there exists a simple factor of $\text{Lie}(G_{\mathbb{Q}_p}^{\text{ad}})$ that surjects onto two or more distinct simple factors of $\text{Lie}(H_{A, \mathbb{Q}_p}^{\text{ad}})$, then the proof of Theorem 7.2 shows that the torus $Z^0(H_A)$ is small enough so that we reach a contradiction with Theorem 1.1.4. In the proof of Theorem 7.2 we will use very intimately the construction of the injective map f_2 of Shimura pairs and the proof of Theorem 1.3.2, in order to get the desired control on the subtorus $Z^0(H_A)$ of $Z^0(G_2)$.

The essence of the proofs of Theorems 1.3.6 and 1.3.7 is the same main point mentioned above. The proof of Theorem 1.3.6 is presented in Subsection 7.3. Theorem 1.3.7 is a particular case of Theorem 7.6.

Acknowledgments. We would like to thank UC at Berkeley, U of Utah, U of Arizona, and Binghamton U for providing us with good conditions with which to write this work. We would like to thank Tankeev for making us aware of Theorem 1.1.6, Serre and Deligne for some comments, Borovoi for mentioning the reference [Sa2], and the referee and Larsen for many valuable comments and suggestions. We are obliged to [Pi]. This research was partially supported by the NSF grant DMS 97-05376.

2. Some complements on Shimura pairs

In Subsection 2.1 we gather extra conventions and notations to be used in Sections 2 to 6. In Subsections 2.2 and 2.3 we include complements on reflex fields and types of Shimura pairs. In Subsection 2.4 we include complements on injective maps into Shimura pairs that define Siegel modular varieties. The notations of Section 2 will be independent from the ones of Subsections 1.1, 1.3, and 1.5.

2.1. Extra conventions and notations. Let $p \in \mathbb{N}$ be a prime. For each field K and every cocharacter μ_0 of a reductive group H over K , let $[\mu_0]$ be the $H(K)$ -conjugacy class of μ_0 . A pair of the form $(H, [\mu_1])$, with μ_1 a cocharacter of $H_{\overline{K}}$, is called a *group pair* over K . If D is a semisimple K -algebra, let $\mathbf{GL}_1(D)$ be the reductive group over K of invertible elements of D . If K is a p -adic field, we call H *unramified* if it is quasi-split and splits over an unramified extension of K . If K is a number field, let K_w be the completion of K with respect to a finite prime w of K . Let $k(w)$ be the residue field of w . Let $B(k)$ be the field of fractions of the ring $W(k)$ of Witt vectors with coefficients in a perfect field k . Let $\mathbb{F} := \overline{\mathbb{F}_p}$. Let $\mathbb{A}_f := \widehat{\mathbb{Z}} \otimes_{\mathbb{Z}} \mathbb{Q}$ be the $\mathbb{Q}$ -algebra of finite adèles of $\mathbb{Q}$. If an abstract group C acts on a set V , let $V^C := \{y \in V | y \text{ fixed by } C\}$.

Let (G, X) be a Shimura pair. Let $E(G, X)$ be the reflex field of (G, X) . The Shimura variety $\mathrm{Sh}(G, X)$ is identified with the canonical model over $E(G, X)$ of the complex Shimura variety $\mathrm{Sh}(G, X)_{\mathbb{C}}$ (see [De1], [De2], [Mi1], [Mi3], and [Mi4]).

Let $a, b \in \mathbb{N} \cup \{0\}$ with $a + b > 0$. Let $\mathbf{SU}(a, b)$ be the simply connected semisimple group over $\mathbb{Q}$ whose $\mathbb{Q}$ -valued points are the $\mathbb{Q}(i)$ -valued points of $\mathbf{SL}_{a+b, \mathbb{Q}}$ that leave invariant the hermitian form $-z_1 \bar{z}_1 - \cdots - z_a \bar{z}_a + z_{a+1} \bar{z}_{a+1} + \cdots + z_{a+b} \bar{z}_{a+b}$ over $\mathbb{Q}(i)$. Let $\mathbf{SO}(a, b)$ be the semisimple group over $\mathbb{Q}$ of $a + b$ by $a + b$ matrices of determinant 1 that leave invariant the quadratic form $-x_1^2 - \cdots - x_a^2 + x_{a+1}^2 + \cdots + x_{a+b}^2$ on $\mathbb{Q}^{a+b}$. Let $\mathbf{SO}(a) := \mathbf{SO}(0, a)$. Let $\mathbf{SO}^*(2a)$ be the semisimple group over $\mathbb{Q}$ whose group of $\mathbb{Q}$ -valued points is the subgroup of $\mathbf{SO}(2n)(\mathbb{Q}(i))$ that leaves invariant the skew hermitian form $-z_1 \bar{z}_{n+1} + z_{n+1} \bar{z}_1 - \cdots - z_n \bar{z}_{2n} + z_{2n} \bar{z}_n$ over $\mathbb{Q}(i)$ (z_i 's and x_i 's are related here over $\mathbb{Q}(i)$ via $z_i = x_i$).

The *minuscule weights* of a split, simple Lie algebra of classical Lie type $\mathfrak{L}$ over a field of characteristic 0 are: ϖ_i with $i \in \{1, \dots, n\}$ if $\mathfrak{L} = A_n$, ϖ_n if $\mathfrak{L} = B_n$, ϖ_1 if $\mathfrak{L} = C_n$, and ϖ_1, ϖ_{n-1} , and ϖ_n if $\mathfrak{L} = D_n$ with $n \geq 3$ (see [Sa1], [Bou2, pp. 127–129], [De2], [Se1]).

2.2. On reflex fields and Shimura types. Let (G, X) be a Shimura pair. We have $\mathbb{S}(\mathbb{R}) = \mathbb{G}_{m, \mathbb{R}}(\mathbb{C})$. We identify $\mathbb{S}(\mathbb{C}) = \mathbb{G}_{m, \mathbb{R}}(\mathbb{C}) \times \mathbb{G}_{m, \mathbb{R}}(\mathbb{C})$ in such a way that the monomorphism $\mathbb{S}(\mathbb{R}) \hookrightarrow \mathbb{S}(\mathbb{C})$ induces the map $z \rightarrow (z, \bar{z})$ (here $z \in \mathbb{G}_{m, \mathbb{R}}(\mathbb{C})$). Let $x \in X$; it is a homomorphism $x : \mathbb{S} \rightarrow G_{\mathbb{R}}$. Let $\mu_x : \mathbb{G}_{m, \mathbb{C}} \rightarrow G_{\mathbb{C}}$ be the cocharacter given on complex points by the rule $z \rightarrow x_{\mathbb{C}}(z, 1)$. Let μ_0 be a cocharacter of $G_{\overline{\mathbb{Q}}}$ such that we have $[\mu_{0, \mathbb{C}}] = [\mu_x]$. If K is a field of characteristic 0, the $G(\overline{K})$ -conjugacy class $[\mu_{0, \overline{K}}]$ of $\mu_{0, \overline{K}}$ will be denoted simply by μ_K . If K is a p -adic field, let $[\mu_p] := [\mu_K] = [\mu_{0, \overline{K}}]$.

Let $\mathfrak{C}(G, X)$ be the set of cocharacters of $G_{\mathbb{C}}$ that are $G(\mathbb{C})$ -conjugates of the extensions to $\mathbb{C}$ of all Galois conjugates of μ_0 (i.e., of all cocharacters of the form $\gamma(\mu_0) : \mathbb{G}_{m, \overline{\mathbb{Q}}} \rightarrow G_{\overline{\mathbb{Q}}}$ for some element $\gamma \in \text{Gal}(\overline{\mathbb{Q}})$). As the notation suggests, the set $\mathfrak{C}(G, X)$ does not depend on the choice of the cocharacter $\mu_0 \in [\mu_{\mathbb{Q}}] = [\mu_0]$.

The *reflex field* $E(G, X)$ of (G, X) is the field of definition of $[\mu_{\mathbb{C}}] = [\mu_x]$ (see [De1], [De2], [Mi2], and [Mi3]). It is the composite field of $E(G^{\text{ad}}, X^{\text{ad}})$ and $E(G^{\text{ab}}, X^{\text{ab}})$, cf. [De1, Prop. 3.8 (i)]. Thus, we now recall how one computes $E(G, X)$, if G is either a torus or a simple, adjoint group. If G is a torus, then $X = \{x\}$ and $E(G, X)$ is the field of definition of μ_x itself. From now on until Subsection 2.4 we will assume that G is a simple, adjoint group over $\mathbb{Q}$; thus (G, X) is a simple, adjoint Shimura pair.

Let $F(G, X)$ be a totally real number subfield of $\overline{\mathbb{Q}} \subseteq \mathbb{C}$ such that we have

$$G = \text{Res}_{F(G, X)/\mathbb{Q}} G[F(G, X)],$$

with $G[F(G, X)]$ as an absolutely simple, adjoint group over $F(G, X)$ (cf. [De2, Subsubsection. 2.3.4]); the subfield $F(G, X)$ of $\overline{\mathbb{Q}}$ is unique up to $\text{Gal}(\overline{\mathbb{Q}})$ -conjugation. Let I_0 be the split, simple, adjoint group over $\mathbb{Q}$ of the same Lie type as every simple factor of $G_{\overline{\mathbb{Q}}}$. Let $\text{Aut}(I_0)$ be the group over $\mathbb{Q}$ of automorphisms of I_0 . The group I_0 is the group of inner automorphisms of I_0 and thus it is a normal subgroup of $\text{Aut}(I_0)$. The quotient group $\text{Aut}(I_0)/I_0$ is either trivial or $\mathbb{Z}/2\mathbb{Z}$ or the symmetric group S_3 . Let $I(G, X)$ be the smallest field extension of $F(G, X)$ such that the group $G[F(G, X)]_{I(G, X)}$ is an inner form of $I_{0, I(G, X)}$; the degree $[I(G, X) : F(G, X)]$ divides the order of the finite group $\text{Aut}(I_0)/I_0$.

Let T be a maximal torus of G . Let B be a Borel subgroup of $G_{\overline{\mathbb{Q}}}$ that contains $T_{\overline{\mathbb{Q}}}$. Let $\mathfrak{D}$ be the Dynkin diagram of $\text{Lie}(G_{\overline{\mathbb{Q}}})$ with respect to $T_{\overline{\mathbb{Q}}}$ and B . As $G_{\overline{\mathbb{Q}}} = \prod_{i \in \text{Hom}(F(G, X), \mathbb{R})} G[F(G, X)] \times_{F(G, X), i} \overline{\mathbb{Q}}$, we have a disjoint union

$$\mathfrak{D} = \cup_{i \in \text{Hom}(F(G, X), \mathbb{R})} \mathfrak{D}_i,$$

where $\mathfrak{D}_i$ is the Dynkin diagram of $\text{Lie}(G[F(G, X)] \times_{F(G, X), i} \overline{\mathbb{Q}})$. For a vertex $\mathfrak{n}$ of $\mathfrak{D}$, let $\mathfrak{g}_{\mathfrak{n}}$ be the 1 dimensional Lie subalgebra of $\text{Lie}(B)$ that corresponds to $\mathfrak{n}$.

The Galois group $\text{Gal}(\overline{\mathbb{Q}})$ acts on $\mathfrak{D}$ as follows. If $\gamma \in \text{Gal}(\overline{\mathbb{Q}})$, then $\gamma(\mathfrak{n})$ is the vertex of $\mathfrak{D}$ defined by the identity $\mathfrak{g}_{\gamma(\mathfrak{n})} = i_{g_{\gamma}}(\gamma(\mathfrak{g}_{\mathfrak{n}}))$, where $i_{g_{\gamma}}$ is the inner conjugation of $\text{Lie}(G_{\overline{\mathbb{Q}}})$ defined by an element $g_{\gamma} \in G(\overline{\mathbb{Q}})$ that normalizes $T_{\overline{\mathbb{Q}}}$ and that has the property that $g_{\gamma}\gamma(B)g_{\gamma}^{-1} = B$. Here $\gamma(B)$ is the Galois conjugate of B through γ .

Let $\mathfrak{n}_X$ be the set of vertices of $\mathfrak{D}$ such that the cocharacter of $T_{\overline{\mathbb{Q}}}$ that acts on $\mathfrak{g}_{\mathfrak{n}}$ trivially if $\mathfrak{n} \notin \mathfrak{n}_X$ and via the identical character of $\mathbb{G}_{m, \overline{\mathbb{Q}}}$ if $\mathfrak{n} \in \mathfrak{n}_X$, belongs to $[\mu_0]$. The

image of $x : \mathbb{S} \rightarrow G_{\mathbb{R}}$ in a simple factor $\mathcal{F}$ of $G_{\mathbb{R}}$ is trivial if and only if the group $\mathcal{F}$ is compact (this is so as the centralizer of $\text{Im}(x)$ in $G_{\mathbb{R}}$ is a maximal compact subgroup of $G_{\mathbb{R}}$, cf. [De2, p. 259]). Thus, as the group $G_{\mathbb{R}}$ is not compact, the set $\mathfrak{n}_X$ is non-empty.

As the set $\mathfrak{n}_X$ is non-empty and as the Hodge $\mathbb{Q}$ -structure on $\text{Lie}(G)$ defined by each $x \in X$ is of type $\{(-1, 1), (0, 0), (1, -1)\}$, the group I_0 is of either classical or E_6 or E_7 Lie type (see [De2, Table 1.3.9]). Moreover, if $\mathfrak{n}_X$ contains a vertex of $\mathfrak{D}_i$, then (with the standard notations of [Bou1, Planches I to VI]) this vertex is: an arbitrary vertex if I_0 is of A_n Lie type, vertex 1 if I_0 is of B_n Lie type, vertex n if I_0 is of C_n Lie type, an extremal vertex if I_0 is of D_n Lie type, vertex 1 or 6 if I_0 is of E_6 Lie type, and vertex 7 if I_0 is of E_7 Lie type (cf. [De2, Table 1.3.9]). The reflex field $E(G, X)$ is the fixed field of the open subgroup of $\text{Gal}(\mathbb{Q})$ that is the stabilizer of $\mathfrak{n}_X$, cf. [De2, Prop. 2.3.6].

The following three definitions conform with [De2] and [Mi3].

2.2.1. Definitions. (a) If I_0 is of A_n , B_n , C_n , E_6 , or E_7 Lie type, then we say (G, X) is of A_n , B_n , C_n , E_6 , or E_7 type (respectively).

Let now I_0 be of D_n Lie type, with $n \geq 4$. If $n = 4$, we say (G, X) is of $D_4^{\mathbb{R}}$ (resp. of $D_4^{\mathbb{H}}$) type if for each embedding $i : F(G, X) \hookrightarrow \mathbb{R}$, the orbit of $\mathfrak{n}_X$ under $\text{Gal}(\mathbb{Q})$ contains only one vertex (resp. only two vertices) of $\mathfrak{D}_i$. If $n \geq 5$, we say (G, X) is of $D_n^{\mathbb{R}}$ type if for each embedding $i : F(G, X) \hookrightarrow \mathbb{R}$, the orbit of $\mathfrak{n}_X$ under $\text{Gal}(\mathbb{Q})$ contains only the vertex 1 of $\mathfrak{D}_i$. If $n \geq 5$, we say (G, X) is of $D_n^{\mathbb{H}}$ type if for each embedding $i : F(G, X) \hookrightarrow \mathbb{R}$, the orbit of $\mathfrak{n}_X$ under $\text{Gal}(\mathbb{Q})$ contains only vertices of $\mathfrak{D}_i$ that belong to the set $\{n-1, n\}$. If $n \geq 4$, we say (G, X) is of D_n^{mixed} type if it is neither of $D_n^{\mathbb{H}}$ type nor of $D_n^{\mathbb{R}}$ type.

(b) An adjoint Shimura pair is said to be of *abelian type* if it is a product of simple, adjoint Shimura pairs of A_n , B_n , C_n , $D_n^{\mathbb{H}}$, or $D_n^{\mathbb{R}}$ type.

(c) Let (G, X) be of $D_n^{\mathbb{R}}$ type, $n \geq 4$. If $n = 4$ we index the vertices of each $\mathfrak{D}_i$ in such a way that the $\text{Gal}(\mathbb{Q})$ -orbit of $\mathfrak{n}_X$ contains only the vertex 1 of $\mathfrak{D}_i$. Let $*$ be a field of characteristic 0 such that the group G_* is split. By the two half spin representations of a simple factor of $\text{Lie}(G_*)$, we mean the two representations associated to the highest weights ϖ_{n-1} and ϖ_n . By the spin representation of a simple factor of $\text{Lie}(G_*)$, we mean the direct sum of the two half spin representations of the mentioned simple factor. Similarly, we speak about the two half spin representations and about the spin representation of the simply connected, semisimple group cover of a simple factor of G_* .

2.2.2. Extra fields. There exists a unique totally real number field $E_1(G, X)$ such that $E(G, X)$ is either it or a totally imaginary quadratic extension of it, cf. [De1, Prop. 3.8 (ii)]. We say (G, X) is: (i) *without involution* if $E(G, X) = E_1(G, X)$, and (ii) *with involution* if $E(G, X) \neq E_1(G, X)$. We have $E(G, X) = E_1(G, X)$ if and only if the set $\mathfrak{n}_X$ is fixed by the automorphism of $\overline{\mathbb{Q}}$ defined by the complex conjugation, cf. loc. cit. The last automorphism of $\overline{\mathbb{Q}}$ acts on each $\mathfrak{D}_i$ via the standard opposition (it is either trivial or an involution) defined by the element $-w_0$ of [Bou1, item (XI), Planches I to IV], cf. [De1, Prop. 3.8 (ii)]. Thus, if (G, X) is not of some A_{2n+1} type, then from the description of elements w_0 we get that (G, X) is with involution if and only if it is of one of the following types: A_{2n} , $D_{2n+3}^{\mathbb{H}}$, D_{2n+3}^{mixed} , or E_6 . If (G, X) is of A_{2n+1} type, then it is without involution if and only if all simple, non-compact factors of $G_{\mathbb{R}}$ are $\text{SU}(n+1, n+1)_{\mathbb{R}}^{\text{ad}}$ groups (i.e., if and only if for each element $i \in \text{Hom}(F(G, X), \mathbb{R})$, the set $\mathfrak{n}_X$ contains either no vertex of

$\mathfrak{D}_i$ or only the vertex $n + 1$).

Let $EF_1(G, X)$ (resp. $EF(G, X)$) be the composite field of $F(G, X)$ with $E_1(G, X)$ (resp. with $E(G, X)$) inside $\overline{\mathbb{Q}}$. Let $G_1(G, X)$ (resp. $G(G, X)$) be the Galois extension of $\mathbb{Q}$ generated by $EF_1(G, X)$ (resp. by $EF(G, X)$). Let $G_2(G, X)$ be the Galois extension of $\mathbb{Q}$ generated by $F(G, X)$. Both number fields $G_1(G, X)$ and $G_2(G, X)$ are totally real.

2.2.3. A construction. For a totally real number field F_1 that contains $F(G, X)$ let

$$G^{F_1} := \text{Res}_{F_1/\mathbb{Q}} G[F(G, X)]_{F_1}.$$

We have a natural monomorphism $G[F(G, X)] \hookrightarrow \text{Res}_{F_1/F(G, X)} G[F(G, X)]_{F_1}$ over $F(G, X)$ which at the level of $F(G, X)$ -valued points is defined by the inclusion

$$GF(G, X) \hookrightarrow G[F(G, X)](F_1) = \text{Res}_{F_1/F(G, X)} G[F(G, X)]_{F_1}(F(G, X)).$$

The $\text{Res}_{F(G, X)/\mathbb{Q}}$ of this natural monomorphism is a monomorphism $G \hookrightarrow G^{F_1}$ that extends to an injective map

$$(1) \quad (G, X) \hookrightarrow (G^{F_1}, X^{F_1})$$

of Shimura pairs. Here X^{F_1} is the $G^{F_1}(\mathbb{R})$ -conjugacy class of the composite of any $x \in X$ with the monomorphism $G_{\mathbb{R}} \hookrightarrow G_{\mathbb{R}}^{F_1}$. As $G_{\mathbb{R}}^{F_1} \simeq G_{\mathbb{R}}^{[F_1:F(G, X)]}$, the hermitian symmetric domain X^{F_1} is isomorphic to $X^{[F_1:F(G, X)]}$. We have $F(G^{F_1}, X^{F_1}) = F_1$.

Let $(\mathfrak{D}^{F_1}, \mathfrak{n}_X^{F_1})$ be the analogue of the pair $(\mathfrak{D}, \mathfrak{n}_X)$ but for the Shimura pair (G^{F_1}, X^{F_1}) and the pair (T^{F_1}, B^{F_1}) , where T^{F_1} is the maximal torus of G^{F_1} which is the centralizer of T in G^{F_1} and where B^{F_1} is the unique Borel subgroup of $G_{\overline{\mathbb{Q}}}^{F_1}$ that contains both $T_{\overline{\mathbb{Q}}}^{F_1}$ and B . Let $s^{F_1} : \mathfrak{D}^{F_1} \rightarrow \mathfrak{D}$ be the $\text{Gal}(\mathbb{Q})$ -invariant surjective map of Dynkin diagrams which is naturally associated to the monomorphism $G_{\overline{\mathbb{Q}}} \hookrightarrow G_{\overline{\mathbb{Q}}}^{F_1} \simeq G_{\overline{\mathbb{Q}}}^{[F_1:F(G, X)]}$; if $i : F(G, X) \hookrightarrow \mathbb{R}$ is an embedding, then $(s^{F_1})^{-1}(\mathfrak{D}_i)$ is the disjoint union of the connected Dynkin diagrams of those simple factors of $\text{Lie}(G_{\overline{\mathbb{Q}}}^{F_1}) = \bigoplus_{i_1 \in \text{Hom}(F_1, \mathbb{R})} \text{Lie}(G^{F_1} \otimes_{F_1, i_1} \overline{\mathbb{Q}})$ that correspond to embeddings $i_1 : F_1 \hookrightarrow \mathbb{R}$ that extend i . We have $(s^{F_1})^{-1}(\mathfrak{n}_X) = \mathfrak{n}_X^{F_1}$, cf. the very definition of X^{F_1} .

The identity $(s^{F_1})^{-1}(\mathfrak{n}_X) = \mathfrak{n}_X^{F_1}$ implies that the fixed fields of the stabilizers of $\mathfrak{n}_X$ and $\mathfrak{n}_X^{F_1}$ in $\text{Gal}(\mathbb{Q})$ are equal; thus $E(G, X) = E(G^{F_1}, X^{F_1})$. The existence of s^{F_1} also implies that the types of (G, X) and (G^{F_1}, X^{F_1}) are equal, cf. Definition 2.2.1 (a).

2.3. Definitions. Let (G, X) be a simple, adjoint Shimura pair. We say (G, X) is:

- (a) of *inner* (resp. *non-inner*) *type*, if $I(G, X)$ is (resp. is not) $F(G, X)$;
- (b) with *apparent $\mathbb{R}$ -involution*, if $EF(G, X) \neq F(G, X)$;
- (c) with *$\mathbb{R}$ -involution*, if $E(G, X)$ is not a subfield of $G_2(G, X)$;

(d) with *$\mathbb{Q}_p$ -involution*, if there exists a finite field extension $F_{p,1}$ of $\mathbb{Q}_p$ with the property that the group pair $(G_{F_{p,1}}, [\mu_p])$ has a simple factor $(G_1, [\mu_1])$ such that G_1 is a non-split, absolutely simple, unramified group over $F_{p,1}$ and μ_1 is the extension to $\overline{\mathbb{Q}_p}$ of a cocharacter of $G_{1, F_{p,2}}$ whose $G_1(F_{p,2})$ -conjugacy class is not fixed by $\text{Gal}(F_{p,2}/F_{p,1})$; here

$F_{p,2}$ is an unramified field extension of $F_{p,1}$ of degree 2 if (G, X) is not of D_4^{mixed} type and of some degree in the set $\{2, 3\}$ if (G, X) is of D_4^{mixed} type;

- (e) of *compact type*, if the $F(G, X)$ -rank of $G[F(G, X)]$ (i.e., the $\mathbb{Q}$ -rank of G) is 0;
- (f) of *strong compact type*, if the $I(G, X)$ -rank of $G[F(G, X)]_{I(G, X)}$ is 0;
- (g) of *p-compact type*, if the $\mathbb{Q}_p$ -rank of $G_{\mathbb{Q}_p}$ is 0.

2.3.1. Remark. In Definitions 2.3 (b) and (c) the use of “ $\mathbb{R}$ -” emphasizes that we are dealing with totally real number fields $F(G, X)$ and $G_2(G, X)$ while the word involution points out to the fact that suitable group pairs are in the natural sense with involution. The group pairs are not necessarily over $F(G, X)$ or $G_2(G, X)$; often we have to be in a $\mathbb{Q}_p$ -context that is similar to the one of Definition 2.3 (d) (cf. Subsubsection 2.3.3 below).

2.3.2. On involutions. We assume that (G, X) is with involution i.e., $[E(G, X) : E_1(G, X)] = 2$. The field $G(G, X)$ is a Galois extension of $\mathbb{Q}$ that is a totally imaginary quadratic extension of $G_1(G, X)$. Thus (G, X) is also with $\mathbb{R}$ -involution. Standard facts on number fields (for instance, see [La, p. 168]) imply that the set of primes $p \in \mathbb{N}$ which split in $G_1(G, X)$, which do not split in $G(G, X)$, and which have the property that $G_{\mathbb{Q}_p}$ is unramified, has positive Dirichlet density. If p is such a prime, then $G_{\mathbb{Q}_p}$ is a product of absolutely simple, unramified, adjoint groups and each prime of $E_1(G, X)$ (resp. of $E(G, X)$) that divides p has residue field $\mathbb{F}_p$ (resp. $\mathbb{F}_{p^2}$). From this and [Mi3, Prop. 4.6 and Cor. 4.7] we easily get that $(G_{\mathbb{Q}_p}, [\mu_p])$ has a simple factor $(G_1, [\mu_1])$ for which the following two properties hold:

- (a) the group G_1 is non-split, absolutely simple, unramified, and splits over $B(\mathbb{F}_{p^2})$;
- (b) the cocharacter μ_1 is the extension to $\overline{B(\mathbb{F}_{p^2})}$ of a cocharacter $\mu_{1,2}$ of $G_{1,B(\mathbb{F}_{p^2})}$ with the property that the $G_1(B(\mathbb{F}_{p^2}))$ -conjugacy class $[\mu_{1,2}]$ is not fixed by $\text{Gal}(B(\mathbb{F}_{p^2})/\mathbb{Q}_p)$.

By taking $F_{p,1}$ and $F_{p,2}$ to be $\mathbb{Q}_p$ and $B(\mathbb{F}_{p^2})$ (respectively), from the property (b) we get that the Shimura pair (G, X) is also with $\mathbb{Q}_p$ -involution.

2.3.3. On $\mathbb{R}$ -involutions. We assume that (G, X) is with $\mathbb{R}$ -involution. As in Subsubsection 2.3.2 we argue that the set of primes $p \in \mathbb{N}$ which split in $G_2(G, X)$, which do not split in $G(G, X)$, and which have the property that $G_{\mathbb{Q}_p}$ is unramified, has positive Dirichlet density. For such a prime p , if (G, X) is not (resp. is) of D_4^{mixed} type we get the existence of a simple factor $(G_1, [\mu_1])$ of $(G_{\mathbb{Q}_p}, [\mu_p])$ such that properties 2.3.2 (a) and (b) hold (resp. hold with $\mathbb{F}_{p^2}$ replaced by $\mathbb{F}_{p^s}$ for some $s \in \{2, 3\}$). Thus (G, X) is with $\mathbb{Q}_p$ -involution. From the property 2.3.2 (a) we get that G_1 is not an inner form of $I_{0,\mathbb{Q}_p}$ and thus (G, X) is also of non-inner type.

2.3.4. On the non-inner type and the D_{2n} Lie type. If (G, X) is of B_n , C_n , or E_7 type, then $I(G, X) = F(G, X)$. If $n > 1$ and $a \in \{0, \dots, n+1\}$, then the group $\text{SU}(a, n+1-a)_{\mathbb{R}}^{\text{ad}}$ is not an inner form of $\text{PGL}_{n+1,\mathbb{R}}$. Thus, as $F(G, X)$ is a totally real number field, (G, X) is of non-inner type if it is of A_n type with $n > 1$. If (G, X) is of E_6 type, then it is of non-inner type (cf. Subsubsections 2.2.2, 2.3.2, and 2.3.3). Thus the notion non-inner type is of interest only when the simple, adjoint Shimura pair (G, X) is of $D_{2n}^{\mathbb{H}}$, $D_{2n}^{\mathbb{R}}$, or D_{2n}^{mixed} type with $n \geq 2$.

Let now G be such that I_0 is of D_{2n} Lie type, with $n \geq 2$. We have $E(G, X) = E_1(G, X)$, cf. Subsubsection 2.2.2. Each simple factor of $G_{\mathbb{R}}^{\text{ad}}$ is an inner form of $\mathbf{SO}(4n)_{\mathbb{R}}^{\text{ad}}$ (cf. [De2, Subsubsect. 1.3.6]) and thus also of $I_{0, \mathbb{R}}$. Therefore the field $I(G, X)$ is totally real. If $i_0 : F(G, X) \hookrightarrow \mathbb{R}$ is the embedding defined naturally by the inclusion $F(G, X) \hookrightarrow \mathbb{C}$, then (G, X) is of inner type if and only if the action of $\text{Gal}(F(G, X))$ on $\mathfrak{D}_{i_0}$ is trivial. Thus (G, X) is of inner type if and only if the action of $\text{Gal}(\mathbb{Q})$ on the set of extremal vertices of $\mathfrak{D}$ has exactly three orbits.

If (G, X) is of $D_{2n}^{\mathbb{R}}$ type, then the notions of apparent $\mathbb{R}$ -involution and of non-inner type are unrelated (i.e., no one implies the other one). Also, if (G, X) is of $D_{2n}^{\mathbb{R}}$ type, then it is never with $\mathbb{R}$ -involution (cf. [De2, Rm. 2.3.12]) or with $\mathbb{Q}_p$ -involution (this is trivial). If (G, X) is of non-inner $D_{2n}^{\mathbb{H}}$ or D_{2n}^{mixed} type, then each element of $\text{Gal}(F(G, X))$ that does not fix $I(G, X)$ will also not fix $E(G, X)$ and therefore (G, X) is with apparent $\mathbb{R}$ -involution.

2.3.5. Remark. The Shimura pair (G, X) is of compact type if and only if $\text{Sh}(G, X)_{\mathbb{C}}$ is a pro-étale cover of a normal, projective scheme over $\mathbb{C}$; this is implied by [BHC, Thm. 12.3 and Cor. 12.4]. If (G, X) is of p -compact type, then it is also of compact type. If (G, X) is of p -compact type and if all primes of $F(G, X)$ that divide p split in $I(G, X)$, then (G, X) is also of strong compact type.

2.3.6. Example. We assume that $[I(G, X) : F(G, X)] = 2$ and that $G[F(G, X)]$ is the adjoint group of the special unitary group $\mathbf{SU}_1(D, h_D)$, where D is a central division algebra over $I(G, X)$ of dimension $(n+1)^2$ that is equipped with an involution $*$ which is of the second kind and for which we have an identity $F(G, X) = \{x \in I(G, X) | x^* = x\}$ and where h_D is a non-degenerate hermitian form of index 0 relative to $*$. Then $G[F(G, X)]_{I(G, X)}$ is $\mathbf{GL}_1(D)^{\text{ad}}$ and thus has rank 0. Thus the Shimura pair (G, X) is of strong compact type.

2.3.7. Example. We assume that $G_{\mathbb{R}}$ has at least one simple, compact factor and (G, X) is of strong compact type. Let F_1 be a totally real finite field extension of $F(G, X)$ such that the $F_{1, I}$ -rank of $G[F(G, X)]_{F_{1, I}}$ is positive, where $F_{1, I}$ is the composite field of F_1 and $I(G, X)$. We have $I(G^{F_1}, X^{F_1}) = F_{1, I}$. The Shimura pair (G^{F_1}, X^{F_1}) is of compact type (as $G_{\mathbb{R}}^{F_1} \simeq G_{\mathbb{R}}^{[F_1 : F(G, X)]}$ has at least one simple, compact factor) but is not of strong compact type (as the $F_{1, I}$ -rank of $G[F(G, X)]_{F_{1, I}}$ is positive).

2.4. Hodge quasi products. Let $I := \{1, \dots, n\}$. For each element $i \in I$ we consider an injective map

$$f_i : (G_i, X_i) \hookrightarrow (\mathbf{GSp}(W_i, \psi_i), S_i)$$

of Shimura pairs. Let $\mathcal{B}_i := \text{End}(W_i)^{G_i(\mathbb{Q})}$. We say f_i is a *PEL type embedding* if the group G_i is the identity component of the subgroup of $\mathbf{GSp}(W_i, \psi_i)$ that fixes $\mathcal{B}_i$. A Shimura pair that admits a PEL type embedding into a Shimura pair that defines a Siegel modular variety, is called a *Shimura pair of PEL type* (cf. [Sh], [De1], [Zi], [Ko2]).

What follows next extends [Va1, Subsect. 2.5, Ex. 5] (loc. cit. dealt with $n = 2$). Let

$$(W, \psi) := \oplus_{i \in I} (W_i, \varepsilon_i \psi_i),$$

where $\varepsilon_i \in \{-1, 1\}$. Let $G_I := \prod_{i \in I} G_i$. It is a subgroup of both $\prod_{i \in I} \mathbf{GSp}(W_i, \psi_i)$ and $\mathbf{GL}_W$. Let G_0 be the identity component of the intersection $G_I \cap \mathbf{GSp}(W, \psi)$. Let $x : \mathbb{S} \hookrightarrow G_{I, \mathbb{R}}$ be a monomorphism that defines an element of $\prod_{i \in I} X_i$; it factors through $G_{0, \mathbb{R}}$. We choose the ε_i 's (with $i \in I$) such that either $+2\pi i \psi$ or $-2\pi i \psi$ is a polarization of the Hodge $\mathbb{Q}$ -structure on W defined by x . Thus, if X_0 (resp. if S) is the $G_0(\mathbb{R})$ -conjugacy (resp. the $\mathbf{GSp}(W, \psi)(\mathbb{R})$ -conjugacy) class of x , we get an injective map

$$f_0 : (G_0, X_0) \hookrightarrow (\mathbf{GSp}(W, \psi), S)$$

of Shimura pairs which we call a Hodge quasi product of f_i 's. We emphasize that both $\pm\psi$ and f_0 depend uniquely on the choice of x . We denote each such f_0 by $\times_{i \in I}^{\mathcal{H}} f_i$. We refer to the Shimura pair (G_0, X_0) as a *Hodge quasi product* of (G_i, X_i) 's and we denote it by $\times_{i \in I}^{\mathcal{H}} (G_i, X_i)$. If $n = 2$, then we also denote f_0 by $f_1 \times^{\mathcal{H}} f_2$ and (G_0, X_0) by $(G_1, X_1) \times^{\mathcal{H}} (G_2, X_2)$.

We call a $\mathbb{Z}$ -lattice L of W well adapted for f_0 if it is of the form $\oplus_{i \in I} L_i$, where for each $i \in I$ the $\mathbb{Z}$ -lattice L_i of W_i is such that we get a perfect alternating form $\psi_i : L_i \otimes_{\mathbb{Z}} L_i \rightarrow \mathbb{Z}$.

2.4.1. Facts. (a) *The group G_I is the extension of $\mathbb{G}_{m, \mathbb{Q}}^{n-1}$ by G_0 . We have $Z^0(G_0) = \mathbb{G}_{m, \mathbb{Q}}$ if and only if for all $i \in I$ we have $Z^0(G_i) = \mathbb{G}_{m, \mathbb{Q}}$.*

(b) *We assume that for each $i \in I$ the injective map f_i is a PEL type embedding. Then f_0 is also a PEL type embedding.*

Proof: As $\prod_{i \in I} \mathbf{GSp}(W_i, \psi_i)$ is the extension of $\mathbb{G}_{m, \mathbb{Q}}^{n-1}$ by $\mathbf{GSp}(W, \psi) \cap (\prod_{i \in I} \mathbf{GSp}(W_i, \psi_i))$, the first part of (a) follows. The image of each $x_i \in X_i$ contains $Z(\mathbf{GL}_{W_i \otimes_{\mathbb{Q}} \mathbb{R}})$. Thus $Z^0(G_i)$ contains the 1 dimensional split torus $Z(\mathbf{GL}_{W_i})$. The first part of (a) implies that we have a short exact sequence $0 \rightarrow Z^0(G_0) \rightarrow \prod_{i \in I} Z^0(G_i) \rightarrow \mathbb{G}_{m, \mathbb{Q}}^{n-1} \rightarrow 0$. From the last two sentences we get that the second part of (a) also holds.

Part (b) is implied by the fact that G_0 is the identity component of the subgroup of $\mathbf{GSp}(W, \psi)$ that fixes the semisimple $\mathbb{Q}$ -subalgebra $\prod_{i \in I} \mathcal{B}_i$ of $\text{End}(W)$. $\square$

3. Basic techniques and the proof of Theorem 1.3.1

In Subsections 3.1 to 3.3 we review basic facts about Conjecture 1.1.1. In Subsection 3.4 we prove Theorem 1.3.1. Until Section 4 we use the notations of Subsection 1.1. Thus $G_{\mathbb{Q}_p}$ is a connected subgroup of $\mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p}$.

3.1. Simple properties. (a) *The Mumford–Tate conjecture holds for A if and only if it holds for an abelian variety isogenous to $A_{E'}$, where E' is a finite field extension of E .*

(b) *The following three assertions are equivalent:*

(i) *The abelian variety $A_{\overline{E}}$ has complex multiplication (i.e., the semisimple $\mathbb{Q}$ -algebra $\text{End}(A_{\overline{E}}) \otimes_{\mathbb{Z}} \mathbb{Q}$ contains a commutative, semisimple $\mathbb{Q}$ -algebra of dimension $2 \dim(A)$).*

(ii) *The group H_A is a torus.*

(iii) *The group $G_{\mathbb{Q}_p}$ is a torus.*

The equivalence of (i) (resp. (iii)) and (ii) follows from definitions (resp. from Theorem 1.1.4).

(c) As $Z(G_{\mathbb{Q}_p})$ is a subgroup of $Z(H_{A,\mathbb{Q}_p})$ (cf. Theorems 1.1.3 and 1.1.4), the monomorphism $G_{\mathbb{Q}_p} \hookrightarrow H_{A,\mathbb{Q}_p}$ gives birth naturally to a monomorphism $G_{\mathbb{Q}_p}^{\text{ad}} \hookrightarrow H_{A,\mathbb{Q}_p}^{\text{ad}}$.

3.2. Frobenius tori. For the constructions reviewed here we refer to [Ch]. Let v , A_v , and l be as before Theorem 1.1.5. Until Subsection 3.4 we will assume that $p \neq l$. Let $\bar{v}$ be a prime of $\bar{E}$ that divides v . Let $F(\bar{v}) \in \text{Gal}(E)$ be a Frobenius automorphism of $\bar{v}$. The pair $(\bar{v}, F(\bar{v}))$ is uniquely determined up to $\text{Gal}(E)$ -conjugation. As v is a prime of good reduction for A , the restriction of ρ to the inertia group of $\bar{v}$ is trivial (cf. [BLR, Ch. 7, 7.4, Thm. 5]). Thus the element $\rho(F(\bar{v})) \in \mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p}(\mathbb{Q}_p)$ is well defined regardless of the choice of $F(\bar{v})$; it and its positive, integral powers will be referred as *Frobenius elements*. As $\mathbb{Z}_p$ -modules, via $\bar{v}$ we identify $L_A \otimes_{\mathbb{Z}} \mathbb{Z}_p = T_p(A)$ with the Tate module $T_p(A_{v,k(v)})$ of $A_{v,k(v)}$; thus the Frobenius endomorphism Fr_v of A_v acts on $L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p$ as $\rho(F(\bar{v}))$ does.

Let $J_v := \mathbf{GL}_1(\text{End}(A_v) \otimes_{\mathbb{Z}} \mathbb{Q})$. Let T_v be the identity component of the smallest subgroup of J_v that has Fr_v as a $\mathbb{Q}$ -valued point. It is a $\mathbb{Q}$ -model of the identity component of the Zariski closure in $\mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p}$ of the subgroup $\{\rho(F(\bar{v})^s) | s \in \mathbb{Z}\}$ of $\mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p}(\mathbb{Q}_p)$. As Fr_v is a semisimple endomorphism of $L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p$ (see [Tat]), T_v is a torus. Thus $T_{v,\mathbb{Q}_p}$ is a torus of $G_{\mathbb{Q}_p}$, uniquely determined up to $\text{Im}(\rho)$ -conjugation. The torus T_v depends only on some positive, integral power π_v of Fr_v , cf. [Ch, Subsects. 3.a and 3.b]; here we can replace Fr_v by the Frobenius endomorphism of any other abelian variety $B_{v'}$ over a finite field extension $k(v')$ of $k(v)$ that is isogenous to $A_{v,k(v)}$.

If the finite field $k(v')$ is large enough, then we can choose $B_{v'}$ such that there exists an abelian scheme B_R over a complete discrete valuation ring R of mixed characteristic $(0, l)$ and of residue field $k(v')$, which lifts $B_{v'}$ and has complex multiplication. Let $B := B_R \times_R R[\frac{1}{l}]$. We can assume that E is a subfield of $R[\frac{1}{l}]$. We fix an embedding $R[\frac{1}{l}] \hookrightarrow \mathbb{C}$ that extends the embedding i_E of Subsection 1.1. Let H_B be the Mumford–Tate group of $B_{\mathbb{C}}$; as B_R has complex multiplication, H_B is a torus. Let C_B be the centralizer of H_B in $\mathbf{GL}_{H_1(B(\mathbb{C}), \mathbb{Q})}$. Let D_B be the centralizer of C_B (i.e., the double centralizer of H_B) in $\mathbf{GL}_{H_1(B(\mathbb{C}), \mathbb{Q})}$. As B_R has complex multiplication and as $\text{Lie}(C_B)$ is naturally identified with the Lie algebra of $\mathbb{Q}$ -endomorphisms of $B_{\mathbb{C}}$, we have a natural Lie monomorphism $\text{Lie}(C_B) \hookrightarrow \text{End}(B_{v'}) \otimes_{\mathbb{Z}} \mathbb{Q}$. Each invertible element of the $\mathbb{Q}$ -subalgebra of $\text{End}(B_{v'}) \otimes_{\mathbb{Z}} \mathbb{Q}$ generated by the Frobenius endomorphism of $B_{v'}$, is naturally identified with an element of the center of $\text{Lie}(C_B)$ and thus also with a $\mathbb{Q}$ -valued point of D_B ; thus T_v is naturally identified with a subtorus of D_B . If B_R is equipped with a polarization λ_{B_R} and if $\mathcal{B}$ is a semisimple $\mathbb{Q}$ -subalgebra of $\text{End}(H_1(B(\mathbb{C}), \mathbb{Q}))^{H_B(\mathbb{Q})}$, then T_v is in fact a torus of the connected component of the intersection of $\mathbf{GSp}(H_1(B(\mathbb{C}), \mathbb{Q}), \psi_B)$ with the centralizer of $\mathcal{B}$ in $\mathbf{GL}_{H_1(B(\mathbb{C}), \mathbb{Q})}$. Here ψ_B is the alternating form on $H_1(B(\mathbb{C}), \mathbb{Q})$ induced by λ_{B_R} .

The following Lemma is due in essence to Bogomolov and Serre, cf. [Bog] and [Se3, pp. 31, 43, and 44].

3.3. Lemma. *There exists a subtorus T^0 of $Z^0(H_A)$ that does not depend on p and such that $G_{\mathbb{Q}_p}$ is the subgroup of $H_{A,\mathbb{Q}_p}$ generated by $G_{\mathbb{Q}_p}^{\text{der}}$ and $T_{\mathbb{Q}_p}^0$.*

Proof: By replacing E with a finite field extension of it, we can assume that $\text{End}(A) = \text{End}(A_{\overline{\mathbb{Q}}})$. Let C_A be the centralizer of $Z^0(H_A)$ in $\mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}}$. The group C_A is reductive and $Z(C_A)$ is a torus. The commutative, semisimple $\mathbb{Q}$ -algebra $Z(C_A)(\mathbb{Q})$ is a $\mathbb{Q}$ -subalgebra of

$$\text{End}(L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p) = \text{End}(T_p(A) \otimes_{\mathbb{Z}_p} \mathbb{Q}_p) = \text{End}(T_p(A_{v, k(v)}))$$

which is generated by $\mathbb{Q}$ -linear combinations of étale $\mathbb{Q}_p$ -realizations of endomorphisms of A and thus also of A_v . Therefore $Z(C_A)$ is naturally a torus of $\mathbb{J}_v$. Let C_v be the centralizer of $Z(C_A)$ in $\mathbb{J}_v$; it is a reductive group over $\mathbb{Q}$ (cf. [Bor, Ch. IV, 13.17, Cor. 2 (a)]). We identify $\mathbb{J}_{v, \mathbb{Q}_p}$ and $C_{v, \mathbb{Q}_p}$ with subgroups of $\mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}_p}$ that contain $Z(C_A)_{\mathbb{Q}_p}$.

We have $\pi_v \in C_v(\mathbb{Q})$. We denote also by π_v its étale $\mathbb{Q}_p$ -realization (i.e., the corresponding integral power of $\rho(F(\overline{v}))$). Thus $\pi_v \in G_{\mathbb{Q}_p}(\mathbb{Q}_p) \leq H_A(\mathbb{Q}_p) \leq C_A(\mathbb{Q}_p)$. After replacing π_v with a positive, integral power of it that divides the order of the finite group $C_A^{\text{der}} \cap Z(C_A)$, we can write $\pi_v \in C_A(\mathbb{Q}_p)$ as a product

$$\pi_v = \pi_0 \pi_1,$$

where $\pi_0 \in Z(C_A)(\mathbb{Q}_p)$ and $\pi_1 \in C_A^{\text{der}}(\mathbb{Q}_p)$. As $\pi_v \in H_A(\mathbb{Q}_p)$ and as H_A is a reductive subgroup of C_A such that $Z^0(H_A)$ is a subtorus of $Z(C_A)$, after a similar replacement of π_v we get that $\pi_0 \in Z^0(H_A)(\mathbb{Q}_p)$ and $\pi_1 \in H_A^{\text{der}}(\mathbb{Q}_p)$. Let C'_v be the unique reductive subgroup of C_v such that we have a natural isogeny $C'_v \times_{\mathbb{Q}} Z(C_A) \rightarrow C_v$. As $\pi_v \in C_v(\mathbb{Q})$, after a similar replacement of π_v we get that $\pi_0 \in Z(C_A)(\mathbb{Q})$ and $\pi_1 \in C'_v(\mathbb{Q})$.

Thus $\pi_0 \in Z^0(H_A)(\mathbb{Q}_p) \cap Z(C_A)(\mathbb{Q}) = Z^0(H_A)(\mathbb{Q})$. Let T^0 be the identity component of the smallest subgroup of $Z^0(H_A)$ which has π_0 as a $\mathbb{Q}$ -valued point; it is a torus.

We now take v such that $T_{v, \mathbb{Q}_p}$ is a maximal torus of $G_{\mathbb{Q}_p}$, cf. Theorem 1.1.5. The images of $Z^0(G_{\mathbb{Q}_p})$, $G_{\mathbb{Q}_p}$, and $T_{v, \mathbb{Q}_p}$ in $H_{A, \mathbb{Q}_p}^{\text{ab}}$ are the same torus $T_{\mathbb{Q}_p}^1$. As the images of π_v and π_0 in $H_{A, \mathbb{Q}_p}^{\text{ab}}(\mathbb{Q}_p)$ coincide, $T_{\mathbb{Q}_p}^1$ is the extension to $\mathbb{Q}_p$ of $T^1 := \text{Im}(T^0 \rightarrow H_A^{\text{ab}})$. Thus $Z^0(G_{\mathbb{Q}_p})$ is a subtorus of $Z^0(H_A)_{\mathbb{Q}_p}$ (cf. Theorems 1.1.2 and 1.1.4) that has the same image in $H_{A, \mathbb{Q}_p}^{\text{ab}}$ as $T_{\mathbb{Q}_p}^0$. This implies that $Z^0(G_{\mathbb{Q}_p}) = T_{\mathbb{Q}_p}^0$. Thus $G_{\mathbb{Q}_p}$ is the subgroup of $H_{A, \mathbb{Q}_p}$ generated by $G_{\mathbb{Q}_p}^{\text{der}}$ and $T_{\mathbb{Q}_p}^0$.

As $\pi_0 \in Z(C_A)(\mathbb{Q})$ depends only on v , the torus T^0 does not depend on the prime $p \neq l$. The fact that T^0 also does not depend on l , is checked in the standard way by considering another finite prime v_1 of E which is of good reduction for A , whose characteristic is not l , and whose Frobenius torus T_{v_1} has the same rank as T_v . $\square$

3.3.1. Remark. If $T_{v, \mathbb{Q}_p}$ is not a maximal torus of $G_{\mathbb{Q}_p}$, then the same argument shows that the image of $T_{v, \mathbb{Q}_p}$ in $H_{A, \mathbb{Q}_p}^{\text{ab}}$ is the extension to $\mathbb{Q}_p$ of a subtorus of H_A^{ab} .

3.4. Proof of 1.3.1. We prove Theorem 1.3.1. Let H_A^0 be the subgroup of H_A generated by H_A^{der} and the torus T^0 of Lemma 3.3. As $Z^0(G_{\mathbb{Q}_p}) = T_{\mathbb{Q}_p}^0$, to prove Theorem 1.3.1 it suffices to show that we have an identity $T^0 = Z^0(H_A)$. We include a proof of the identity $T^0 = Z^0(H_A)$ that is also used to get several other properties which play key roles in the subsequent Sections. In this Subsection we do not assume that $p \neq l$.

For Hodge cycles on abelian varieties over reduced $\mathbb{Q}$ -schemes we refer to [De3]. It suffices to prove the identity $T^0 = Z^0(H_A)$ under the extra assumptions that $E^{\text{conn}} = E$ and that all Hodge cycles on $A_{\mathbb{C}}$ are also Hodge cycles on A , cf. [De3, Prop. 2.9 and Thm. 2.11]. As the torus T^0 does not depend on p , until Section 4 we will choose the prime v to be unramified over l . We identify E_v with $B(k(v))$. Let $A_{W(k(v))}$ be the abelian scheme over $W(k(v))$ whose generic fibre is $A_{B(k(v))}$. Let (M, ϕ) be the F -crystal (i.e., the contravariant Dieudonné module) of the special fibre A_v of $A_{W(k(v))}$. Thus M is a free $W(k(v))$ -module of rank $2 \dim(A)$ and ϕ is a Frobenius-linear monomorphism of M . Let F^1 be the Hodge filtration of $M = H_{\text{crys}}^1(A_v/W(k(v))) = H_{\text{dR}}^1(A_{W(k(v))}/W(k(v)))$ defined by $A_{W(k(v))}$. We identify $L_A^* \otimes_{\mathbb{Z}} \mathbb{Q}_l$ with $H_{\text{ét}}^1(A_{\overline{B(k(v))}}, \mathbb{Q}_l)$ as $\mathbb{Q}_l$ -vector spaces.

For a vector space V over a field K let $\mathcal{T}(V) := \bigoplus_{s,t \in \mathbb{N} \cup \{0\}} V^{\otimes s} \otimes_K V^{*\otimes t}$. We identify $\text{End}(V) = V \otimes_K V^* \subseteq \mathcal{T}(V)$. The action of ϕ on $M[\frac{1}{l}]$ extends in the standard (tensorial) way to an action of ϕ on $\mathcal{T}(M[\frac{1}{l}])$. Let $(v_{\alpha})_{\alpha \in \mathcal{J}}$ be a family of tensors of $\mathcal{T}(L_A^* \otimes_{\mathbb{Z}} \mathbb{Q}_l)$ such that the reductive group $G_{\mathbb{Q}_l}$ is the subgroup of $\mathbf{GL}_{L_A \otimes_{\mathbb{Z}} \mathbb{Q}_l}$ (equivalently of $\mathbf{GL}_{L_A^* \otimes_{\mathbb{Z}} \mathbb{Q}_l}$) that fixes v_{α} for all $\alpha \in \mathcal{J}$, cf. [De3, Prop. 3.1 (c)].

Let t_{α} be the tensor of $\mathcal{T}(M[\frac{1}{l}])$ that corresponds to v_{α} via Fontaine comparison theory (see [Fo] and [Ts]). It is fixed by the action of ϕ on $\mathcal{T}(M[\frac{1}{l}])$ and moreover it belongs to the F^0 -filtration of the tensor product filtration of $\mathcal{T}(M[\frac{1}{l}])$ defined naturally by the filtration $(F^i[\frac{1}{l}])_{i \in \{0,1,2\}}$ of $M[\frac{1}{l}]$, where $F^2 := 0$ and $F^0 := M$, and by the filtration of $M[\frac{1}{l}]^*$ that is the dual of $(F^i[\frac{1}{l}])_{i \in \{0,1,2\}}$.

Let $\tilde{G}_{B(k(v))}$ be the subgroup of $\mathbf{GL}_{M[\frac{1}{l}]}$ that fixes t_{α} for all $\alpha \in \mathcal{J}$; it is a form of $G_{B(k(v))}$ and thus it is a reductive group. Based on Theorem 1.1.2, we can assume that there exists a subset $\mathcal{J}_1$ of $\mathcal{J}$ such that $(v_{\alpha})_{\alpha \in \mathcal{J}_1}$ is the family of all $\mathbb{Q}_l$ -étale realizations of Hodge cycles on $A_{B(k(v))}$ that belong to $\mathcal{T}(L_A^* \otimes_{\mathbb{Z}} \mathbb{Q}_l)$ (i.e., it is the family of all tensors of $\mathcal{T}(L_A^* \otimes_{\mathbb{Z}} \mathbb{Q})$ fixed by H_A). From this and [Bl, Thm. (0.3)] we get that $(t_{\alpha})_{\alpha \in \mathcal{J}_1}$ is the family of all de Rham realizations of Hodge cycles on $A_{B(k(v))}$ that belong to $\mathcal{T}(M[\frac{1}{l}])$. Let $\tilde{H}_{A,B(k(v))}$ be the subgroup of $\mathbf{GL}_{M[\frac{1}{l}]}$ that fixes t_{α} for all $\alpha \in \mathcal{J}_1$; it is a “de Rham” form of $H_{A,B(k(v))}$. The existence of an isomorphism between the two pairs $(L_A^* \otimes_{\mathbb{Z}} B(k(v)), (v_{\alpha})_{\alpha \in \mathcal{J}})$ and $(M[\frac{1}{l}], (t_{\alpha})_{\alpha \in \mathcal{J}})$ is measured by a class δ_v in the pointed set $H^1(B(k(v)), G_{B(k(v))})$, cf. Fontaine comparison theory. As $G_{B(k(v))}$ acts trivially on $T_{B(k(v))}^0 = Z^0(G_{B(k(v))})$ (via inner conjugation), the image of δ_v in $H^1(B(k(v)), \text{Aut}(T_{B(k(v))}^0))$ is trivial. Thus we can identify $T_{B(k(v))}^0 = Z^0(G_{B(k(v))})$ with the subgroup $Z^0(\tilde{G}_{B(k(v))})$ of either $\tilde{G}_{B(k(v))}$ or $\tilde{H}_{A,B(k(v))}$.

Let $\mu_v : \mathbb{G}_{m,W(k(v))} \rightarrow \mathbf{GL}_M$ be the inverse of the canonical split cocharacter of (M, F^1, ϕ) defined in [Wi, p. 512]. We have a direct sum decomposition $M = F^1 \oplus \tilde{F}^0$ such that $\tilde{F}^0$ is fixed by μ_v and $\beta \in \mathbb{G}_{m,W(k(v))}(W(k(v)))$ acts via μ_v on F^1 as the multiplication with β^{-1} . For $\alpha \in \mathcal{J}$, the tensor t_{α} is fixed by μ_v (cf. the functorial aspects of [Wi, p. 513]). Thus $\mu_{v,B(k(v))}$ factors through $\tilde{G}_{B(k(v))}$.

Let $i_{\overline{\mathbb{Q}_l}} : \overline{\mathbb{Q}_l} \hookrightarrow \mathbb{C}$ be an embedding that extends the embedding i_E of Subsection 1.1. We view $\mathbb{C}$ as a $B(k(v))$ -algebra via the restriction $i_{B(k(v))}$ of $i_{\overline{\mathbb{Q}_l}}$ to $B(k(v))$. The canonical identification $H^1(A(\mathbb{C}), \mathbb{Z}) = L_A^*$ allows us to view H_A and $H_{A,\mathbb{C}}$ naturally as subgroups of the linear groups $\mathbf{GL}_{H^1(A(\mathbb{C}), \mathbb{Q})}$ and $\mathbf{GL}_{H^1(A(\mathbb{C}), \mathbb{C})}$ (respectively). Let P be the parabolic

subgroup of $\tilde{H}_{A,B(k(v))}$ that is the normalizer of $F^1[\frac{1}{l}]$. Under the standard identification of $M_{\mathbb{C}} := M[\frac{1}{l}] \otimes_{B(k(v))} \mathbb{C} = H_{\text{dR}}^1(A_{\mathbb{C}}/\mathbb{C})$ with $H^1(A(\mathbb{C}), \mathbb{C})$, the groups $\tilde{H}_{A,\mathbb{C}}$ and $P_{\mathbb{C}}$ get identified with $H_{A,\mathbb{C}}$ and respectively with the parabolic subgroup of $H_{A,\mathbb{C}}$ that normalizes the direct summand $F^{1,0} = F^1 \otimes_{W(k(v))} \mathbb{C}$ of the Hodge decomposition $M_{\mathbb{C}} = F^{1,0} \oplus F^{0,1}$ of $A_{\mathbb{C}}$. Both μ_A and $\mu_{v,\mathbb{C}}$ are (identified with) cocharacters of $P_{\mathbb{C}}$.

Let $g \in P(\mathbb{C})$ be such that $g\mu_A g^{-1}$ commutes with $\mu_{v,\mathbb{C}}$. As the commuting cocharacters $g\mu_A g^{-1}$ and $\mu_{v,\mathbb{C}}$ act in the same way on $F^{1,0}$ and on $M_{\mathbb{C}}/F^{1,0}$, they coincide. As $\mu_{v,B(k(v))}$ factors through $\tilde{G}_{B(k(v))}$ and as $T_{B(k(v))}^0 = Z^0(\tilde{G}_{B(k(v))})$, the cocharacter $\mu_{v,\mathbb{C}} = g\mu_A g^{-1}$ factors through the normal subgroup $H_{A,\mathbb{C}}^0$ of $H_{A,\mathbb{C}}$. From the last two sentences, we get that both $g\mu_A g^{-1}$ and μ_A factor through $H_{A,\mathbb{C}}^0$. Due to the smallest property of H_A (see Subsection 1.1) and the fact that μ_A factors through $H_{A,\mathbb{C}}^0$, we get that $H_A = H_A^0$. Thus $T^0 = Z^0(H_A)$. This ends the proof of Theorem 1.3.1. $\square$

3.4.1. Simple properties. (a) Conjecture 1.1.1 holds if A has complex multiplication over $\overline{\mathbb{Q}}$. This well known fact also follows from the property 3.1 (b) and Theorems 1.1.2 and 1.3.1.

(b) The monomorphism $\tilde{G}_{\overline{\mathbb{Q}_l}} \hookrightarrow \tilde{H}_{A,\overline{\mathbb{Q}_l}}$ is naturally identified up to $H_A(\overline{\mathbb{Q}_l})$ -conjugation with the monomorphism $G_{\overline{\mathbb{Q}_l}} \hookrightarrow H_{A,\overline{\mathbb{Q}_l}}$, cf. the existence of the class δ_v . As μ_v factors through $\tilde{G}_{B(k(v))}$ and as the cocharacters $\mu_{v,\mathbb{C}}$ and μ_A of $H_{A,\mathbb{C}}$ are $H_A(\mathbb{C})$ -conjugate, there exists a cocharacter $\mu_{v,\overline{\mathbb{Q}_l}}^{\text{ét}} : \mathbb{G}_{m,\overline{\mathbb{Q}_l}} \rightarrow G_{\overline{\mathbb{Q}_l}}$ whose extension to $\mathbb{C}$ (via $i_{\overline{\mathbb{Q}_l}} : \overline{\mathbb{Q}_l} \hookrightarrow \mathbb{C}$) is $H_A(\mathbb{C})$ -conjugate to μ_A .

(c) We use the notations of Subsection 3.4. The element $\phi^{[k(v):\mathbb{F}_l]} \in \tilde{G}_{B(k(v))}(B(k(v)))$ is the crystalline realization of Fr_v and thus it is a semisimple element. This implies that the torus $T_{v,B(k(v))}$ is naturally identified with the smallest subtorus of $\tilde{G}_{B(k(v))}$ that has $\phi^{s[k(v):\mathbb{F}_l]}$ as a $B(k(v))$ -valued point, where $s \in \mathbb{N}$ is such that $\pi_v = Fr_v^s$. Thus $T_{v,\overline{\mathbb{Q}_l}}$ is naturally identified with a torus of $\tilde{G}_{\overline{\mathbb{Q}_l}}$ and therefore (cf. (b)) also of $G_{\overline{\mathbb{Q}_l}}$.

3.4.2. Corollary. *We assume that $E^{\text{conn}} = E$, that v is unramified over l , that $k(v) = \mathbb{F}_l$, and that T_v has the same rank as $G_{\mathbb{Q}_l}$. We also assume that T_v is (isomorphic to) a torus of H_A in such a way that there exists an element $g_v \in H_A(\overline{\mathbb{Q}_l})$ with the property that the torus $g_v T_{v,\overline{\mathbb{Q}_l}} g_v^{-1}$ of $H_{A,\overline{\mathbb{Q}_l}}$ is a maximal torus of $G_{\overline{\mathbb{Q}_l}}$. Let $i_{\overline{\mathbb{Q}_l}} : \overline{\mathbb{Q}_l} \hookrightarrow \mathbb{C}$ be an embedding that extends the embedding $i_E : E \hookrightarrow \mathbb{C}$ of Subsection 1.1. Let $\mathfrak{C}_l(H_A, X_A)$ be the set of cocharacters of $H_{A,\overline{\mathbb{Q}_l}}$ whose extensions to $\mathbb{C}$ via $i_{\overline{\mathbb{Q}_l}}$ belong to the set $\mathfrak{C}(H_A, X_A)$ of cocharacters of $H_{A,\mathbb{C}}$ defined in Subsection 2.2. Then the torus $T_{v,\overline{\mathbb{Q}_l}}$ of $H_{A,\overline{\mathbb{Q}_l}}$ is generated by images of certain cocharacters in the set $\mathfrak{C}_l(H_A, X_A)$.*

Proof: We can choose the element g_v such that the cocharacter $\mu_{v,\overline{\mathbb{Q}_l}}^{\text{ét}}$ of the property 3.4.1 (b) factors through $g_v T_{v,\overline{\mathbb{Q}_l}} g_v^{-1}$. As the cocharacters $\mu_{v,\mathbb{C}}$ and μ_A are $P(\mathbb{C})$ -conjugate (see Subsection 3.4) and thus also $H_A(\mathbb{C})$ -conjugate, the set $\mathfrak{C}_l(H_A, X_A)$ is the set of $H_A(\overline{\mathbb{Q}_l})$ -conjugates of the $\text{Gal}(\overline{\mathbb{Q}}/\overline{\mathbb{Q}_l})$ -conjugates of the cocharacter $g_v^{-1} \mu_{v,\overline{\mathbb{Q}_l}}^{\text{ét}} g_v$ of the extension of T_v to $\overline{\mathbb{Q}_l}$ (equivalently to $\overline{\mathbb{Q}}$). But up to $G_{\mathbb{Q}_l}(\overline{\mathbb{Q}_l})$ -conjugation, we can choose the cocharacter $\mu_{v,\overline{\mathbb{Q}_l}}^{\text{ét}} : \mathbb{G}_{m,\overline{\mathbb{Q}_l}} \rightarrow G_{\overline{\mathbb{Q}_l}}$ such that $g_v^{-1} \mu_{v,\overline{\mathbb{Q}_l}}^{\text{ét}} g_v$ is a strong Hodge cocharacter of $T_{v,\overline{\mathbb{Q}_l}}$ in the sense of [Pi, Def. 3.14] (see [Pi, Prop. 3.13 and paragraph before Def. 3.14]). Thus the

fact that $T_{v, \overline{\mathbb{Q}}_l}$ is generated by images of certain cocharacters in the set $\mathfrak{C}_l(H_A, X_A)$ is only a weaker form of [Pi, Thm. 3.16]. $\square$

4. Injective maps into unitary Shimura pairs of PEL type

In Subsections 4.1 to 4.9 we include different rational versions of the embedding results of [Sa1]; the approach is close in spirit to [Sa2], [De2, Prop. 2.3.10], and [Va1, Subsects. 6.5 and 6.6]. Corollary 4.10 classifies all adjoint Shimura pairs that are adjoints of Shimura pairs of PEL type. Until Section 5 the notations are independent of the ones of Subsections 1.1, 1.3, or 1.5.

Until Subsection 4.9 let (G, X) be a simple, adjoint Shimura pair of abelian type. Thus G is a simple, adjoint group over $\mathbb{Q}$. Let $F := F(G, X)$. We write $G = \text{Res}_{F/\mathbb{Q}} G[F]$, cf. Subsection 2.2. For a totally real number field F_1 that contains F , let $G^{F_1} := \text{Res}_{F_1/\mathbb{Q}} G[F]_{F_1}$ and $(G, X) \hookrightarrow (G^{F_1}, X^{F_1})$ be as in Subsubsection 2.2.3.

4.1. Proposition. *We assume that the simple, adjoint Shimura pair (G, X) is of A_n type. Then there exists an injective map*

$$f_2: (G_2, X_2) \hookrightarrow (\mathbf{GSp}(W, \psi), S)$$

of Shimura pairs such that the following two properties hold:

- (i) *the adjoint Shimura pair $(G_2^{\text{ad}}, X_2^{\text{ad}})$ is isomorphic to (G, X) ;*
- (ii) *the group G_2 is the subgroup of $\mathbf{GSp}(W, \psi)$ that fixes $\mathcal{B} := \text{End}(W)^{G_2(\mathbb{Q})}$ (thus f_2 is a PEL type embedding).*

Proof: Let $G[F]^{\text{sc}}$ be the simply connected semisimple group cover of $G[F]$. The group $G^{\text{sc}} := \text{Res}_{F/\mathbb{Q}} G[F]^{\text{sc}}$ is the simply connected semisimple group cover of G . Let T be a maximal torus of G^{sc} . Let F_0 be a finite field extension of F such that the group $G[F]_{F_0}^{\text{sc}}$ is split (i.e., it is an $\mathbf{SL}_{n+1, F_0}$ group). For instance, we can take F_0 to be the smallest Galois extension of $\mathbb{Q}$ such that the torus T_{F_0} is split. Let K be a totally imaginary quadratic extension of F that is linearly disjoint from F_0 ; if $n = 1$ and $G[F]$ is non-split (i.e., if (G, X) is of compact A_1 type), then we choose K such that $G[F]_K$ is also non-split. If $n > 1$ (resp. $n = 1$) let E_0 be F_0 (resp. be $F_0 \otimes_F K$). Let W_0 be an F_0 -vector space of dimension $n + 1$. Let $W_2 := W_0 \otimes_{F_0} E_0$; when we view W_2 as a $\mathbb{Q}$ -vector space we denote it by W . We identify $G[F]_{F_0}^{\text{sc}} = \mathbf{GL}_{W_0}^{\text{der}}$ and thus also $G[F]_{E_0}^{\text{sc}} = \mathbf{GL}_{W_2}^{\text{der}}$. We have natural monomorphisms $G^{\text{sc}} \hookrightarrow \text{Res}_{E_0/\mathbb{Q}} G[F]_{E_0}^{\text{sc}} \hookrightarrow \mathbf{GL}_W^{\text{der}}$ and thus we can view G^{sc} as a subgroup of $\mathbf{GL}_W$. Next we follow the proof of [De2, Prop. 2.3.10].

Let $\mathcal{S}$ be the set of extremal vertices of the Dynkin diagram $\mathfrak{D}$ of $\text{Lie}(G_{\overline{\mathbb{Q}}})$ with respect to $T_{\overline{\mathbb{Q}}}$ and a Borel subgroup of $G_{\overline{\mathbb{Q}}}$ that contains $T_{\overline{\mathbb{Q}}}$. The Galois group $\text{Gal}(\mathbb{Q})$ acts on $\mathcal{S}$ (see Subsection 2.2). If $n > 1$ we can identify $\mathcal{S}$ with the $\text{Gal}(\mathbb{Q})$ -set $\text{Hom}_{\mathbb{Q}}(\mathcal{K}, \overline{\mathbb{Q}})$, where $\mathcal{K} := I(G, X)$ is a totally imaginary quadratic extension of F . If $n = 1$ let $\mathcal{K} := K$. Always $\mathcal{K}$ is a subfield of E_0 and thus W_2 is naturally a $\mathcal{K}$ -vector space. Therefore the torus $\mathcal{T} := \text{Res}_{\mathcal{K}/\mathbb{Q}} \mathbb{G}_{m, \mathcal{K}}$ acts naturally on W . If $n > 1$ this action over $\overline{\mathbb{Q}}$ can be described as follows: if V_0 is a simple $G_{\overline{\mathbb{Q}}}^{\text{sc}}$ -submodule of $W \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}$ whose highest weight is the fundamental

weight associated to a vertex $\mathbf{n} \in \mathcal{S}$, then $\mathcal{T}_{\overline{\mathbb{Q}}}$ acts on V_0 via its character that corresponds naturally to $\mathbf{n}$. As $\mathcal{K}$ is a totally imaginary quadratic extension of F , for $n \geq 1$ the extension to $\mathbb{R}$ of the quotient torus $\mathcal{T}/\text{Res}_{F/\mathbb{Q}}\mathbb{G}_{m,F}$ is compact. This implies that the maximal subtorus $\mathcal{T}_c$ of $\mathcal{T}$ which over $\mathbb{R}$ is compact, is isomorphic to $\mathcal{T}/\text{Res}_{F/\mathbb{Q}}\mathbb{G}_{m,F}$ and therefore it has rank $[F : \mathbb{Q}]$.

The centralizer C of G^{sc} in $\mathbf{GL}_W$ is a reductive group and $Z(C)$ is a torus of rank equal to $2[F : \mathbb{Q}]$ if $n > 1$ (resp. to $[F : \mathbb{Q}]$ if $n = 1$). If $n > 1$ (resp. $n = 1$) we view $\mathcal{T}$ and $\mathcal{T}_c$ as subtori of $Z(C)$ (resp. of C). Let $\mathcal{T}_{c,+}$ be the torus of C generated by $\mathcal{T}_c$ and $Z(\mathbf{GL}_W)$; its rank is $[F : \mathbb{Q}] + 1$ and it commutes with G^{sc} . Let G_2 be the reductive subgroup of $\mathbf{GL}_W$ that is generated by G^{sc} and $\mathcal{T}_{c,+}$; thus we have $G_2^{\text{der}} = G^{\text{sc}}$ and $Z^0(G_2) = \mathcal{T}_{c,+}$. The centralizer C_2 of G_2 in $\mathbf{GL}_W$ is also a reductive group. Let $\mathcal{B}$ be the semisimple $\mathbb{Q}$ -subalgebra of $\text{End}(W)$ defined by the elements of $\text{Lie}(C_2)$. We have $\mathcal{B} = \text{End}(W)^{G_2(\mathbb{Q})}$.

We fix an element $x \in X$. We will construct a monomorphism $x_2 : \mathbb{S} \hookrightarrow G_{2,\mathbb{R}}$ such that it defines a Hodge $\mathbb{Q}$ -structure on W of type $\{(-1, 0), (0, -1)\}$ and it lifts the homomorphism $x : \mathbb{S} \rightarrow G_{2,\mathbb{R}}^{\text{ad}} = G_{\mathbb{R}}$.

Let $\mathfrak{R} := \text{Hom}_{\mathbb{Q}}(F, \mathbb{R})$. We have $G_{\mathbb{R}}^{\text{sc}} = \prod_{i \in \mathfrak{R}} G[F]^{\text{sc}} \times_{F,i} \mathbb{R}$. For $i \in \mathfrak{R}$ let V_i be the real vector subspace of $W \otimes_{\mathbb{Q}} \mathbb{R}$ generated by its simple $G[F]^{\text{sc}} \times_{F,i} \mathbb{R}$ -submodules. We have a product decomposition $\mathcal{T}_{\mathbb{R}} = \prod_{i \in \mathfrak{R}} T_i$ in 2 dimensional tori such that for each pair $(i, i') \in \mathfrak{R}^2$, the torus T_i acts trivially on $V_{i'}$ if and only if $i \neq i'$. Each T_i is isomorphic to $\mathbb{S}$ and acts faithfully on V_i . We have

$$\mathcal{T}_{c,\mathbb{R}} = \prod_{i \in \mathfrak{R}} T_{c,i},$$

where $T_{c,i}$ is the 1 dimensional compact subtorus of T_i . The direct sum decomposition

$$W \otimes_{\mathbb{Q}} \mathbb{R} = \oplus_{i \in \mathfrak{R}} V_i$$

is normalized by $G_{2,\mathbb{R}}$. We have a unique direct sum decomposition

$$(2) \quad V_i \otimes_{\mathbb{R}} \mathbb{C} = V_i^+ \oplus V_i^-$$

such that $\mathcal{T}_{\mathbb{C}}$ acts on each V_i^u via a unique character that is uniquely determined by the pair (i, u) (here $u \in \{-, +\}$). Each $G_{2,\mathbb{C}}^{\text{der}}$ -module V_i^u is isotypic (i.e., is a direct sum of isomorphic simple $G_{2,\mathbb{C}}^{\text{der}}$ -modules). More precisely, the highest weight of the representation of the factor $G[F]^{\text{sc}} \times_{F,i} \mathbb{C}$ of $G_{2,\mathbb{C}}^{\text{der}}$ on V_i^u is $\varpi_{s_i(u)}$, where $s_i : \{+, -\} \rightarrow \{1, n\}$ is a surjective function. Thus $I_i := \text{Im}(T_i \rightarrow \mathbf{GL}_{V_i})$ is the center of the centralizer of $C_{2,\mathbb{R}}$ in $\mathbf{GL}_{V_i}$.

We will take $x_2 : \mathbb{S} \hookrightarrow G_{2,\mathbb{R}}$ such that its restriction to the split subtorus $\mathbb{G}_{m,\mathbb{R}}$ of $\mathbb{S}$ induces an isomorphism $\mathbb{G}_{m,\mathbb{R}} \xrightarrow{\sim} Z(\mathbf{GL}_{W \otimes_{\mathbb{Q}} \mathbb{R}})$ and the following two properties hold:

- (a) if $i \in \mathfrak{R}$ is such that $G[F] \times_{F,i} \mathbb{R}$ is non-compact, then the homomorphism $\mathbb{S} \rightarrow \mathbf{GL}_{V_i}$ defined by x_2 is constructed as in the proof of [De2, Prop. 2.3.10] (thus this homomorphism factors through $\text{Im}(G_{2,\mathbb{R}} \rightarrow \mathbf{GL}_{V_i})$, defines a Hodge $\mathbb{R}$ -structure on V_i of type $\{(-1, 0), (0, -1)\}$, and it is uniquely determined by these two properties and the fact that it lifts the homomorphism $\mathbb{S} \rightarrow G[F] \times_{F,i} \mathbb{R}$ defined by x);

- (b) if $i \in \mathfrak{R}$ is such that $G[F] \times_{F,i} \mathbb{R}$ is compact, then the homomorphism $\mathbb{S} \rightarrow \mathbf{GL}_{V_i}$ defined by x_2 is a monomorphism whose image is naturally identified with I_i (and thus whose image in $G[F] \times_{F,i} \mathbb{R}$ is trivial).

For the rest of the proof it is irrelevant which one of the two possible natural identifications of (b) we choose; each such choice is obtained naturally from the other one via the standard non-trivial automorphism of the compact subtorus of $\mathbb{S}$.

Let X_2 be the $G_2(\mathbb{R})$ -conjugacy class of x_2 . The homomorphism x_2 lifts $x \in X$, cf. (a) and (b) and the fact (see Subsection 2.2) that the image of x in each simple, compact factor of $G_{\mathbb{R}}$ is trivial. Therefore the pair (G_2, X_2) is a Shimura pair whose adjoint is naturally isomorphic to (G, X) . Thus (i) holds. The torus $\mathcal{T}$ splits over the field of CM type which is the Galois extension of $\mathbb{Q}$ generated by $\mathcal{K}$. Due to (a) and (b), the homomorphism x_2 defines a Hodge $\mathbb{Q}$ -structure on W of type $\{(-1, 0), (0, -1)\}$. From the last two sentences and [De2, Cor. 2.3.3], we get that there exists a non-degenerate alternating form ψ on W such that we have an injective map $f_2 : (G_2, X_2) \hookrightarrow (\mathbf{GSp}(W, \psi), S)$ of Shimura pairs.

The representation of $G_{2, \overline{\mathbb{Q}}}$ on $W \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}$ is a direct sum of irreducible representations of dimension $n + 1$ which (as $\overline{\mathbb{Q}}$ -vector spaces) are isotropic with respect to ψ . For all $n \geq 1$, the number of non-isomorphic such irreducible representations is $2[F : \mathbb{Q}]$. Thus the subgroup $G'_{2, \overline{\mathbb{Q}}}$ of $\mathbf{GSp}(W \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}, \psi)$ that centralizes $\mathcal{B} \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}$ (equivalently $C_{2, \overline{\mathbb{Q}}}$) is a reductive group that has the following properties: it contains $G_{2, \overline{\mathbb{Q}}}$, $G'_{2, \overline{\mathbb{Q}}} \stackrel{\text{der}}{\simeq} \mathbf{SL}_{n+1, \overline{\mathbb{Q}}}^{[F: \mathbb{Q}]}$, and $Z^0(G'_{2, \overline{\mathbb{Q}}}) \simeq \mathbb{G}_{m, \overline{\mathbb{Q}}}^{[F: \mathbb{Q}] + 1}$. By reasons of dimensions we have $G_{2, \overline{\mathbb{Q}}} = G'_{2, \overline{\mathbb{Q}}}$. Therefore the subgroup of $\mathbf{GSp}(W, \psi)$ that centralizes $\mathcal{B}$ (equivalently C_2) is G_2 . Thus (ii) also holds. $\square$

4.1.1. Two properties. (a) The double centralizer G_5 of G_2 in $\mathbf{GL}_W$ is such that $G_{5, \overline{\mathbb{Q}}}$ is isomorphic to $\mathbf{GL}_{n+1, \overline{\mathbb{Q}}}^{2[F: \mathbb{Q}]}$ and G_5^{ad} is $\text{Res}_{\mathcal{K}/\mathbb{Q}} G[F]_{\mathcal{K}}$. The semisimple $\mathbb{Q}$ -algebra $\mathcal{B}$ has $\mathcal{K}$ as its center and thus it is simple. Moreover, if $n > 1$ (resp. $n = 1$) and (G, X) is of strong compact type (resp. of compact type), then from the description of $\mathcal{K}$ in the proof of Proposition 4.1 we get that the $\mathcal{K}$ -rank of $G[F]_{\mathcal{K}}$ is 0; thus the $\mathbb{Q}$ -rank of G_5^{ad} is also 0.

(b) We assume that there exists a number $a \in \mathbb{N}$ such that $n = 2a - 1$ and $G_{\mathbb{R}}^{\text{sc}} = G_{2, \mathbb{R}}^{\text{der}}$ is a product of $\mathbf{SU}(a, a)_{\mathbb{R}}$ groups. Then the monomorphism $x_2 : \mathbb{S} \hookrightarrow G_{2, \mathbb{R}}$ factors through the subgroup of $G_{2, \mathbb{R}}$ generated by $G_{2, \mathbb{R}}^{\text{der}}$ and $Z(\mathbf{GL}_{W \otimes_{\mathbb{Q}} \mathbb{R}})$, cf. [De2, Rms. 2.3.12 and 2.3.13]. In other words, the homomorphism $\mathbb{S} \rightarrow \mathbf{GL}_{V_i}$ of the property 4.1 (a) factors through the subgroup of $\mathbf{GL}_{V_i}$ generated by $\text{Im}(G_{2, \mathbb{R}}^{\text{der}} \rightarrow \mathbf{GL}_{V_i})$ and $Z(\mathbf{GL}_{V_i})$.

4.1.2. A variant of 4.1. We refer to the proof of Proposition 4.1 with $n > 1$ and with (G, X) without involution. We take E_0 to be $F_0 \otimes_F K$, with K a totally imaginary quadratic extension of F that is linearly disjoint from $\mathcal{K} = I(G, X)$. Let $\mathcal{T} := \text{Res}_{\mathcal{K} \otimes_F K / \mathbb{Q}} \mathbb{G}_{m, \mathcal{K} \otimes_F K}$. As in the proof of Proposition 4.1, W is naturally a $\mathcal{K} \otimes_F K$ -module and thus a $\mathcal{T}$ -module. Let $\mathcal{T}_c$ and $\mathcal{T}_{c,+}$ be the subtori of $\mathcal{T}$ obtained as in the proof of Proposition 4.1; their ranks are $2[F : \mathbb{Q}]$ and $2[F : \mathbb{Q}] + 1$ (respectively). We take G_2 to be defined as in the proof of Proposition 4.1. We have a unique direct sum decomposition in $G_{2, \mathbb{C}}$ -modules

$$V_i \otimes_{\mathbb{R}} \mathbb{C} = V_i^{+,+} \oplus V_i^{+,-} \oplus V_i^{-,+} \oplus V_i^{-,-}$$

such that all irreducible subrepresentations of the representation of $G_{2,\mathbb{C}}^{\text{der}}$ (resp. of $Z^0(G_{2,\mathbb{C}})$) on $V_i^{u,+} \oplus V_i^{u,-}$ (resp. on $V_i^{+,u} \oplus V_i^{-,u}$) are isomorphic; here $u \in \{-, +\}$.

The proof of [De2, Prop. 2.3.10] guarantees the existence of a monomorphism $x_2 : \mathbb{S} \hookrightarrow G_{2,\mathbb{R}}$ such that it defines a Hodge $\mathbb{Q}$ -structure on W of type $\{(-1, 0), (0, -1)\}$ and it lifts the homomorphism $x : \mathbb{S} \rightarrow G_{2,\mathbb{R}}^{\text{ad}} = G_{\mathbb{R}}$. As in the proof of Proposition 4.1 we argue that there exists a non-degenerate alternating form ψ on W such that we have an injective map $f_2 : (G_2, X_2) \hookrightarrow (\mathbf{GSp}(W, \psi), S)$ of Shimura pairs. The representation of $G_{2,\overline{\mathbb{Q}}}$ on $W \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}$ is a direct sum of irreducible representations which (as $\overline{\mathbb{Q}}$ -vector spaces) are isotropic with respect to ψ . But this time the number of non-isomorphic such irreducible representations is $4[F : \mathbb{Q}]$. Thus the subgroup G_4 of $\mathbf{GSp}(W, \psi)$ that centralizes $\mathcal{B} = \text{End}(W)^{G_2(\mathbb{Q})}$ is not G_2 but a reductive group with the property that $G_{4,\overline{\mathbb{Q}}}^{\text{der}}$ is isomorphic to $(G_{2,\overline{\mathbb{Q}}}^{\text{der}})^2$.

As $G_2 \leq G_4 \leq \mathbf{GSp}(W, \psi)$ and as obviously there exists no simple factor of G_4^{ad} that over $\mathbb{R}$ is compact, the $G_4(\mathbb{R})$ -conjugacy class X_4 of x_2 has the property that we have injective maps $(G_2, X_2) \hookrightarrow (G_4, X_4) \hookrightarrow (\mathbf{GSp}(W, \psi), S)$ of Shimura pairs. The injective map $(G_4, X_4) \hookrightarrow (\mathbf{GSp}(W, \psi), S)$ of Shimura pairs is a PEL type embedding, cf. the very definition of G_4 . As $F_0 \otimes_F K$ is a field (cf. our linearly disjoint assumption), the center of the $\mathbb{Q}$ -algebra $\mathcal{B}$ is a $\mathbb{Q}$ -subalgebra of $F_0 \otimes_F K$ and thus it is a field. Therefore the $\mathbb{Q}$ -algebra $\mathcal{B}$ is simple. Thus the injective map $(G, X) = (G_2^{\text{ad}}, X_2^{\text{ad}}) \hookrightarrow (G_4^{\text{ad}}, X_4^{\text{ad}})$ of Shimura pairs is as in (1) of Subsubsection 2.2.3, with F_1 as the unique totally real quadratic extension of F contained in $F_0 \otimes_F K$. We have identities $E(G_2^{\text{ad}}, X_2^{\text{ad}}) = E(G, X) = E(G^{F_1}, X^{F_1}) = E(G_4^{\text{ad}}, X_4^{\text{ad}})$, cf. Subsubsection 2.2.3.

4.1.3. Remark. If in Proposition 4.1 (resp. in Subsubsection 4.1.2) we have $n > 1$ and we can take G_2 (resp. G_4) such that $Z^0(G_2) = \mathbb{G}_{m,\mathbb{Q}}$ (resp. such that $Z^0(G_4) = \mathbb{G}_{m,\mathbb{Q}}$), then the property 4.1 (ii) (resp. the part of Subsubsection 4.1.2 that involves PEL type embeddings) does not hold anymore.

4.2. The group $\tilde{G}$. Until Subsection 4.9 we will assume that (G, X) is of B_n , C_n , $D_n^{\mathbb{H}}$, or $D_n^{\mathbb{R}}$ type. All simple, non-compact factors of $G_{\mathbb{R}}$ are isomorphic. We now introduce the “simplest” absolutely simple, adjoint group over $\mathbb{Q}$ with the property that $\tilde{G}_{\mathbb{R}}$ is isomorphic to each simple, non-compact factor of $G_{\mathbb{R}}$. Depending on the type of (G, X) we take $\tilde{G}$ as follows (the values of n are such that we do not get repetitions among the Shimura types).

- If (G, X) is of B_n type ($n > 1$), then $\tilde{G} := \mathbf{SO}(2, 2n - 1)$.
- If (G, X) is of C_n type ($n > 2$), then $\tilde{G} := \mathbf{Sp}_{2n,\mathbb{Q}}^{\text{ad}}$.
- If (G, X) is of $D_n^{\mathbb{H}}$ type ($n \geq 4$), then $\tilde{G} := \mathbf{SO}^*(2n)^{\text{ad}}$.
- If (G, X) is of $D_n^{\mathbb{R}}$ type ($n \geq 4$), then $\tilde{G} := \mathbf{SO}(2, 2n - 2)^{\text{ad}}$.

Let $\tilde{G}^{\text{sc}}$ be the simply connected semisimple group cover of $\tilde{G}$. Let $\tilde{x} : \mathbb{S} \rightarrow \tilde{G}_{\mathbb{R}}$ be a homomorphism whose $\tilde{G}(\mathbb{R})$ -conjugacy class $\tilde{X}$ has the property that the pair $(\tilde{G}, \tilde{X})$ is a Shimura pair. We know that $\tilde{x}$ is unique up to $\tilde{G}(\mathbb{R})$ -conjugation and possibly up to a replacement of it by its inverse, cf. [De2, Prop. 1.2.7 and Cor. 1.2.8].

4.3. The group $G(A)$. Until Subsection 4.9 we will have the same restrictions on n as in Subsection 4.2. Let the group $G(A)$ over $\mathbb{Q}$ be defined by the rules:

- if (G, X) is of B_n or $D_n^{\mathbb{R}}$ type, then $G(A) := \mathbf{SU}(2^{n-1}, 2^{n-1})$;
- if (G, X) is of C_n or $D_n^{\mathbb{H}}$ type, then $G(A) := \mathbf{SU}(n, n)$.

These rules conform with [Sa1, Subsects. 3.3 to 3.7 or p. 461]. Subsections [Sa1, 1.1 and 3.3 to 3.7] are stated in terms of simple Lie algebras over $\mathbb{R}$ and show the existence of a Lie monomorphism $dh_{\mathbb{R}} : \mathrm{Lie}(\tilde{G}_{\mathbb{R}}^{\mathrm{sc}}) \hookrightarrow \mathrm{Lie}(G(A)_{\mathbb{R}})$ that satisfies the condition (H_2) of [Sa1, end of Subsect. 1.1]. This condition says that we have an identity

$$(3) \quad (dh_{\mathbb{R}} \circ d\tilde{x})(\sqrt{-1}) = d\tilde{y}(\sqrt{-1}) \in \mathrm{Lie}(G(A)_{\mathbb{R}}^{\mathrm{ad}}),$$

where $\sqrt{-1} \in \mathbb{C} = \mathrm{Lie}(\mathbb{S})$, where $\tilde{x}$ is as in the end of Subsection 4.2, and where $\tilde{y} : \mathbb{S} \rightarrow G(A)_{\mathbb{R}}^{\mathrm{ad}}$ is a homomorphism that induces a monomorphism $\mathbb{S}/\mathbb{G}_{m, \mathbb{R}} \hookrightarrow G(A)_{\mathbb{R}}^{\mathrm{ad}}$ whose image is the identity component of the center of a maximal compact subgroup of $G(A)_{\mathbb{R}}^{\mathrm{ad}}$. Here $d\tilde{x}$ and $d\tilde{y}$ are the Lie homomorphisms induced naturally by $\tilde{x}$ and $\tilde{y}$ (respectively). Let

$$h_{\mathbb{R}} : \tilde{G}_{\mathbb{R}}^{\mathrm{sc}} \rightarrow G(A)_{\mathbb{R}}$$

be the homomorphism of finite kernel whose differential at the level of Lie algebras is $dh_{\mathbb{R}}$.

The only difference here from [Sa1, Subsects. 3.3 to 3.7 or p. 461] is the following one. See [He, Ch. X, Sect. 6, Table V] for the classification of hermitian symmetric domains. If (G, X) is of $D_n^{\mathbb{R}}$ type, then $\tilde{X}$ is two copies of the hermitian symmetric domain $BDI(2, 2n-2)$ associated to the group $\mathbf{SO}(2, 2n-2)_{\mathbb{R}}^{\mathrm{ad}}$. In [Sa1, Subsect. 3.5] this hermitian symmetric domain is embedded into the hermitian symmetric domain $AIII(2^{n-2}, 2^{n-2})$ associated to the group $\mathbf{SU}(2^{n-2}, 2^{n-2})_{\mathbb{R}}^{\mathrm{ad}}$. We have two such embeddings: they correspond to the two half spin representations of $\tilde{G}_{\mathbb{C}}^{\mathrm{sc}}$. We get our $h_{\mathbb{R}}$ by working with the spin representation of $\tilde{G}_{\mathbb{C}}^{\mathrm{sc}}$ i.e., by composing the homomorphism $\tilde{h}_{\mathbb{R}} : \tilde{G}_{\mathbb{R}}^{\mathrm{sc}} \rightarrow \mathbf{SU}(2^{n-2}, 2^{n-2})_{\mathbb{R}} \times_{\mathbb{R}} \mathbf{SU}(2^{n-2}, 2^{n-2})_{\mathbb{R}}$ defined by the mentioned two embeddings with a standard monomorphism

$$\Delta : \mathbf{SU}(2^{n-2}, 2^{n-2})_{\mathbb{R}} \times_{\mathbb{R}} \mathbf{SU}(2^{n-2}, 2^{n-2})_{\mathbb{R}} \hookrightarrow \mathbf{SU}(2^{n-1}, 2^{n-1})_{\mathbb{R}}.$$

It is easy to see that Δ is uniquely determined up to $\mathbf{SU}(2^{n-1}, 2^{n-1})_{\mathbb{R}}(\mathbb{R})$ -conjugation.

Let V_A be a complex vector space such that $G(A)_{\mathbb{C}}$ is (isomorphic to) $\mathbf{GL}_{V_A}^{\mathrm{der}}$. If (G, X) is of C_n or $D_n^{\mathbb{H}}$ (resp. B_n or $D_n^{\mathbb{R}}$) type, then the representation of $\tilde{G}_{\mathbb{C}}^{\mathrm{sc}}$ on V_A is the standard (resp. is the spin) representation of dimension $2n$ (resp. 2^n).

4.3.1. Simple properties. (a) We assume that (G, X) is of $D_n^{\mathbb{R}}$ type. Thus $\tilde{G}^{\mathrm{sc}}$ is $\mathbf{Spin}(2, 2n-2)$. We check that $h_{\mathbb{R}}$ is the composite $h_{\mathbb{R}}^{(1)} : \mathbf{Spin}(2, 2n-2)_{\mathbb{R}} \rightarrow G(A)_{\mathbb{R}}$ of a standard monomorphism $\mathbf{Spin}(2, 2n-2)_{\mathbb{R}} \hookrightarrow \mathbf{Spin}(2, 2n-1)_{\mathbb{R}}$ with the monomorphism $\mathbf{Spin}(2, 2n-1)_{\mathbb{R}} \hookrightarrow G(A)_{\mathbb{R}}$ we mentioned above but for the B_n type case. The representation of $\tilde{G}_{\mathbb{C}}^{\mathrm{sc}}$ on V_A is the spin representation, regardless if this representation is obtained via $h_{\mathbb{R}}$ or via $h_{\mathbb{R}}^{(1)}$. Thus the double centralizer of $\mathrm{Im}(h_{\mathbb{R}}^{(1)})$ in $G(A)_{\mathbb{R}}$ has a derived group D whose extension to $\mathbb{C}$ is isomorphic to $\mathbf{SL}_{2^{n-1}, \mathbb{C}}^2$. As $h_{\mathbb{R}}^{(1)}$ satisfies at the level of

Lie algebras the condition (H_2) and as $\text{Im}(h_{\mathbb{R}}^{(1)}) \leq D \leq G(A)_{\mathbb{R}}$, D has a compact torus of the same rank as D . Thus each simple factor of D^{ad} is absolutely simple and of the form $\mathbf{SU}(2^{n-1} - a, a)_{\mathbb{R}}^{\text{ad}}$ for some $a \in \{0, \dots, 2^{n-1}\}$ (see [He, Ch. X, Sect. 6, Table V]). The homomorphism $\tilde{G}_{\mathbb{R}}^{\text{sc}} \rightarrow D$ satisfies at the level of Lie algebras the condition (H_2) (as $h_{\mathbb{R}}^{(1)}$ has this property). Thus from [Sa1, Subsect. 3.3] we get that D is a product of two copies of $\mathbf{SU}(2^{n-2}, 2^{n-2})_{\mathbb{R}}$ (i.e., a is 2^{n-2}) and that the homomorphism $\tilde{G}_{\mathbb{R}}^{\text{sc}} \rightarrow D$ can be identified with $\tilde{h}_{\mathbb{R}}$ of Subsection 4.3. Also Δ can be identified with the monomorphism $D \hookrightarrow G(A)_{\mathbb{R}}$. We conclude that $h_{\mathbb{R}}$ can be identified with $h_{\mathbb{R}}^{(1)}$.

(b) The finite group $\text{Ker}(h_{\mathbb{R}})$ is trivial if and only if (G, X) is of B_n , C_n , or $D_n^{\mathbb{R}}$ type (cf. the description of the $\tilde{G}_{\mathbb{C}}^{\text{sc}}$ -module V_A).

(c) From now on we take $h_{\mathbb{R}}$ such that it is the extension to $\mathbb{R}$ of a homomorphism

$$h: \tilde{G}^{\text{sc}} \rightarrow G(A)$$

over $\mathbb{Q}$ (thus the notation $h_{\mathbb{R}}$ is justified). The fact that such a choice of $h_{\mathbb{R}}$ is possible follows from the constructions in [Sa1, Subsects. 3.3 to 3.7]: one only has to take Δ to be defined over $\mathbb{Q}$ and to replace in loc. cit. $\mathbb{R}$, $\mathbb{C}$, and $\mathbb{H}$ by $\mathbb{Q}$, $\mathbb{Q}(i)$, and respectively by the standard quaternion algebra $\mathbb{Q}(i, j)$ over $\mathbb{Q}$. Let

$$\tilde{G}^d := \tilde{G}^{\text{sc}} / \text{Ker}(h).$$

4.3.2. Fact. *Let K be a field of characteristic 0. Let $\tilde{G}_K^d \hookrightarrow G(A)_K$ be the monomorphism defined by $h_K: \tilde{G}_K^{\text{sc}} \rightarrow G(A)_K$. Then the centralizer of a maximal torus of $\tilde{G}_K^d$ in $G(A)_K$, is a maximal torus of $G(A)_K$.*

Proof: We can assume that K is $\mathbb{C}$. Each such centralizer is a reductive group, cf. [Bor, Ch. IV, 13.17, Cor. 2 (a)]. Thus to prove the Fact, it suffices to show that the centralizer of a maximal torus $\tilde{T}$ of $\tilde{G}_{\mathbb{C}}^d$ in $\mathbf{GL}_{V_A}$ is a torus. It suffices to check that all weight spaces of the representation of $\tilde{T}$ on V_A have dimension 1. The fact that this holds is a basic property of minuscule weights (cf. [Bou2, pp. 127–129]) but for readers' convenience we recall one simple way to argue it.

If $(\tilde{G}, \tilde{X})$ is of C_n (resp. $D_n^{\mathbb{H}}$) type, then the monomorphism $\tilde{G}_{\mathbb{C}}^d \hookrightarrow \mathbf{GL}_{V_A}$ is a standard $\mathbf{Sp}_{2n, \mathbb{C}} \hookrightarrow \mathbf{GL}_{2n, \mathbb{C}}$ (resp. $\mathbf{SO}_{2n, \mathbb{C}} \hookrightarrow \mathbf{GL}_{2n, \mathbb{C}}$) monomorphism and it is well known that all the mentioned weight spaces have dimension 1. If $(\tilde{G}, \tilde{X})$ is of B_n or $D_n^{\mathbb{R}}$ type, then the representation of $\tilde{G}_{\mathbb{C}}^d$ on V_A is a spin representation and thus (cf. [FH, Props. 20.15 and 20.20]) all the mentioned weight spaces have dimension 1. $\square$

4.4. Lemma. *There exists a totally real finite field extension F_1 of F of degree a divisor of 2 and such that the group $G[F]_{F_1}$ is an inner form of $\tilde{G}_{F_1}$.*

Proof: The group $G[F]$ is a form of $\tilde{G}_F$. If (G, X) is of B_n or C_n type, then this form is inner and thus we can take $F_1 = F$. Let now (G, X) be of $D_n^{\mathbb{H}}$ or $D_n^{\mathbb{R}}$ type ($n \geq 4$). Let $\text{Aut}(\tilde{G}_F)$ be the group scheme of automorphisms of $\tilde{G}_F$. The group $\tilde{G}_F$ is a normal subgroup of $\text{Aut}(\tilde{G}_F)$ and the quotient group $Q := \text{Aut}(\tilde{G}_F) / \tilde{G}_F$ is $\mathbb{Z}/2\mathbb{Z}$ if $n \geq 5$ and is the symmetric group S_3 if $n = 4$. We have a short exact sequence $1 \rightarrow Q_1 \rightarrow Q \rightarrow \mathbb{Z}/2\mathbb{Z} \rightarrow 1$,

where Q_1 is $\mathbb{Z}/3\mathbb{Z}$ if $n = 4$ and is the trivial group if $n > 5$. Let $Aut_1(\tilde{G}_F)$ be the normal subgroup of $Aut(\tilde{G}_F)$ which contains $\tilde{G}_F$ and whose group of connected components is Q_1 .

Let $\delta_0 \in H^1(F, Aut(\tilde{G}_F))$ be the class that defines the form $G[F]$ of $\tilde{G}_F$. In the étale topology of $\text{Spec}(F)$, the non-trivial torsors of $\mathbb{Z}/2\mathbb{Z}$ correspond to quadratic extensions of F . For each embedding $F \hookrightarrow \mathbb{R}$, the group $G[F]_{\mathbb{R}}$ is either $\tilde{G}_{\mathbb{R}}$ or the compact form of $\tilde{G}_{\mathbb{R}}$; thus $G[F]_{\mathbb{R}}$ is an inner form of $\tilde{G}_{\mathbb{R}}$ (cf. [De2, Subsubsection. 2.3.4]). This implies that there exists a smallest totally real finite field extension F_1 of F of degree a divisor of 2 and such that the image of δ_0 in $H^1(F_1, Aut(\tilde{G}_F)_{F_1})$ is the image of a class $\delta_1 \in H^1(F_1, Aut_1(\tilde{G}_F)_{F_1})$. The Shimura pairs (G, X) and (G^{F_1}, X^{F_1}) have the same type that is either $D_n^{\mathbb{H}}$ or $D_n^{\mathbb{R}}$, cf. Subsubsection 2.2.3. If $n = 4$, then from the very definition of the $D_4^{\mathbb{R}}$ and $D_4^{\mathbb{H}}$ types we get that the image of δ_1 in $H^1(F_1, Q_1)$ is trivial. We conclude that for all $n \geq 4$, δ_1 is the image of a class in $H^1(F_1, \tilde{G}_{F_1})$. Thus $G[F]_{F_1}$ is an inner form of $\tilde{G}_{F_1}$ for all $n \geq 4$. $\square$

4.5. The twisting process. Let $\delta \in H^1(F_1, \tilde{G}_{F_1})$ be the class that defines the inner form $G[F]_{F_1}$ of $\tilde{G}_{F_1}$. Let $\tilde{G}^e$ be the image of $\tilde{G}^d$ in $G(A)^{\text{ad}}$. If (G, X) is not of $D_n^{\mathbb{R}}$ type, then the representation of $\tilde{G}_{\mathbb{C}}^d$ on V_A is irreducible and this implies that $Z(\tilde{G}^d)$ is a subgroup of $Z(G(A))$; thus $\tilde{G}^e$ is an adjoint group.

In this paragraph we assume that (G, X) is of $D_n^{\mathbb{R}}$ type. The group $\tilde{G}_{\mathbb{R}}^e$ is the image of $\mathbf{Spin}(2, 2n-2)_{\mathbb{R}}$ in $\mathbf{SO}(2, 2n-1)_{\mathbb{R}}$ (cf. property 4.3.1 (a)) and thus it is $\mathbf{SO}(2, 2n-2)_{\mathbb{R}}$. This implies that $\tilde{G}_e$ is $\mathbf{SO}(2, 2n-2)$ and thus $\mu_{2, \mathbb{Q}} \xrightarrow{\sim} \ker(\tilde{G}^d \rightarrow \tilde{G}^e)$. The compact inner form $\mathbf{SO}(2n)_{\mathbb{R}}$ of $\mathbf{SO}(2, 2n-2)_{\mathbb{R}}$ is obtained by twisting through a class in the set $H^1(\mathbb{R}, \mathbf{SO}(2, 2n-2)_{\mathbb{R}})$. Thus the obstruction of lifting δ to a class $\delta_e \in H^1(F_1, \tilde{G}_{F_1}^e)$, is measured by a class $\delta_2 \in H^2(F_1, \mu_{2, F_1})$ whose image in $H^2(\mathbb{R}, \mu_{2, \mathbb{R}})$ under each embedding $F_1 \hookrightarrow \mathbb{R}$ is trivial. Let $\mathcal{D}_2$ be the division algebra of center F_1 that corresponds to δ_2 . Based on [Ha, Lem. 5.5.3] applied to the reductive group $\mathbf{GL}_1(\mathcal{D}_2)$, we get the existence of a totally real finite field extension F_2 of F_1 which has a degree that divides 2 and which is a maximal commutative F_1 -subalgebra of $\mathcal{D}_2$. By replacing F_1 with F_2 we can assume that δ_2 is the trivial class.

Thus, regardless of the type of (G, X) , by requesting only that the degree $[F_1 : F]$ divides 4 we can assume that δ is the image of a class $\delta_e \in H^1(F_1, \tilde{G}_{F_1}^e)$. Let $h_e : \tilde{G}^e \hookrightarrow G(A)^{\text{ad}}$ be the monomorphism defined by h . We use δ_e to twist h_{e, F_1} . By performing such a twist and by lifting the resulting monomorphism via isogenies, we get a monomorphism

$$h_{F_1}^t : G[F]_{F_1}^d \hookrightarrow G(A)_{F_1}^t.$$

Here $G[F]_{F_1}^d$ is the isogeny cover of $G[F]_{F_1}$ that corresponds to the isogeny cover $\tilde{G}^d$ of $\tilde{G}$ and $G(A)_{F_1}^t$ is the resulting inner form of $G(A)_{F_1}$ (the upper index t stands for twist).

4.6. Lemma. *If by extension of scalars under an embedding $F_1 \hookrightarrow \mathbb{R}$, the group $G[F]_{F_1}$ becomes a compact group (resp. it becomes $\tilde{G}_{\mathbb{R}}$), then under the same extension of scalars $G(A)_{F_1}^t$ becomes a compact group (resp. it becomes $G(A)_{\mathbb{R}}$).*

Proof: We have to prove that if we twist $h_{e, \mathbb{R}} : \tilde{G}_{\mathbb{R}}^e \hookrightarrow G(A)_{\mathbb{R}}^{\text{ad}}$ through the compact inner form of $\tilde{G}_{\mathbb{R}}^e$, then the inner form J^{ad} of $G(A)_{\mathbb{R}}^{\text{ad}}$ we get is the compact form. This can be proved using Cartan decompositions (see [He]). Here we include a proof of this that relies

on Lemma 1.1.7. Based on the property 4.3.1 (a), the case of $D_n^{\mathbb{R}}$ type gets reduced to the B_n type case. Thus it suffices to consider the cases when (G, X) is of B_n , C_n , or $D_n^{\mathbb{H}}$ type. Let $\mathcal{J}$ be the simply connected semisimple group cover of $\mathcal{J}^{\text{ad}}$.

Let $\mathcal{C}$ be a maximal compact subgroup of $\mathcal{J}$. Let V_A be as before Subsubsection 4.3.1. We identify $\mathcal{J}_{\mathbb{C}} = \mathbf{GL}_{V_A}^{\text{der}}$. If $T_{\mathbb{R}}$ is a maximal compact torus of $\tilde{G}_{\mathbb{R}}^e$, then the map $H^1(\mathbb{R}, T_{\mathbb{R}}) \rightarrow H^1(\mathbb{R}, \tilde{G}_{\mathbb{R}}^e)$ is surjective (see [Ko1, Lem. 10.2]). Thus, as $T_{\mathbb{R}}$ is contained in a maximal compact torus $T(A)$ of $G(A)_{\mathbb{R}}^{\text{ad}}$, we get that $T(A)$ is isomorphic to a torus of $\mathcal{J}$. As the maximal compact Lie subgroups of $\mathcal{J}(\mathbb{R})$ are $\mathcal{J}(\mathbb{R})$ -conjugate (see [He, Ch. VI, Sect. 2, Thm. 2.2 (ii)]), $\mathcal{C}$ has a subgroup isomorphic to $T(A)$. Thus $\mathcal{C}$ has the same rank as $\mathcal{J}$.

The representation of $\tilde{G}_{\mathbb{C}}^d$ on V_A is irreducible. Thus the $\tilde{G}_{\mathbb{C}}^d$ -module V_A is simple. But up to $\mathcal{J}(\mathbb{C})$ -conjugation, we can assume that $\tilde{G}_{\mathbb{C}}^d$ is a subgroup of $\mathcal{C}_{\mathbb{C}}$. Thus the $\mathcal{C}_{\mathbb{C}}$ -module V_A is also simple. From Lemma 1.1.7 applied to the inclusions $\text{Lie}(\mathcal{C}_{\mathbb{C}}) \hookrightarrow \text{Lie}(\mathcal{J}_{\mathbb{C}}) \hookrightarrow \text{End}(V_A)$, we get that $\text{Lie}(\mathcal{J}_{\mathbb{C}}) = \text{Lie}(\mathcal{C}_{\mathbb{C}})$. Thus $\mathcal{J} = \mathcal{C}$ i.e., the group $\mathcal{J}$ is compact. $\square$

4.7. The Shimura pair (G_1, X_1) . Let G_1 be the adjoint group of $G_1^{\text{sc}} := \text{Res}_{F_1/\mathbb{Q}} G(A)_{F_1}^t$. Each simple factor of $G_{1, \mathbb{R}}$ is either compact or an $\mathbf{SU}(a, a)_{\mathbb{R}}^{\text{ad}}$ group for some $a \in \mathbb{N}$, cf. Lemma 4.6. This implies the existence of an adjoint Shimura pair of the form (G_1, X_1) ; it is uniquely determined (cf. [De2, Prop. 1.2.7 and Cor. 1.2.8 (ii)]).

4.8. Theorem. *We recall that (G, X) is a simple, adjoint Shimura pair of B_n , C_n , $D_n^{\mathbb{H}}$, or $D_n^{\mathbb{R}}$ type and that F_1 is a totally real finite field extension of F of degree a divisor of 4. There exist injective maps*

$$f_2 : (G_2, X_2) \hookrightarrow (G_3, X_3) \hookrightarrow (G_4, X_4) \xrightarrow{f_4} (\mathbf{GSp}(W_1, \psi_1), S_1)$$

of Shimura pairs such that the following seven properties hold:

- (i) we have $(G_2^{\text{ad}}, X_2^{\text{ad}}) = (G, X)$, $(G_3^{\text{ad}}, X_3^{\text{ad}}) = (G^{F_1}, X^{F_1})$, $(G_4^{\text{ad}}, X_4^{\text{ad}}) = (G_1, X_1)$, and $G_4^{\text{der}} = G_1^{\text{sc}}$;
- (ii) we have a natural monomorphism $G_2^{\text{ad}} \hookrightarrow G_3^{\text{ad}}$ which under the identifications of (i) is nothing else but the natural monomorphism $G \hookrightarrow G^{F_1}$;
- (iii) we have $G_3^{\text{der}} = \text{Res}_{F_1/\mathbb{Q}} G[F]_{F_1}^d$ and under this identification and of the ones of (i), the monomorphism $G_3^{\text{der}} \hookrightarrow G_4^{\text{der}}$ becomes the $\text{Res}_{F_1/\mathbb{Q}}$ of $h_{F_1}^t$ (of Subsection 4.5);
- (iv) if (G, X) is not (resp. is) of $D_n^{\mathbb{H}}$ type, then we have $E(G_2, X_2) = E(G_3, X_3) = E(G_4, X_4)$ (resp. we have $E(G_2, X_2) = E(G_3, X_3)$);
- (v) both $Z^0(G_2)$ and $Z^0(G_3)$ are subtori of $Z^0(G_4)$;
- (vi) if (G, X) is not of $D_n^{\mathbb{H}}$ type, then both G_2^{der} and G_3^{der} are simply connected;
- (vii) the group G_4 is the subgroup of $\mathbf{GSp}(W_1, \psi_1)$ that fixes all elements of the semisimple $\mathbb{Q}$ -algebra $\mathcal{B} := \text{End}(W_1)^{G_4(\mathbb{Q})}$ (thus f_4 is a PEL type embedding).

Proof: Let $f_4 : (G_4, X_4) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ be obtained as in Proposition 4.1 but working with (G_1, X_1) (instead of (G, X)). Thus (vii) holds (cf. property 4.1 (ii)) and we have $G_4^{\text{der}} = G_1^{\text{sc}}$ and $(G_4^{\text{ad}}, X_4^{\text{ad}}) = (G_1, X_1)$. We take G_3 to be generated by $Z^0(G_4)$ and by the semisimple subgroup $\text{Res}_{F_1/\mathbb{Q}} G[F]_{F_1}^d$ of G_4^{der} . This takes care of (iii) and

we have $Z^0(G_3) = Z^0(G_4)$. We take G_2 to be the reductive subgroup of G_3 such that $Z^0(G_2) = Z^0(G_4)$ and (ii) holds. Thus (v) also holds.

The monomorphism $G_{3,\mathbb{R}}^{\text{der}} \hookrightarrow G_{4,\mathbb{R}}^{\text{der}}$ is a product of $[F_1 : \mathbb{Q}]$ monomorphisms that can be identified either with the monomorphism $\tilde{G}_{\mathbb{R}}^d \hookrightarrow G(A)_{\mathbb{R}}$ defined naturally by $h_{\mathbb{R}}$ or with a monomorphism between compact groups, cf. Lemma 4.6. Thus from (3) and the uniqueness part of Subsection 4.7, we get the existence of elements $x \in X$ and $x_1 \in X_1$ such that the image of $dx(\sqrt{-1}) \in \text{Lie}(G_{\mathbb{R}})$ in $\text{Lie}(G_{1,\mathbb{R}})$ is $dx_1(\sqrt{-1}) \in \text{Lie}(G_{1,\mathbb{R}})$. But $G_4^{\text{ad}}(\mathbb{Q})$ is dense in $G_4^{\text{ad}}(\mathbb{R})$ and therefore it permutes transitively the connected components of $X_1 = X_4^{\text{ad}}$. Thus by composing the monomorphism $G_4 \hookrightarrow \mathbf{GSp}(W_1, \psi_1)$ with an isomorphism $G_4 \xrightarrow{\sim} G_4$ defined by an element of $G_4^{\text{ad}}(\mathbb{Q})$, we can choose x and x_1 such that x_1 is the image in $X_1 = X_4^{\text{ad}}$ of a point $x_4 \in X_4$. Therefore the monomorphism $x_4 : \mathbb{S} \hookrightarrow G_{4,\mathbb{R}}$ factors through $G_{2,\mathbb{R}}$ in such a way that the resulting homomorphism $\mathbb{S} \rightarrow G_{2,\mathbb{R}}^{\text{ad}} = G_{\mathbb{R}}$ is an element of X . Thus by taking X_2 (resp. X_3) to be the $G_2(\mathbb{R})$ -conjugacy (resp. the $G_3(\mathbb{R})$ -conjugacy) class of the factorization of x_4 through $G_{2,\mathbb{R}}$ (resp. through $G_{3,\mathbb{R}}$), we get that (i) holds.

To check (iv), let $* \in \{2, 3, 4\}$. The reflex field of $E(G_*, X_*)$ is the composite field of $E(G_*^{\text{ad}}, X_*^{\text{ad}})$ and $E(G_*^{\text{ab}}, X_*^{\text{ab}})$, cf. Subsection 2.2. But $(G_*^{\text{ab}}, X_*^{\text{ab}})$ and thus also $E(G_*^{\text{ab}}, X_*^{\text{ab}})$ does not depend on $*$. Moreover, we have $E(G_2^{\text{ad}}, X_2^{\text{ad}}) = E(G_3^{\text{ad}}, X_3^{\text{ad}})$ (cf. Subsubsection 2.2.3). We now assume that (G, X) is not of $D_n^{\mathbb{H}}$ type. Let $(\mathfrak{D}_*, \mathfrak{n}_{X_*})$ be the analogue of $(\mathfrak{D}, \mathfrak{n}_X)$ of Subsection 2.2 but for the Shimura pair $E(G_*^{\text{ad}}, X_*^{\text{ad}})$. We recall from [De2, Rm. 2.3.12] that $E(G_3^{\text{ad}}, X_3^{\text{ad}})$ (resp. $E(G_4^{\text{ad}}, X_4^{\text{ad}})$) is the fixed field of the subgroup of $\text{Gal}(\mathbb{Q})$ that stabilizes the union of the connected components of $\mathfrak{D}_3$ (resp. of $\mathfrak{D}_4$) that correspond to compact factors of $G_{3,\mathbb{R}}^{\text{ad}}$ (resp. of $G_{4,\mathbb{R}}^{\text{ad}}$). From this and Lemma 4.6 we get $E(G_3^{\text{ad}}, X_3^{\text{ad}}) = E(G_4^{\text{ad}}, X_4^{\text{ad}})$. Thus (iv) holds.

Condition (vi) follows from the construction of $h_{F_1}^t$ and the property 4.3.1 (b). $\square$

4.8.1. Remark. We assume that $n \in 2 + 2\mathbb{N}$ and that (G, X) is of inner $D_n^{\mathbb{R}}$ type. Then we have a variant of Subsections 4.3 to 4.8 in which we work with only one fixed half spin representation of $\tilde{G}_{\mathbb{C}}^{\text{sc}}$. Thus the group $G(A)$ becomes $\mathbf{SU}(2^{n-2}, 2^{n-2})$, we have $[F_1 : F] \leq 2$ (i.e., in Lemma 4.4 we have $F_1 = F$ and in Subsection 4.5 we do not have to replace F_1 by F_2), and the groups $G_{2,\overline{\mathbb{Q}}}^{\text{der}}$ and $G_{3,\overline{\mathbb{Q}}}^{\text{der}}$ are products of $\mathbf{Spin}_{2n,\overline{\mathbb{Q}}}/\mu_{2,\overline{\mathbb{Q}}}$ groups which for $n \neq 4$ are not $\mathbf{SO}_{2n,\overline{\mathbb{Q}}}$ groups.

Also if (G, X) is of non-inner $D_4^{\mathbb{R}}$ type, then we have a variant of Subsections 4.3 to 4.8 in which we construct $h : \tilde{G}^{\text{sc}} \rightarrow G(A)$ as for the $D_4^{\mathbb{H}}$ type. Thus $G(A)$ is $\mathbf{SU}(4, 4)$, $F_1 \neq F$, and due to the non-inner assumption the groups $G_{2,\overline{\mathbb{Q}}}^{\text{der}}$ and $G_{3,\overline{\mathbb{Q}}}^{\text{der}}$ are products of $\mathbf{Spin}_{8,\overline{\mathbb{Q}}}$ groups.

4.8.2. Remarks. (a) In the proof of Proposition 4.1, the degree $[F_0 : \mathbb{Q}]$ is bounded in terms of $\dim(T_{\mathbb{Q}})$ and thus also of the rank of $G_{\mathbb{C}}$. Moreover, in Theorem 4.8 we have $[F_1 : F] \leq 4$. Thus there exists an effectively computable number $M(G, X) \in \mathbb{N}$ such that in Proposition 4.1 (resp. Theorem 4.8), we can take the injective map $f_2 : (G_2, X_2) \hookrightarrow (\mathbf{GSp}(W, \psi), S)$ (resp. $f_2 : (G_2, X_2) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$) of Shimura pairs in such a way that the following inequality $\dim_{\mathbb{Q}}(W_1) \leq M(G, X)$ holds.

(b) We assume that (G, X) is of $D_n^{\mathbb{H}}$ type. We denote also by $h : \tilde{G}^d \hookrightarrow G(A)$ the

monomorphism defined by h . Then there exist monomorphisms over $\mathbb{R}$

$$(4) \quad \tilde{G}_{\mathbb{R}}^d \xrightarrow{h_{\mathbb{R}}} G(A)_{\mathbb{R}} \xrightarrow{h_{1,\mathbb{R}}} \mathbf{Sp}_{4n,\mathbb{R}} \xrightarrow{h_{2,\mathbb{R}}} \mathbf{SU}(2n, 2n)_{\mathbb{R}}$$

which at the level of Lie algebras satisfy the condition (H_2) . Argument: see [Sa1, Subsect. 1.4] for $h_{1,\mathbb{R}}$; as $h_{2,\mathbb{R}}$ we can take the analogue of $h_{\mathbb{R}}$ but for the C_{2n} type.

By reasons of ranks of maximal compact subgroups, $\mathbf{SU}(2n, 0)_{\mathbb{R}}$ is not a subgroup of a non-compact form of $\mathbf{Sp}_{4n,\mathbb{R}}$. Thus Lemma 4.6 also holds in the context of $h_{1,\mathbb{R}}$. As in the property 4.3.1 (c) we argue that (4) is the extension to $\mathbb{R}$ of monomorphisms h , h_1 , and h_2 over $\mathbb{Q}$. Therefore Subsections 4.3 to 4.8 can be adapted in the context of the monomorphisms h , h_1 , and h_2 to get that we can choose f_2 and f_4 such that the injective map $f_4 : (G_4, X_4) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ of Shimura pairs factors as injective maps $(G_4, X_4) \hookrightarrow (G_5, X_5) \hookrightarrow (G_6, X_6) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ of Shimura pairs and moreover, besides properties 4.8 (i) to (v), we also have the following four extra properties:

- (i) the group G_6 is the subgroup of $\mathbf{GSp}(W_1, \psi_1)$ that centralizes the semisimple $\mathbb{Q}$ -algebra $\mathcal{B}_6 := \text{End}(W_1)^{G_6(\mathbb{Q})}$;
- (ii) the monomorphisms $G_3^{\text{der}} \hookrightarrow G_4^{\text{der}}$, $G_4^{\text{der}} \hookrightarrow G_5^{\text{der}}$, and $G_5^{\text{der}} \hookrightarrow G_6^{\text{der}}$ are $\text{Res}_{F_1/\mathbb{Q}}$ of inner twists of the monomorphisms h_{F_1} , h_{1,F_1} , and h_{2,F_1} (respectively);
- (iii) the torus $Z^0(G_*)$ does not depend on $* \in \{2, \dots, 6\}$;
- (iv) the group G_3 is the identity component of the subgroup of G_5 that centralizes the semisimple $\mathbb{Q}$ -algebra $\mathcal{B}_3 := \text{End}(W_1)^{G_3(\mathbb{Q})}$.

Property (iv) is an easy consequence of (iii). As $G_{5,\overline{\mathbb{Q}}}^{\text{der}}$ is a product of $\mathbf{Sp}_{4n,\overline{\mathbb{Q}}}$ groups, the relative PEL property expressed by (iv) is a fundamental tool in showing that many properties enjoyed by (G_5, X_5) are as well enjoyed by (G_3, X_3) and thus implicitly by (G_2, X_2) (we have in mind mainly applications to the Langlands–Rapoport conjecture mentioned in Subsubsect. 1.2.1). In other words, this relative PEL type property “compensates” the fact that G_2^{der} and G_3^{der} are not simply connected.

(c) We assume that (G, X) is of $D_n^{\mathbb{R}}$ type. The monomorphism $G_{3,\overline{\mathbb{Q}}}^{\text{der}} \hookrightarrow G_{4,\overline{\mathbb{Q}}}^{\text{der}}$ is a product of $[F_1 : \mathbb{Q}]$ copies of the spin (representation) monomorphism $\mathbf{Spin}_{2n,\overline{\mathbb{Q}}} \hookrightarrow \mathbf{SL}_{2n,\overline{\mathbb{Q}}}$. Thus the identity component Z_3 of the centralizer of G_3^{der} in G_4^{der} is a torus of rank $[F_1 : \mathbb{Q}]$. Let $G_{3.5}$ be the reductive subgroup of G_4 generated by G_3 and Z_3 . Let $G_{3.75}$ be the reductive subgroup of $\mathbf{GSp}(W_1, \psi_1)$ that centralizes $\text{End}(W_1)^{G_3(\mathbb{Q})}$. We have monomorphisms $G_{3.5} \hookrightarrow G_{3.75} \hookrightarrow G_4$ which give birth to an identity $Z^0(G_{3.5}) = Z^0(G_{3.75})$. Moreover, the group $G_{3.75,\overline{\mathbb{Q}}}^{\text{der}}$ is isomorphic to $\mathbf{SL}_{2n-1,\overline{\mathbb{Q}}}^{2[F_1:\mathbb{Q}]}$.

4.8.3. Special cases. We refer to Theorem 4.8. All simple factors of $G_{1,\mathbb{R}}$ are either compact or are $\mathbf{SU}(a, a)_{\mathbb{R}}$ groups, where $a \in \mathbb{N}$. Thus (G_1, X_1) is without involution. For the rest of this Subsubsection we will assume that all simple factors of $G_{\mathbb{R}}$ are non-compact. From Lemma 4.6 we get that all simple factors of $G_{1,\mathbb{R}}$ and $G_{2,\mathbb{R}}^{\text{ad}}$ are also non-compact. Thus the homomorphism $x_4 : \mathbb{S} \rightarrow G_{4,\mathbb{R}}$ of the proof of Theorem 4.8 factors through the extension to $\mathbb{R}$ of the subgroup of G_4 generated by G_4^{der} and by $Z(\mathbf{GL}_{W_1})$, cf. property 4.1.1 (b). Thus we can choose f_2 such that we have $Z^0(G_2) = Z^0(G_4) = Z(\mathbf{GL}_{W_1}) \xrightarrow{\sim} \mathbb{G}_{m,\mathbb{Q}}$.

In this paragraph we take (G, X) to be of either C_n or $D_n^{\mathbb{H}}$ type. The double centralizer D_2 of G_2 in $\mathbf{GL}_{W_1}$ is a reductive group whose extension to $\overline{\mathbb{Q}}$ is a product of $[F : \mathbb{Q}]$ copies of $\mathbf{GL}_{2n, \overline{\mathbb{F}}}$. The involution of $\text{End}(W_1 \otimes_{\mathbb{Q}} \overline{\mathbb{Q}})$ induces a non-trivial involution of the Lie algebra of each such copy of $\mathbf{GL}_{2n, \overline{\mathbb{F}}}$. This implies that $\text{Lie}(D_2 \cap \mathbf{GSp}(W_1, \psi_1))$ is $\text{Lie}(G_2)$. Thus G_2 is the identity component of D_2 and therefore the injective map $f_2 : (G_2, X_2) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ of Shimura pairs is a PEL type embedding.

In this paragraph we take (G, X) to be of inner $D_4^{\mathbb{R}}$ type. If we work with the variant of Theorem 4.8 mentioned in Remark 4.8.1, then similar arguments show that we can assume that $Z^0(G_2) = Z^0(G_4) = Z(\mathbf{GL}_{W_1}) \simeq \mathbb{G}_{m, \mathbb{Q}}$ and that $f_2 : (G_2, X_2) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ is a PEL type embedding.

4.9. Variants. Using Fact 2.4.1 (b) we get that Proposition 4.1 and Theorem 4.8 have variants for the case when (G, X) is an arbitrary Shimura pair of adjoint, abelian type. To state such variants, let $(G, X) = \prod_{i \in I} (G_i, X_i)$ be the product decomposition into simple, adjoint Shimura pairs indexed by a set I which is disjoint from $\mathbb{Z}$. For $i \in I$ let $F_i := F(G_i, X_i)$ and $\tilde{G}_i := G_i[F_i]$, cf. Subsection 2.2.

(a) There exists a product $(G_1, X_1) = \prod_{i \in I} (G_{i,1}, X_{i,1})$ of simple, adjoint, unitary Shimura pairs and there exist injective maps

$$f_2 : (G_2, X_2) \hookrightarrow (G_3, X_3) \hookrightarrow (G_4, X_4) \xrightarrow{f_4} (\mathbf{GSp}(W_1, \psi_1), S_1)$$

of Shimura pairs such that the following seven properties hold:

(i) we have $(G_2^{\text{ad}}, X_2^{\text{ad}}) = (G, X)$ and $(G_3^{\text{ad}}, X_3^{\text{ad}}) = \prod_{i \in I} (G_i^{F_{i,1}}, X_i^{F_{i,1}})$, with $F_{i,1}$ as a totally real number field that contains F_i ;

(ii) we get a monomorphism $G_2^{\text{ad}} \hookrightarrow G_3^{\text{ad}}$ that is a product indexed by $i \in I$ of the natural monomorphisms $G_i \hookrightarrow G_i^{F_{i,1}}$;

(iii) the monomorphism $G_3^{\text{der}} \hookrightarrow G_4^{\text{der}}$ is a product indexed by $i \in I$ of monomorphisms that are either isomorphisms or are Weil restrictions of monomorphisms obtained as $h_{F_1}^t$ in Subsection 4.5, depending on the fact that (G_i, X_i) is or is not of A_{n_i} type for some $n_i \in \mathbb{N}$;

(iv) we have $(G_4^{\text{ad}}, X_4^{\text{ad}}) = (G_1, X_1)$, the group G_4^{der} is a simply connected semisimple group, and f_4 is a PEL type embedding that is a Hodge quasi product indexed naturally by elements of I of PEL type embeddings (cf. Fact 2.4.1 (b));

(v) if for all $i \in I$ (resp. if there exists $i \in I$ such that) the Shimura pair (G_i, X_i) is not (resp. is) of $D_n^{\mathbb{H}}$ type, then $E(G_2, X_2) = E(G_3, X_3) = E(G_4, X_4)$ (resp. $E(G_2, X_2) = E(G_3, X_3)$); also if for all $i \in I$ the pair (G_i, X_i) is of some A_{n_i} type, then $G_2 = G_3 = G_4$;

(vi) both $Z^0(G_2)$ and $Z^0(G_3)$ are subtori of $Z^0(G_4)$;

(vii) if none of the Shimura pairs (G_i, X_i) (with $i \in I$) is of some $D_n^{\mathbb{H}}$ type, then G_2^{der} and G_3^{der} are simply connected semisimple groups.

(b) In Theorem 4.8 we can replace F_1 by any other totally real number field F_2 that contains it. If there is $i \in I$ such that the simple, adjoint Shimura pair (G_i, X_i) is of A_n type, then for each totally real number field extension $F_{i,2}$ of $F_{i,1}$, the simple, adjoint

Shimura pair $(G_i^{F_{i,2}}, X_i^{F_{i,2}})$ is also of A_n type and therefore one can apply Proposition 4.1 to it. Thus, as in the proof of Theorem 4.8 we argue that for such an element $i \in I$ there exist injective maps $f_{i,2} : (G_{i,2}, X_{i,2}) = (G_{i,3}, X_{i,3}) \hookrightarrow (G_{i,4}, X_{i,4}) \xrightarrow{f_{i,4}} (\mathbf{GSp}(W_{i,1}, \psi_{i,1}), S_{i,1})$ of Shimura pairs such that the analogues of properties 4.8 (i) to (vii) hold for them (therefore $(G_{i,2}^{\text{ad}}, X_{i,2}^{\text{ad}}) = (G_i, X_i)$, $(G_{i,4}, X_{i,4}) = (G_i^{F_{i,2}}, X_i^{F_{i,2}})$, $f_{i,4}$ is a PEL type embedding, etc.)

Based on the previous paragraph, by using composites of fields and injective maps as in (1) we get an uniformized variant of (a) in which to the property (iii) of (a) we can add the requirement that the field $F_{i,1}$ does not depend on $i \in I$ (thus in general we have no upper bounds for the degrees $[F_{i,1} : F_i]$).

(c) We have variants of (a) and (b) in which for the A_n type without involution and with $n > 1$ we use Subsubsection 4.1.2 instead of Proposition 4.1. In what follows we will only use the variant of (a) that is modeled on Subsubsection 4.1.2. For this variant we only have to perform the following two changes to (a):

- in the property (iii) of (a) we have to replace “either isomorphisms” by “either monomorphisms which over $\overline{\mathbb{Q}}$ are products of diagonal embeddings of the form $\mathbf{SL}_{n_i+1, \overline{\mathbb{Q}}} \hookrightarrow \mathbf{SL}_{n_i+1, \overline{\mathbb{Q}}}^{s_i}$, where $s_i \in \{1, 2\}$,”;
- in the property (v) of (a) we have to replace “some A_{n_i} type” by “some A_1 type or some A_{n_i} type with involution”.

4.10. Corollary. *Let (G_0, X_0) be an adjoint Shimura pair. Then (G_0, X_0) is the adjoint of a Shimura pair of PEL type if and only if each simple factor (G, X) of (G_0, X_0) is such that one of the following two conditions holds:*

- (i) *the Shimura pair (G, X) is of A_n type, or*
- (ii) *the Shimura pair (G, X) is of C_n , $D_n^{\mathbb{H}}$, or inner $D_4^{\mathbb{R}}$ type and the group $G_{\mathbb{R}}$ has no simple, compact factor.*

Proof: We first prove the “only if” part. Let $\tilde{f} : (\tilde{G}, \tilde{X}) \hookrightarrow (\mathbf{GSp}(\tilde{W}, \tilde{\psi}), \tilde{S})$ be an embedding of PEL type. Let (G, X) be a simple factor of $(\tilde{G}^{\text{ad}}, \tilde{X}^{\text{ad}})$. Let $\tilde{\mathcal{B}} := \text{End}(\tilde{W})^{\tilde{G}(\mathbb{Q})}$. We show that the assumption that $\tilde{\mathcal{B}}$ has two distinct factors $\tilde{\mathcal{B}}_1$ and $\tilde{\mathcal{B}}_2$ which are permuted by the involution of $\text{End}(\tilde{W})$ defined by $\tilde{\psi}$, leads to a contradiction. Let $\tilde{W} = \tilde{W}_0 \oplus \tilde{W}_1 \oplus \tilde{W}_2$ be the unique direct sum decomposition in $\tilde{\mathcal{B}}$ -modules such that $\tilde{\mathcal{B}}_1 \oplus \mathcal{B}_2$ acts trivially on $\tilde{W}_0$ and for $i \in \{1, 2\}$ the $\tilde{\mathcal{B}}_i$ -module $\tilde{W}_i$ is a direct sum of simple $\tilde{\mathcal{B}}_i$ -modules. The $\mathbb{G}_{m, \mathbb{Q}}$ subgroup of $\mathbf{GL}_{\tilde{W}}$ which fixes $\tilde{W}_0$, which acts on $\tilde{W}_1$ via the identical character of $\mathbb{G}_{m, \mathbb{Q}}$, and which acts on $\tilde{W}_2$ via the inverse of the identical character of $\mathbb{G}_{m, \mathbb{Q}}$, is contained in both $Z(\tilde{G})$ and $\mathbf{Sp}(\tilde{W}, \tilde{\psi})$. This contradicts the fact that the centralizer of each element $\tilde{x} \in \tilde{X}$ in $\mathbf{Sp}(\tilde{W}, \tilde{\psi})_{\mathbb{R}}$ is a compact group.

Thus each simple factor of $\tilde{\mathcal{B}}$ is normalized by the involution of $\text{End}(\tilde{W})$ defined by $\tilde{\psi}$. Based on this, we can assume that $\tilde{\mathcal{B}}$ is a simple $\mathbb{Q}$ -algebra; thus either $\tilde{G}^{\text{ad}} = G$ or $\tilde{G}_{\mathbb{C}}^{\text{der}}$ is a product of $\mathbf{SO}_{4, \mathbb{C}}$ groups. If $\tilde{G}_{\mathbb{C}}^{\text{der}}$ is a product of $\mathbf{SO}_{4, \mathbb{C}}$ groups, then (G, X) is of A_1 type. Thus we can assume that $\tilde{G}^{\text{ad}} = G$ and that (G, X) is not of A_n type. From this and [De2, Table 2.3.8] we get that (G, X) is of either B_n type with $n \geq 2$ or C_n type with $n \geq 2$ or $D_n^{\mathbb{H}}$ type with $n \geq 4$ or $D_n^{\mathbb{R}}$ type with $n \geq 4$. From [Sh, p. 320] we get that

$\tilde{G}_{\mathbb{R}}^{\text{der}}$ is a product of non-compact groups that are isomorphic to either $\mathbf{Sp}_{2n, \mathbb{R}}$ (with $n \geq 2$) or $\mathbf{SO}^*(2n)_{\mathbb{R}}$ (with $n \geq 4$); here the restrictions on n are as per the type of (G, X) . Thus either (ii) holds or (G, X) is of non-inner $D_4^{\mathbb{R}}$ type. But if (G, X) is of $D_4^{\mathbb{R}}$ type, then $\tilde{G}_{\overline{\mathbb{Q}}}^{\text{der}}$ is a product of $\mathbf{SO}_{8, \overline{\mathbb{Q}}}$ groups. If (G, X) is of non-inner $D_4^{\mathbb{R}}$ type, then the representation of each simple factor of $\text{Lie}(G_{\overline{\mathbb{Q}}})$ on $\tilde{W} \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}$ involves both half spin representations (see Definition 2.2.1 (c)) and thus $\tilde{G}_{\overline{\mathbb{Q}}}^{\text{der}}$ is a product of $\mathbf{Spin}_{8, \overline{\mathbb{Q}}}$ groups and therefore we reached a contradiction. Thus (ii) holds. This proves the “only if” part.

Based on Fact 2.4.1 (b), it suffices to check the “if” part only in the case when (G_0, X_0) is simple. If $(G, X) := (G_0, X_0)$ is such that (i) (resp. (ii)) holds, then from Proposition 4.1 (resp. from Subsubsection 4.8.3), we get that (G_0, X_0) is the adjoint of a Shimura pair of PEL type. $\square$

5. Applications to Frobenius tori

We now apply Section 4 to the study of Frobenius tori of abelian varieties over number fields. At the very end of Subsections 5.1 to 5.8, we prove Theorem 1.3.2 and Corollary 1.3.3. In Subsections 5.9 and 5.10 we deal with complements on the shifting process mentioned in Subsection 1.6 and elaborated in Subsection 5.4; in particular, see Proposition 5.10 for three main advantages one gains in general by performing it.

We begin by introducing a setting which is independent of Section 4. Let $E, A, L_A, H_A, \rho, G_{\mathbb{Q}_p}$, and E^{conn} be as in Subsection 1.1. Let W_A, λ_A, ψ_A , and f_A be as in Subsection 1.5. In this Section we will assume that λ_A is a principal polarization of A . Let

$$K(4) := \{g \in \mathbf{GSp}(L_A, \psi_A)(\widehat{\mathbb{Z}}) \mid g \bmod 4 \text{ is the identity}\}.$$

It is known that $\text{Sh}(\mathbf{GSp}(W_A, \psi_A), S_A)/K(4)$ is the $\mathbb{Q}$ -scheme that parameterizes isomorphism classes of principally polarized abelian schemes that are of relative dimension $\dim_{\mathbb{Q}}(W_A)/2$ and that have level-4 symplectic similitude structure, cf. [De1, Thm. 4.21]. Let $K_A(4) := H_A(\mathbb{A}_f) \cap K(4)$. We have a natural finite morphism

$$\text{Sh}(H_A, X_A)/K_A(4) \rightarrow \text{Sh}(\mathbf{GSp}(W_A, \psi_A), S_A)/K(4)$$

of $\mathbb{Q}$ -schemes (see [De1, Cor. 5.4]). Let $(\mathcal{A}, \lambda_{\mathcal{A}})$ be the pull back to $\text{Sh}(H_A, X_A)/K_A(4)$ of the universal principally polarized abelian scheme over $\text{Sh}(\mathbf{GSp}(W_A, \psi_A), S_A)/K(4)$. By replacing E with a finite field extension of it, we can assume that $E = E^{\text{conn}}$ and that there exists a morphism

$$m: \text{Spec}(E) \rightarrow \text{Sh}(H_A, X_A)/K_A(4)$$

such that $m^*(\mathcal{A}, \lambda_{\mathcal{A}})$ is (A, λ_A) . As $\text{Sh}(H_A, X_A)(\mathbb{C}) = H_A(\mathbb{Q}) \backslash (X_A \times H_A(\mathbb{A}_f))$ (see [De2, Prop. 2.1.10 and Cor. 2.1.11]), we can assume that the composite of the morphism $\text{Spec}(\mathbb{C}) \rightarrow \text{Spec}(E)$ with m is the point

$$[h_A, 1_{W_A}] \in \text{Sh}(H_A, X_A)/K_A(4)(\mathbb{C}) = H_A(\mathbb{Q}) \backslash (X_A \times H_A(\mathbb{A}_f)/K_A(4));$$

here 1_{W_A} is the identity element of $H_A(\mathbb{A}_f)$. As $E = E^{\text{conn}}$, we have $\text{Im}(\rho) \leq G_{\mathbb{Q}_p}(\mathbb{Q}_p)$.

5.1. Definition. Let B and C be two abelian varieties over a field K embeddable into $\mathbb{C}$. We fix an embedding $i_K : K \hookrightarrow \mathbb{C}$. Let H_B , H_C , and $H_{B \times_K C}$ be the Mumford–Tate groups of $B_{\mathbb{C}}$, $C_{\mathbb{C}}$, and $B_{\mathbb{C}} \times_{\mathbb{C}} C_{\mathbb{C}}$ (respectively). We say B and C are *adjoint-isogenous* if the natural projection epimorphisms $H_{B \times_K C} \twoheadrightarrow H_B$ and $H_{B \times_K C} \twoheadrightarrow H_C$ induce isomorphisms at the level of adjoint groups (thus we can identify $H_B^{\text{ad}} = H_C^{\text{ad}} = H_{B \times_K C}^{\text{ad}}$).

5.1.1. Remark. The choice of another embedding $K \hookrightarrow \mathbb{C}$ corresponds to the replacements of H_B , H_C , and $H_{B \times_K C}$ by suitable forms of them. Thus the fact that the abelian varieties B and C are or are not adjoint-isogenous does not depend on the choice of the embedding $i_K : K \hookrightarrow \mathbb{C}$.

5.2. New injective maps. Until Subsection 5.8 we will assume that the adjoint group H_A^{ad} is non-trivial. Let

$$f_2 : (G_2, X_2) \hookrightarrow (G_3, X_3) \hookrightarrow (G_4, X_4) \xrightarrow{f_4} (\mathbf{GSp}(W_1, \psi_1), S_1)$$

be injective maps of Shimura pairs constructed as in the Variants 4.9 (a), (b), or (c) but starting from the adjoint Shimura pair $(G_2^{\text{ad}}, X_2^{\text{ad}}) \stackrel{ID}{=} (H_A^{\text{ad}}, X_A^{\text{ad}})$. Thus (G_4, X_4) is unitary and the injective map $(G_4, X_4) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ of Shimura pairs is a PEL type embedding. Until Proposition 5.10 it is irrelevant if we use Variant 4.9 (a), (b), or (c) to construct these injective maps of Shimura pairs. Let

$$\mathcal{B} := \text{End}(W_1)^{G_4(\mathbb{Q})}.$$

As the group $G_2^{\text{ad}}(\mathbb{Q})$ is dense in $G_2^{\text{ad}}(\mathbb{R})$, by replacing f_2 with its composite with an automorphism $(G_2, X_2) \xrightarrow{\sim} (G_2, X_2)$ defined by an element of $G_2^{\text{ad}}(\mathbb{Q})$, we can assume that the intersection $X_2 \cap X_A$ (taken inside $X_2^{\text{ad}} = X_A^{\text{ad}}$) contains the element $h_A \in X_A$.

Let L_1 be a $\mathbb{Z}$ -lattice of W_1 such that we get a perfect alternating form $\psi_1 : L_1 \otimes_{\mathbb{Z}} L_1 \rightarrow \mathbb{Z}$. Let $K_2(4) := G_2(\mathbb{A}_f) \cap \{g \in \mathbf{GSp}(L_1, \psi_1)(\widehat{\mathbb{Z}}) \mid g \bmod 4 \text{ is the identity}\}$. As before Definition 5.1, we get naturally a principally polarized abelian scheme $(\mathcal{A}_2, \lambda_{\mathcal{A}_2})$ over $\text{Sh}(G_2, X_2)/K_2(4)$.

5.3. Hodge twists. Let $(W_0, \psi_0) := (W_A \oplus W_1, \psi_A \oplus \psi_1)$. Let G_0 be the identity component of the intersection of $\mathbf{GSp}(W_0, \psi_0)$ with the inverse image in $H_A \times_{\mathbb{Q}} G_2$ of the subgroup H_A^{ad} of $H_A^{\text{ad}} \times_{\mathbb{Q}} G_2^{\text{ad}}$ embedded in $H_A^{\text{ad}} \times_{\mathbb{Q}} G_2^{\text{ad}}$ diagonally via ID . Let X_0 be the $G_0(\mathbb{R})$ -conjugacy class of the monomorphism $\mathbb{S} \hookrightarrow G_{0, \mathbb{R}}$ defined by $(h_A, h_A) \in X_A \times X_2$ (this makes sense due to ID and the relation $h_A \in X_A \cap X_2$). We get a (diagonal) injective map

$$f_0 : (G_0, X_0) \hookrightarrow (\mathbf{GSp}(W_0, \psi_0), S_0)$$

of Shimura pairs that is the composite of the injective map $f_3 : (G_0, X_0) \hookrightarrow (H_A, X_A) \times^{\mathcal{H}} (G_2, X_2)$ of Shimura pairs with a Hodge quasi product $f_A \times^{\mathcal{H}} f_2$. The composite of f_3 with the natural injective map $(H_A, X_A) \times^{\mathcal{H}} (G_2, X_2) \hookrightarrow (H_A, X_A) \times (G_2, X_2)$ of Shimura

pairs and with the projections of $(H_A, X_A) \times (G_2, X_2)$ onto its factors, give birth at the level of adjoint Shimura pairs to identifications

$$(G_0^{\text{ad}}, X_0^{\text{ad}}) \stackrel{ID}{=} (G_2^{\text{ad}}, X_2^{\text{ad}}) \stackrel{ID}{=} (H_A^{\text{ad}}, X_A^{\text{ad}}).$$

Due to these identifications, we call (G_0, X_0) a *Hodge twist* of (H_A, X_A) and (G_2, X_2) .

Let $L_0 := L_A \oplus L_1$. It is a $\mathbb{Z}$ -lattice of W_0 and we have a perfect alternating form $\psi_0: L_0 \otimes_{\mathbb{Z}} L_0 \rightarrow \mathbb{Z}$. Let $K_0(4) := G_0(\mathbb{A}_f) \cap (K(4) \times K_2(4))$. As before Definition 5.1, we get naturally (starting from f_0 and L_0) a principally polarized abelian scheme $(\mathcal{A}_0, \lambda_{\mathcal{A}_0})$ over $\text{Sh}(G_0, X_0)/K_0(4)$.

5.4. The shifting process. Let B be an abelian variety over E which is adjoint-isogenous to A . Let H_B be the Mumford–Tate group of $B_{\mathbb{C}}$; we have a natural identification $H_B^{\text{ad}} = H_A^{\text{ad}}$. Let $\tilde{G}_{\mathbb{Q}_p}$ be the identity component of the algebraic envelope of the p -adic Galois representation attached to B ; it is a subgroup of $H_{B, \mathbb{Q}_p}$.

5.4.1. Proposition. *The images of $G_{\mathbb{Q}_p}^{\text{der}}$ and $\tilde{G}_{\mathbb{Q}_p}^{\text{der}}$ in $H_{A, \mathbb{Q}_p}^{\text{ad}} = H_{B, \mathbb{Q}_p}^{\text{ad}}$ coincide. Thus to prove the part of the Mumford–Tate conjecture that involves derived groups for the pair (A, p) is the same thing as proving it for the pair (B, p) (i.e., we have $G_{\mathbb{Q}_p}^{\text{der}} = H_{A, \mathbb{Q}_p}^{\text{der}}$ if and only if $\tilde{G}_{\mathbb{Q}_p}^{\text{der}} = H_{B, \mathbb{Q}_p}^{\text{der}}$).*

Proof: The images of $G_{\mathbb{Q}_p}^{\text{der}}$ and $\tilde{G}_{\mathbb{Q}_p}^{\text{der}}$ in $H_{A, \mathbb{Q}_p}^{\text{ad}} = H_{B, \mathbb{Q}_p}^{\text{ad}}$ coincide with the image of the identity component of the algebraic envelope of the p -adic Galois representation attached to $A \times_E B$ in $H_{A \times_E B, \mathbb{Q}_p}^{\text{ad}} = H_{A, \mathbb{Q}_p}^{\text{ad}} = H_{B, \mathbb{Q}_p}^{\text{ad}}$. Thus we have $G_{\mathbb{Q}_p}^{\text{der}} = H_{A, \mathbb{Q}_p}^{\text{der}}$ if and only if $\tilde{G}_{\mathbb{Q}_p}^{\text{der}} = H_{B, \mathbb{Q}_p}^{\text{der}}$. $\square$

5.4.2. Choice of B . As $h_A \in X_2 \cap X_A$, as $(h_A, h_A) \in X_0$, and as the functorial morphism $\text{Sh}(G_0, X_0)/K_0(4) \rightarrow \text{Sh}(H_A, X_A)/K_A(4)$ is finite, by replacing E with a finite field extension of it we can assume that there exists a morphism

$$m_0: \text{Spec}(E) \rightarrow \text{Sh}(G_0, X_0)/K_0(4)$$

such that the composite of the natural morphism $\text{Spec}(\mathbb{C}) \rightarrow \text{Spec}(E)$ with m_0 is the point $[(h_A, h_A), 1_{W_0}] \in \text{Sh}(G_0, X_0)/K_0(4)(\mathbb{C})$.

Let B be the abelian variety over E obtained from $\mathcal{A}_2$ by pull back via m_0 and the natural morphism $\text{Sh}(G_0, X_0)/K_0(4) \rightarrow \text{Sh}(G_2, X_2)/K_2(4)$. We have an identity

$$m_0^*(\mathcal{A}_0) = A \times_E B.$$

The Mumford–Tate group of $A_{\mathbb{C}} \times_{\mathbb{C}} B_{\mathbb{C}}$ (resp. of $B_{\mathbb{C}}$) is a subgroup of G_0 (resp. of G_2) that surjects onto $G_0^{\text{ad}} = G_2^{\text{ad}} = H_A^{\text{ad}}$. This implies that the abelian varieties A and B are adjoint-isogenous and that $\tilde{G}_{\mathbb{Q}_p}$ is a subgroup of $H_{B, \mathbb{Q}_p}$ and thus also of $G_{2, \mathbb{Q}_p}$.

5.4.3. Decomposing B . See Subsection 5.2 for the definition of $\mathcal{B}$. The injective map $f_4: (G_4, X_4) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ of Shimura pairs is a Hodge quasi product indexed by the set J of Subsection 1.3, cf. property (iv) of Variant 4.9 (a). Thus from the simple

$\mathbb{Q}$ -algebra statements of the property 4.1.1 (a) and of Subsubsection 4.1.2, we get that the number $s_{\mathcal{B}}$ of simple factors of $\mathcal{B}$ equals to the number of elements of J . If we take L_1 to be well adapted for f_4 (see Subsection 2.4), then B is a product indexed by J of abelian varieties over E that have the following two properties: (i) they are principally polarized, and (ii) their extensions to $\mathbb{C}$ have Mumford–Tate groups whose adjoints are simple.

5.5. Good choices for v . Let v, l, A_v , and T_v be as before Theorem 1.1.5. We choose the prime v of E such that moreover we have $l > \dim(B)$, $l \neq p$, and the following two conditions hold:

- (i) the abelian variety B has good reduction B_v with respect to v ;
- (ii) the prime v is unramified over l and $\mathcal{B}_{(l)} := \mathcal{B} \cap \text{End}(L_1 \otimes_{\mathbb{Z}} \mathbb{Z}_{(l)})$ is a semisimple $\mathbb{Z}_{(l)}$ -algebra that is self dual with respect to the perfect alternating ψ_1 on L_1 .

Let $T_{v,2}$ and $T_{v,0}$ be the Frobenius tori of B_v and $A_v \times_{k(v)} B_v$ (respectively). Let $\pi_v \in T_v(\mathbb{Q}_p)$, $\pi_{v,0} \in T_{v,0}(\mathbb{Q}_p)$, and $\pi_{v,2} \in T_{v,2}(\mathbb{Q}_p)$ be (the same integral power of) Frobenius elements obtained as in Subsection 3.2 via the choice of a Frobenius automorphism of a prime $\bar{v}$ of $\bar{E} = \bar{\mathbb{Q}}$ that divides v . We have natural monomorphisms $T_{v,\mathbb{Q}_p} \hookrightarrow G_{\mathbb{Q}_p}$, $T_{v,2,\mathbb{Q}_p} \hookrightarrow \tilde{G}_{\mathbb{Q}_p} \hookrightarrow G_{2,\mathbb{Q}_p}$, and $T_{v,0,\mathbb{Q}_p} \hookrightarrow G_{0,\mathbb{Q}_p}$. The images of π_v , $\pi_{v,0}$, and $\pi_{v,2}$ in $H_{A,\mathbb{Q}_p}^{\text{ad}}(\mathbb{Q}_p) = G_{0,\mathbb{Q}_p}^{\text{ad}}(\mathbb{Q}_p) = G_{2,\mathbb{Q}_p}^{\text{ad}}(\mathbb{Q}_p)$ coincide.

By replacing E with a finite field extension of it, we can assume that the image of the p -adic Galois representation $\rho_2 : \text{Gal}(E) \rightarrow \mathbf{GL}_{L_1 \otimes_{\mathbb{Z}} \mathbb{Q}_p}(\mathbb{Q}_p)$ attached to B factors through $G_4(\mathbb{Q}_p)$ (i.e., all $\mathbb{Q}$ -endomorphisms of $B_{\mathbb{C}}$ defined by elements of $\mathcal{B}$, are defined over E). Each $\mathbb{Q}$ -endomorphism of B defined by an element of $\mathcal{B}$ defines naturally a $\mathbb{Q}$ -endomorphism of B_v . Let λ_B be the principal polarization of B that is the natural pull back of λ_{A_2} . Let λ_{B_v} be the principal polarization of B_v that is defined naturally by λ_B .

5.5.1. Lemma. *There exist a finite field extension $k(v')$ of $k(v)$, a complete discrete valuation ring R of mixed characteristic $(0, l)$ and of residue field $k(v')$, and a principally polarized abelian scheme $(B', \lambda_{B'})$ over R , such that the following three properties hold:*

(i) *the special fibre $(B'_{v'}, \lambda_{B'_{v'}})$ of $(B', \lambda_{B'})$ is $\mathbb{Z}[\frac{1}{l}]$ -isomorphic to $(B_{v'}, \lambda_{B_{v'}}) := (B_v, \lambda_{B_v})_{k(v')}$ (i.e., it is obtained from $(B_{v'}, \lambda_{B_{v'}})$ via isogenies of order a power of l);*

(ii) *the abelian scheme B' has complex multiplication;*

(iii) *the $\mathbb{Q}$ -endomorphisms of $B'_{v'}$ that correspond to elements of $\mathcal{B}$, lift to $\mathbb{Q}$ -endomorphisms of B' in such a way that the elements of $\mathcal{B}_{(l)}$ correspond to $\mathbb{Z}_{(l)}$ -endomorphisms of B' and the trace forms on $\mathcal{B}_{(l)} \otimes_{\mathbb{Z}_{(l)}} k(v')$ defined by its representations on the tangent spaces of $B_{v'}$ and $B'_{v'}$ at identity elements, are equal.*

Proof: By using a product decomposition of B (and thus implicitly of $(L_1, \psi_1, \mathcal{B}_{(l)})$) as in Subsubsection 5.4.3, to prove the Lemma we can assume that $\mathcal{B}$ is a simple $\mathbb{Q}$ -algebra. Let J_1 be the involution of $\text{End}(W_1)$ defined by the identity $\psi_1(e(x), y) = \psi_1(x, J_1(e)(y))$, where $e \in \text{End}(W_1)$ and $x, y \in W_1$. As G_4 is a subgroup of $\mathbf{GSp}(W_1, \psi_1)$, the involution J_1 normalizes $\mathcal{B}$ and thus also $\mathcal{B}_{(l)}$. As the injective map $f_4 : (G_4, X_4) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ of Shimura pairs is a PEL type embedding, the group G_4 is the centralizer of $\mathcal{B}$ in $\mathbf{GSp}(W_1, \psi_1)$. From this and the fact that the Shimura pair (G_4, X_4) is unitary, we get

that the restriction of the involution $\mathcal{I}_1$ to $\mathcal{B}$ is of type A in the sense of [Zi, Subsect. 2.3]. Thus the Lemma is a particular case of [Zi, Thm. 4.4] applied to the data provided by the injective map $f_4 : (G_4, X_4) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ of Shimura pairs, the perfect alternating form $\psi_1 : L_1 \otimes_{\mathbb{Z}} L_1 \rightarrow \mathbb{Z}$, the involution $\mathcal{I}_1$, the prime l , the semisimple $\mathbb{Z}_{(l)}$ -algebra $\mathcal{B}_{(l)}$, and the principally polarized abelian variety (B_v, λ_{B_v}) over $k(v)$ which is the good reduction of the pull back (B, λ_B) of $(\mathcal{A}_2, \lambda_{\mathcal{A}_2})$. Strictly speaking loc. cit. states only that $(B'_{v'}, \lambda_{B'_{v'}})$ is obtained from (B_v, λ_{B_v}) via isogenies. But one can always assume that these isogenies are of order a power of l . $\square$

5.6. Applications of 5.5.1. We can assume that the field $R[\frac{1}{l}]$ contains E . We fix an embedding $R[\frac{1}{p}] \hookrightarrow \mathbb{C}$ that extends the embedding i_E of Subsection 1.1. Let $W'_1 := H_1(B'(\mathbb{C}), \mathbb{Q})$, let ψ'_1 be the alternating form on W'_1 induced by $\lambda_{B'}$, and let $\mathcal{B} \hookrightarrow \text{End}(W'_1)$ be the $\mathbb{Q}$ -monomorphism defined by the identification (see property 5.5.1 (iii)) of $\mathcal{B}$ with a $\mathbb{Q}$ -subalgebra of $\text{End}(B'_\mathbb{C}) \otimes_{\mathbb{Z}} \mathbb{Q}$. Let G'_4 be the subgroup of $\mathbf{GSp}(W'_1, \psi'_1)$ that fixes all elements of $\mathcal{B}$. From the property 5.5.1 (i) we get that for each prime $\tilde{l} \in \mathbb{N}$ different from l , the two triples $(W'_1, \psi'_1, \mathcal{B})$ and $(W_1, \psi_1, \mathcal{B})$ become isomorphic over $\mathbb{Q}_{\tilde{l}}$. Thus the “difference” between the isomorphism classes of the two triples $(W'_1, \psi'_1, \mathcal{B})$ and $(W_1, \psi_1, \mathcal{B})$ is “measured” by a class $\delta_4 \in H^1(\mathbb{Q}, G_4)$ whose image in $H^1(\mathbb{Q}_{\tilde{l}}, G_{4, \mathbb{Q}_{\tilde{l}}})$ is trivial for all primes $\tilde{l}$ different from l . In particular, we get that G'_4 is an inner form of G_4 .

5.6.1. Fact. *There exists a reductive subgroup $G_4^{(1)}$ of $\mathbf{GL}_{W_1}$ that is the extension of $\mathbb{G}_{m, \mathbb{Q}}^{s_{\mathcal{B}}-1}$ by G_4 and such that the images of the class δ_4 in $H^1(\mathbb{R}, G_{4, \mathbb{R}}^{(1)})$ and $H^1(\mathbb{Q}_q, G_{4, \mathbb{Q}_q}^{(1)})$ are trivial, for all primes $q \in \mathbb{N}$ (see Subsubsection 5.4.3 for the definition of $s_{\mathcal{B}}$).*

Proof: By using a product decomposition of B (and thus implicitly of $(L_1, \psi_1, \mathcal{B}_{(l)})$) as in Subsubsection 5.4.3 and due to the first part of Fact 2.4.1 (a), to prove the Lemma we can assume that $\mathcal{B}$ is a simple $\mathbb{Q}$ -algebra. Thus $s_{\mathcal{B}} = 1$ and $G_4^{(1)} = G_4$. We already know that the image of δ_4 in $H^1(\mathbb{Q}_q, G_{4, \mathbb{Q}_q})$ is trivial provided $q \neq l$. The fact that the images of δ_4 in both pointed sets $H^1(\mathbb{R}, G_{4, \mathbb{R}}^{(1)})$ and $H^1(\mathbb{Q}_l, G_{4, \mathbb{Q}_l}^{(1)})$ are trivial is checked in the third and fourth (respectively) paragraphs of [Ko2, Sect. 8]. To loc. cit. we only have to add that the inequality $l > \dim(B)$ implies that it is irrelevant if in the context of the data provided in the proof of Lemma 5.5.1, we work with the determinant condition of [Ko2, Sect. 5] or with the trace condition of the property 5.5.1 (iii) as in [Zi]. $\square$

From Fact 5.6.1 we get that we have an identification $G_{4, \mathbb{Q}_p} = G'_{4, \mathbb{Q}_p}$ (unique up to inner conjugation) as well as a canonical identification $Z(G_4) = Z(G'_4)$.

Let (G'_4, X'_4) be the Shimura pair, where X'_4 is the $G'_4(\mathbb{R})$ -conjugacy class of the homomorphism $\mathbb{S} \rightarrow G'_{4, \mathbb{R}}$ that defines the Hodge $\mathbb{Q}$ -structure on W'_1 . If we have $s_{\mathcal{B}} = 1$, then there exists an isomorphism $W_1 \otimes_{\mathbb{Q}} \mathbb{R} \xrightarrow{\sim} W'_1 \otimes_{\mathbb{Q}} \mathbb{R}$ which (cf. [Ko2, Lem. 4.2 and the third paragraph of Sect. 8]) has the following two properties:

- (a) it takes ψ_1 to ψ'_1 and it takes b to b for all $b \in \mathcal{B}$ (thus it takes the subgroup $G_{4, \mathbb{R}}$ of $\mathbf{GL}_{W_1 \otimes_{\mathbb{Q}} \mathbb{R}}$ onto the subgroup $G'_{4, \mathbb{R}}$ of $\mathbf{GL}_{W'_1 \otimes_{\mathbb{Q}} \mathbb{R}}$);
- (b) it takes X_4 onto X'_4 .

Thus for $s_B = 1$, the canonical identification $Z(G_4) = Z(G'_4)$ gives birth naturally (cf. (b)) to a canonical identification $(G_4^{\text{ab}}, X_4^{\text{ab}}) = (G_4'^{\text{ab}}, X_4'^{\text{ab}})$ of 0 dimensional Shimura pairs. Based on this and the product decomposition of Subsubsection 5.4.3, regardless of what the number $s_B \in \mathbb{N}$ is, we get:

5.6.2. Fact. *Let $Z_{\min}$ be the smallest subtorus of $Z^0(G_4)$ with the property that all monomorphisms $\mathbb{S} \hookrightarrow G_{4,\mathbb{R}}$ defined by elements of X_4 , factor through the extension to $\mathbb{R}$ of the subgroup of G_4 generated by G_4^{der} and $Z_{\min}$. Let $Z'_{\min}$ be the subtorus of $Z^0(G'_4)$ that is defined similarly to $Z_{\min}$. Then the canonical identification $Z(G_4) = Z(G'_4)$ restricts to a canonical identification $Z_{\min} = Z'_{\min}$.*

5.6.3. Betti realization of $\pi_{v,2}$. We use the above identification $G_{4,\mathbb{Q}_p} = G'_{4,\mathbb{Q}_p}$. Coming back to the Frobenius elements of Subsection 5.5, from properties 5.5.1 (ii) and (iii) we get (cf. the third paragraph of Subsection 3.2 and the relation $\text{Im}(\rho_2) \leq G_4(\mathbb{Q}_p)$) the existence of a torus T'_4 of G'_4 and of an element $\pi'_{v',2} \in T'_4(\mathbb{Q})$ such that the following three properties hold:

(i) the element $\pi'_{v',2}$ is the Betti realization of an endomorphism of B' which lifts a suitable integral power of the Frobenius endomorphism of $B'_{k(v')}$;

(ii) the torus T'_4 is the smallest subgroup of G'_4 which has $\pi'_{v',2}$ as a $\mathbb{Q}$ -valued point;

(iii) there exists an element $g' \in G'_4(\mathbb{Q}_p) = G_4(\mathbb{Q}_p)$ that takes under conjugation $T'_{4,\mathbb{Q}_p}$ to $T_{v,2,\mathbb{Q}_p}$ and $\pi'_{v',2}$ to $\pi_{v,2}$. [Here we have to introduce the element g' as the torus T'_4 of G'_4 is uniquely determined only up to $G'_4(\mathbb{Q})$ -conjugation and as $T_{v,2,\mathbb{Q}_p}$ and $\pi_{v,2}$ are uniquely determined (cf. Subsection 3.2) only up to $\text{Im}(\rho_2)$ -conjugation].

The images of $T_{v,\overline{\mathbb{Q}_p}}$ and $T'_{4,\overline{\mathbb{Q}_p}}$ in $G_4^{\text{ad}}/\overline{\mathbb{Q}_p} = G_4'^{\text{ad}}/\overline{\mathbb{Q}_p}$ are $G_4^{\text{ad}}(\overline{\mathbb{Q}_p})$ -conjugate, cf. (iii) and the end of Subsection 5.5.

5.6.4. A crystalline property. As R is a complete discrete valuation ring of mixed characteristic $(0, l)$ and of residue field $k(v')$, it has a canonical structure of a $W(k(v'))$ -algebra. Let (M, ϕ) be the F -crystal over $k(v)$ of B_v . Thus the pair $(M \otimes_{W(k(v))} B(k(v')), \phi)$ is the F -isocrystal over $k(v')$ of $B'_{v'}$. The canonical and functorial identification $M \otimes_{W(k(v))} R[\frac{1}{p}] = H_{\text{dR}}^1(B'/R)[\frac{1}{p}]$ (see [BO, Thm. 1.3]) allows us to identify the de Rham realization of the endomorphism of B' mentioned in the property 5.6.3 (i) with a suitable integral power of $\phi^{[k(v'):\mathbb{F}_p]}$. From this and the property 3.4.1 (c) applied in the context of the pair (B, v) (instead of (A, v)), we get that a suitable $G'_4(\overline{\mathbb{Q}_l})$ -conjugate of $T'_{4,\overline{\mathbb{Q}_l}}$ is a torus of $\tilde{G}_{\overline{\mathbb{Q}_l}}$.

5.7. Simple properties. (a) Each simple factor of $\text{Lie}(G_{4,\mathbb{R}}^{\text{der}})$ is absolutely simple and only one simple factor of $\text{Lie}(G_{2,\mathbb{R}}^{\text{der}})$ maps injectively into it. Each simple factor of $\text{Lie}(G_{2,\mathbb{R}}^{\text{der}})$ is embedded (“diagonally”) into a product of simple factors of $\text{Lie}(G_{3,\mathbb{R}}^{\text{der}})$ isomorphic to it. The same holds over $\overline{\mathbb{Q}_p}$ instead of over $\mathbb{R}$. Moreover, each simple factor of $\text{Lie}(G_2^{\text{ad}})$ projects injectively into a unique simple factor of $\text{Lie}(G_4^{\text{ad}})$. All these follow from the property (iii) of Variant 4.9 (a) and from Variants 4.9 (b) and (c).

(b) We use the identification $G_{4, \overline{\mathbb{Q}_p}} = G'_{4, \overline{\mathbb{Q}_p}}$ (see paragraph after Fact 5.6.1). Two simple factors of $G_{4, \overline{\mathbb{Q}_p}}^{\text{ad}}$ are connected over $\mathbb{Q}$ through G_4^{ad} (i.e., are factors of the extension to $\overline{\mathbb{Q}_p}$ of the same simple factor of G_4^{ad}) if and only if they are connected over $\mathbb{Q}$ through $G_4'^{\text{ad}}$. This is so as G_4' is an inner form of G_4 .

(c) If $s_B = 1$ and $(G_4^{\text{ad}}, X_4^{\text{ad}})$ is a simple Shimura pair of some A_n type with n odd, then the Hasse principle holds for G_4 (cf. [Ko2, Sect. 7]). Thus the class $\delta_4 \in H^1(\mathbb{Q}, G_4)$ is trivial, cf. Fact 5.6.1.

(d) If all simple factors of $(G_4^{\text{ad}}, X_4^{\text{ad}})$ are of some A_n type with n odd and if in the property 5.5.1 (i) we require only that $B_{v'}$ is isogenous (instead of being $\mathbb{Z}[\frac{1}{l}]$ -isomorphic) to B_v and that properties 5.5.1 (ii) and (iii) hold, then one can easily check based on (c) that we can choose $k(v')$, $B_{v'}$, R , and B' in such a way that δ_4 is the trivial class (regardless of what $s_B \in \mathbb{N}$ is). The last sentence applies if there exists no element $j \in J$ such that (H_j, X_j) is of A_n type with n even, cf. the definition of the group $G(A)$ in Subsection 4.3 and the property 4.8 (iii).

5.8. Proofs of 1.3.2 and 1.3.3. We are ready to prove Theorem 1.3.2 and Corollary 1.3.3. We can assume that the adjoint group H_A^{ad} is non-trivial. Let $N(A) \in \mathbb{N}$ be a number that depends only on A and that has the property that for each prime $l \geq N(A)$, both conditions 5.5 (i) and (ii) hold and moreover A has good reduction with respect to all primes of E that divide l . By replacing E with a finite field extension of it, to prove Theorem 1.3.2 and Corollary 1.3.3 we can assume that A has a principal polarization λ_A (cf. [Mu2, Ch. IV, Sect. 23, Cor. 1]) and that (as $l \geq N(A)$) we are in the context of Subsections 5.2 to 5.7. As T_4' is a torus of G_4' , the images of T_4' in simple factors of the extension to $\overline{\mathbb{Q}_p}$ of a fixed simple factor of $G_4'^{\text{ad}}$ (equivalently of G_4^{ad} , cf. property 5.7 (b)), have equal ranks that do not depend on $p \neq l$. Thus also the images of $T_{v, \overline{\mathbb{Q}_p}}$ in simple factors of the extension to $\overline{\mathbb{Q}_p}$ of a fixed simple factor of $H_A^{\text{ad}} = G_2^{\text{ad}}$, have equal ranks (cf. end of Subsubsection 5.6.3 and the property 5.7 (a)) that do not depend on $p \neq l$. This proves Theorem 1.3.2.

In order to get Corollary 1.3.3 from Theorem 1.3.2, we choose the prime v of E such that $l \geq N(A)$ and T_v has the same rank as $G_{\mathbb{Q}_p}$. Thus the fact that Corollary 1.3.3 holds for $p \neq l$ follows from Theorem 1.3.2. As in the end of Lemma 3.3, using a prime v_1 of E that divides a prime $l_1 \in \mathbb{N}$ that is greater than $N(A)$ and different from l , we get that Corollary 1.3.3 holds even if $p = l$. This proves Corollary 1.3.3. $\square$

5.9. Definition. An irreducible representation of a split, simple Lie algebra of classical Lie type $\mathfrak{L}$ over a field K of characteristic 0 is called an *SD-standard representation*, if the following two conditions hold: (i) its highest weight is a minuscule weight, and (ii) if $\mathfrak{L} = A_n$, then moreover its dimension is $n + 1$ (i.e., its highest weight is either ϖ_1 or ϖ_n). Here SD stands for Satake–Deligne, cf. [Sa2] and [De2, Prop. 2.3.10], [Va1, Subsects. 6.5 and 6.6], and the proofs of Proposition 4.1 and Theorem 4.8. Similarly, we speak about SD-standard representations of split, semisimple groups of classical Lie type over K whose adjoints are simple.

Until the end, we continue to assume that the adjoint group H_A^{ad} is non-trivial. Next we list some extra advantages we get (besides proving Theorem 1.3.2) by replacing the abelian variety A and the injective map $f_A : (H_A, X_A) \hookrightarrow (\mathbf{GSp}(W_A, \psi_A), S_A)$ of Shimura pairs with B and $f_2 : (G_2, X_2) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ (respectively).

5.10. Proposition. (a) *The irreducible subrepresentations of the representation of $\text{Lie}(G_{2,\overline{\mathbb{Q}}}^{\text{der}})$ on $W_1 \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}$ factor through simple factors of $\text{Lie}(G_{2,\overline{\mathbb{Q}}}^{\text{der}})$ and are SD-standard representations of these simple factors.*

(b) *By using Variant 4.9 (a) or (c) (resp. 4.9 (c)) to construct the injective maps of Shimura pairs of Subsection 5.2, we can assume that for each simple factor (H_j, X_j) of $(H_A^{\text{ad}}, X_A^{\text{ad}}) = (G_2^{\text{ad}}, X_2^{\text{ad}})$ of $D_n^{\mathbb{R}}$ type with $n \geq 4$ (resp. of A_n type without involution and with $n \geq 2$) and every simple factor $\mathcal{F}_j$ of $H_{j,\overline{\mathbb{Q}}}$, there exists a $G_{2,\overline{\mathbb{Q}}}$ -submodule $\mathcal{W}$ of $W_1 \otimes_{\mathbb{Q}} \overline{\mathbb{Q}}$ that has the following two properties:*

(i) *the natural representation $Z^0(G_{2,\overline{\mathbb{Q}}}) \rightarrow \mathbf{GL}_{\mathcal{W}}$ factors through $Z(\mathbf{GL}_{\mathcal{W}})$ (i.e., $Z^0(G_{2,\overline{\mathbb{Q}}})$ acts on $\mathcal{W}$ via scalar automorphisms), and*

(ii) *both half spin representations (resp. both irreducible representations of the highest weights ϖ_1 and ϖ_n) of $\text{Lie}(\mathcal{F}_j)$ are among the simple $\text{Lie}(G_{2,\overline{\mathbb{Q}}}^{\text{ad}})$ -submodules of $\mathcal{W}$.*

(c) *If $H_{A,\mathbb{R}}^{\text{ad}}$ has no simple, compact factor and if $(H_A^{\text{ad}}, X_A^{\text{ad}})$ has no simple factor of some A_n type with involution, then we can also choose $f_2 : (G_2, X_2) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ such that we have $Z^0(G_2) = Z(\mathbf{GL}_{W_1}) \simeq \mathbb{G}_{m,\mathbb{Q}}$.*

Proof: Part (a) is a consequence of the proof of Proposition 4.1 and of Subsection 4.3, cf. property (iii) of Variant 4.9 (a) and Variants 4.9 (b) and (c). Part (b) for the $D_n^{\mathbb{R}}$ type with $n \geq 4$ (resp. the A_n type without involution and with $n \geq 2$) is a consequence of the fact that in Subsection 4.3 we defined $h_{\mathbb{R}}$ using both half spin representations (resp. of the first paragraph of Subsubsection 4.1.2). Based on Fact 2.4.1 (a), to check (c) we can assume that the adjoint Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is simple. Thus, if $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is not (resp. is) of A_n Lie type, then (c) is a consequence of Remark 4.1.3 (resp. of the first paragraph of Subsubsection 4.8.3). $\square$

5.10.1. Remark. By replacing (A, f_A) with (B, f_2) , the role of the Mumford–Tate group H_A gets replaced by the one of a Mumford–Tate group which is a reductive, normal subgroup of G_2 that contains G_2^{der} (more precisely by the one of the smallest subgroup of G_2 with the property that all homomorphisms $\mathbb{S} \rightarrow G_{2,\mathbb{R}}$ defined by elements of X_2 factor through its extension to $\mathbb{R}$). Thus under such a replacement we lose to a great extent the “control” on the torus $Z(H_A)$; moreover, we often have $\dim(B) > \dim(A)$. If this looks “unpleasant” or if one would like to “stick” to the injective map f_A of Shimura pairs, then one can consider the PEL-envelope $(\tilde{H}_A, \tilde{X}_A)$ of f_A as defined in [Va1, Rm. 4.3.12]. In other words, one can consider natural injective maps

$$f_A : (H_A, X_A) \hookrightarrow (\tilde{H}_A, \tilde{X}_A) \xrightarrow{\tilde{f}_A} (\mathbf{GSp}(W_A, \psi_A), S_A)$$

of Shimura pairs, where $\tilde{H}_A$ is the identity component of the intersection of $\mathbf{GSp}(W_A, \psi_A)$ with the double centralizer of H_A in $\mathbf{GL}_{W_A}$. Thus $\tilde{f}_A$ is a PEL type embedding and we

get a variant of Subsubsection 5.6.3 for its context: if the analogues of the conditions 5.5 (i) and (ii) hold in the context of A and $\tilde{f}_A$ (instead of B and f_2), then Subsections 5.5.1 to 5.6.3 can be adapted to give us that for $l \gg 0$ we can view π_v as a $\mathbb{Q}$ -valued point of a torus $\tilde{T}'_A$ of a suitable form $\tilde{H}'_A$ of $\tilde{H}_A$. This form is inner provided $\tilde{H}_{A,\mathbb{R}}^{\text{der}}$ has no normal subgroup that is an $\mathbf{SO}^*(2n)_{\mathbb{R}}$ group with $n \geq 2$, cf. the connectedness aspects of [Ko2, Sect. 7]. We do not know how one could get Theorem 1.3.2 via $\tilde{T}'_A$ (i.e., via using $\tilde{f}_A$ instead of f_2 of Subsection 5.2).

Let $\tilde{C}_A$ be the centralizer of $Z^0(H_A)$ in $\mathbf{GSp}(W_A, \psi_A)$. Let $\tilde{H}_{A,1}^{\text{ad}}$ be the maximal factor of $\tilde{C}_A^{\text{ad}}$ with the property that no simple factor of it becomes compact over $\mathbb{R}$. Let $\tilde{H}_{A,1}$ be the maximal normal, reductive subgroup of $\tilde{C}_A$ whose adjoint is $\tilde{H}_{A,1}^{\text{ad}}$. We have a variant of the previous paragraph in which we work with $\tilde{H}_{A,1}$ instead of $\tilde{H}_A$. The monomorphism $\tilde{H}_{A,1} \hookrightarrow \mathbf{GSp}(W_A, \psi_A)$ extends uniquely to an injective map $\tilde{f}_{A,1} : (\tilde{H}_{A,1}, \tilde{X}_{A,1}) \hookrightarrow (\mathbf{GSp}(W_A, \psi_A), S_A)$ of Shimura pairs through which $\tilde{f}_A$ factors. The injective map $\tilde{f}_{A,1}$ of Shimura pairs is a PEL type embedding and thus for $l \gg 0$ we can view π_v as a $\mathbb{Q}$ -valued point of a torus $\tilde{T}'_{A,1}$ of a suitable form of $\tilde{H}_{A,1}$. As the group $\tilde{C}_{A,\mathbb{Q}}^{\text{der}}$ (and thus also $\tilde{H}_{A,1,\mathbb{Q}}^{\text{ad}}$) has no normal subgroup which is an $\mathbf{SO}$ group, this form is inner. Also, one can use the torus $\tilde{T}'_{A,1}$ and [Zi] in a way similar to Subsections 5.5 and 5.6, to re-obtain Remark 3.3.1 for $l \gg 0$.

5.10.2. Remark. Different variants of Proposition 5.10 can be obtained by working with suitable other abelian varieties over E that are adjoint-isogenous to A and that are obtained similarly to B but via another injective map $\tilde{f}_2 : (\tilde{G}_2, \tilde{X}_2) \hookrightarrow (\mathbf{GSp}(\tilde{W}_1, \tilde{\psi}_1), \tilde{S}_1)$ of Shimura pairs for which we have an identity $(\tilde{G}_2^{\text{ad}}, \tilde{X}_2^{\text{ad}}) = (H_A^{\text{ad}}, X_A^{\text{ad}})$. Thus for getting Proposition 5.10, we could have only quoted [De2, Subsubsects. 2.3.10 to 2.3.13]. Also, by quoting loc. cit. and by considering PEL-envelopes as in Remark 5.10.1 one gets directly: (i) less explicit forms of Theorem 4.8 provided the simple, adjoint Shimura pair (G, X) of Section 4 is of B_n , C_n , or $D_n^{\mathbb{H}}$ type, and (ii) less explicit and weaker variants of Theorem 4.8 and Remark 4.8.1 if (G, X) is of $D_n^{\mathbb{R}}$ type. For instance, if (G, X) is of $D_n^{\mathbb{R}}$ type with $n \in 2 + 2\mathbb{N}$, then the half spin representations of the D_n Lie type are self dual (cf. [Bou2, p. 210]) and thus [De2, Subsubsects. 2.3.10 to 2.3.13] does not apply to get directly that in Subsections 4.7 and 4.8 we can choose $(G_1, X_1) = (G_4^{\text{ad}}, X_4^{\text{ad}})$ to be a simple, adjoint Shimura pair.

In addition, the explicit character of Subsections 4.1 to 4.8 and the new features of Proposition 4.1 and Remarks 4.8.2 (b) and (c) offer simplifications in many practical situations (like in Definition 6.1 (b) and in the proofs of Theorems 7.1 and 7.2 below).

6. Non-special A_n types

In this Section we define and study simple, adjoint Shimura pairs of non-special A_n type. See Subsection 6.2 for different numerical aspects and examples that pertain to them. The notations will be independent from the ones of Subsections 1.1 and 1.3. If (G_0, X_0) is a Shimura pair and $x \in X_0$, let the cocharacter $\mu_x : \mathbb{G}_{m,\mathbb{C}} \rightarrow G_{0,\mathbb{C}}$ be defined as in Subsection 2.2. We use $+$ to denote the Lie type of products of split, absolutely simple, adjoint groups.

6.1. Definitions. (a) Let (H_1, X_1) be a Shimura pair. A torus T_1 of H_1 is called *Sh-good* for (H_1, X_1) , if there exists a set $\mathfrak{C}_1$ of cocharacters $\mathbb{G}_{m,\mathbb{C}} \rightarrow H_{1,\mathbb{C}}$ that are $H_1(\mathbb{C})$ -conjugate to the cocharacters $\mu_x : \mathbb{G}_{m,\mathbb{C}} \rightarrow H_{1,\mathbb{C}}$ ($x \in X_1$) and such that T_1 is the smallest subgroup of H_1 with the property that the elements of $\mathfrak{C}_1$ factor through $T_{1,\mathbb{C}}$.

(b) Let (H_0, X_0) be a simple, adjoint Shimura pair of either A_1 type or of A_n type with $n \geq 2$ and with involution (resp. of A_n type with $n \geq 2$ and without involution). Let $f_1 : (H_1, X_1) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ be an injective map of Shimura pairs such that the following four conditions hold:

- we have a natural identity $(H_1^{\text{ad}}, X_1^{\text{ad}}) = (H_0, X_0)$;
- property 5.10 (a) holds (resp. properties 5.10 (a) and (b) hold) for $f_2 = f_1$;
- the intersection $\tilde{H}_1$ of $\mathbf{GSp}(W_1, \psi_1)$ with the double centralizer of H_1 in $\mathbf{GL}_W$ is connected and the monomorphism $H_{1,\mathbb{Q}}^{\text{der}} \hookrightarrow \tilde{H}_{1,\mathbb{Q}}^{\text{der}}$ is an isomorphism (resp. is isomorphic to the diagonal embedding of $H_{1,\mathbb{Q}}^{\text{der}}$ in $H_{1,\mathbb{Q}}^{\text{der}} \times_{\overline{\mathbb{Q}}} H_{1,\mathbb{Q}}^{\text{der}}$);
- H_1 is a Mumford–Tate group (i.e., H_1 is the smallest subgroup of $\mathbf{GSp}(W_1, \psi_1)$ such that all monomorphisms $\mathbb{S} \hookrightarrow H_{1,\mathbb{R}}$ that are elements of X_1 , factor through $H_{1,\mathbb{R}}$).

Let $q \in \mathbb{N}$ be a prime. The image of each element $x \in X_1$ in $\tilde{H}_{1,\mathbb{R}}^{\text{ad}}$ is non-trivial. Thus the group $\tilde{H}_1^{\text{ad}}$ has no simple factor which over $\mathbb{R}$ is compact. From this we get that the $\tilde{H}_1(\mathbb{R})$ -conjugacy class $\tilde{X}_1$ of monomorphisms $\mathbb{S} \hookrightarrow \tilde{H}_{1,\mathbb{R}}$ defined by elements of X_1 , has the property that we get injective maps $(H_1, X_1) \hookrightarrow (\tilde{H}_1, \tilde{X}_1) \xrightarrow{\tilde{f}_1} (\mathbf{GSp}(W_1, \psi_1), S_1)$ of Shimura pairs. Let $\tilde{f}'_1 : (\tilde{H}'_1, \tilde{X}'_1) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ be an injective map of Shimura pairs defined by the twist of $\tilde{f}_1$ via a class in $H^1(\mathbb{Q}, \tilde{H}_1)$ that is locally trivial. We identify $\tilde{X}'_1 = \tilde{X}_1$ and $\tilde{H}_{1,\mathbb{Q}_q} = \tilde{H}'_{1,\mathbb{Q}_q}$.

By an *MT pair* for the triple $(f_1, \tilde{f}'_1, q)$ we mean a pair (T_1, G_1) , where T_1 is a torus of $\tilde{H}'_1$ that is *Sh-good* for $(\tilde{H}'_1, \tilde{X}'_1)$ and where G_1 is a reductive subgroup of $H_{1,\mathbb{Q}_q}$ that has the following two properties:

- (i) the group G_1 has a maximal torus that is $\tilde{H}'_1(\mathbb{Q}_q)$ -conjugate to $T_{1,\mathbb{Q}_q}$;
- (ii) we have an identity $\text{End}(W_1 \otimes_{\mathbb{Q}} \mathbb{Q}_q)^{G_1(\mathbb{Q}_q)} = \text{End}(W_1 \otimes_{\mathbb{Q}} \mathbb{Q}_q)^{H_{1,\mathbb{Q}_q}(\mathbb{Q}_q)}$.

(c) Let (H_0, X_0) be as in (b). We say that (H_0, X_0) is of *non-special A_n type*, if there exists an injective map $f_1 : (H_1, X_1) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ of Shimura pairs as in (b) and a prime $q \in \mathbb{N}$ such that for each twist $\tilde{f}'_1$ of $\tilde{f}_1$ obtained as in (b) and for every *MT pair* (T_1, G_1) for $(f_1, \tilde{f}'_1, q)$, the group $G_{1,\mathbb{Q}_q}^{\text{ad}}$ has at least one simple factor of A_n Lie type.

6.2. Numerical tests. Let (H_0, X_0) be a simple, adjoint Shimura pair of A_n type. In this Subsection we list some numerical properties which are implied by the assumption that (H_0, X_0) is or is not of non-special A_n type and which rely on the classification (see Subsection 2.1) of minuscule weights.

We write

$$H_{0,\mathbb{R}} = \prod_{j \in J_1} \text{SU}(a_j, b_j)_{\mathbb{R}}^{\text{ad}} \times \prod_{j \in J_0} \text{SU}(0, n+1)_{\mathbb{R}}^{\text{ad}},$$

where for $j \in J_1$ we have $a_j, b_j \in \mathbb{N}$, $a_j \leq b_j$, and $a_j + b_j = n + 1$. Here J_0 and J_1 are finite sets that keep track of the simple, non-compact and the simple, compact (respectively) factors of $H_{0,\mathbb{R}}$. As $H_{0,\mathbb{R}}$ has simple, non-compact factors (see Subsection 2.2), the set J_1 is non-empty. For $j \in J_1$ let

$$c_j := \frac{a_j}{b_j}.$$

Let

$$\mathfrak{C} := \{c_j | j \in J_1\} \subseteq \left\{ \frac{1}{n}, \frac{2}{n-1}, \dots, \frac{\lfloor \frac{n+1}{2} \rfloor}{n+1 - \lfloor \frac{n+1}{2} \rfloor} \right\} \text{ and } \mathfrak{M} := \{(a_j, b_j), (b_j, a_j) | j \in J_1\}.$$

The set $\mathfrak{C}$ has at least one element and at most $\lfloor \frac{n+1}{2} \rfloor$ elements. Let

$$c := \min(\mathfrak{C}) \text{ and } d := \max(\mathfrak{C}).$$

If $1 \in \mathfrak{C}$, then n is odd. Let $\mathfrak{S} := \{(r, s) \in \mathbb{N} \times \mathbb{N} | 2s - 1 \leq r\}$. For $(r, s) \in \mathfrak{S}$ let

$$c_{(r,s)} := \frac{\binom{r}{s-1}}{\binom{r}{s}} \text{ and } e_{(r,s)} := \binom{r+1}{s}.$$

6.2.1. General properties of MT pairs. Let $q, f_1, \tilde{f}_1$, and $\tilde{f}'_1$ be as in Definition 6.1 (b). Let (T_1, G_1) be an MT pair for $(f_1, \tilde{f}'_1, q)$. Let $\mathcal{W}_0$ be an irreducible $\tilde{H}_{1,\mathbb{Q}_q}$ -submodule of $W_1 \otimes \overline{\mathbb{Q}_q}$; it is also an irreducible $H_{1,\mathbb{Q}_q}$ -module. Let $\mathcal{G}_0$ and $\mathcal{H}_0$ be the images of $G_{1,\mathbb{Q}_q}$ and $H_{1,\mathbb{Q}_q}$ (respectively) in $\mathbf{GL}_{\mathcal{W}_0}$. We get a faithful representation $\rho_0 : \mathcal{G}_0 \hookrightarrow \mathbf{GL}_{\mathcal{W}_0}$. We list few basic properties of ρ_0 and $\mathcal{H}_0$.

(a) As the property 5.10 (a) holds for $f_2 = f_1$, we have $\dim_{\overline{\mathbb{Q}_q}}(\mathcal{W}_0) = n + 1$.

(b) The identity $\text{End}(W_1 \otimes_{\mathbb{Q}} \mathbb{Q}_q)^{G_1(\mathbb{Q}_q)} = \text{End}(W_1 \otimes_{\mathbb{Q}} \mathbb{Q}_q)^{H_{1,\mathbb{Q}_q}(\mathbb{Q}_q)}$ implies that the representation ρ_0 is irreducible.

(c) As the torus T_1 is Sh -good for $(\tilde{H}'_1, \tilde{X}'_1)$, the pair $(\mathcal{G}_0, \rho_0)$ is an irreducible weak Mumford–Tate pair of weight $\{0, 1\}$. More precisely, the group $\mathcal{G}_0$ is generated by images in $\mathbf{GL}_{\mathcal{W}_0}$ of cocharacters μ_0 of $\tilde{H}_{1,\mathbb{Q}_q} = \tilde{H}'_{1,\mathbb{Q}_q}$ whose extensions to $\mathbb{C}$ via an (any) embedding $\overline{\mathbb{Q}_q} \hookrightarrow \mathbb{C}$, belong to the set $\mathfrak{C}(\tilde{H}'_1, \tilde{X}'_1)$ introduced in Subsection 2.2. Thus all simple factors of $G_{1,\mathbb{Q}_q}^{\text{ad}}$ are of classical Lie type (cf. [Pi, Table 4.2]) and the pair $(\mathcal{G}_0, \rho_0)$ is non-trivial (cf. (a)).

(d) Each cocharacter μ_0 as in (c) acts on $\mathcal{W}_0$ via the identical and the trivial character of $\mathbb{G}_{m,\mathbb{Q}_q}$ and the pair of multiplicities $(m_{\text{id}}, m_{\text{triv}})$ of such characters either belongs to $\mathfrak{M}$ or is such that $\{m_{\text{id}}, m_{\text{triv}}\} = \{0, n + 1\}$, cf. (a) and (c). Therefore, if $d < 1$, then we have $m_{\text{id}} \neq m_{\text{triv}}$ and thus from the list of multiplicities in [Pi, Table 4.2] we get that all simple factors of $G_{1,\mathbb{Q}_q}^{\text{ad}}$ are of some A_m Lie type, where $m \in \{1, \dots, n\}$.

(e) As T_1 is defined over $\mathbb{Q}$, we can choose the cocharacter μ_0 of $\tilde{H}_{1,\mathbb{Q}_q} = \tilde{H}'_{1,\mathbb{Q}_q}$ such that either $(m_{\text{id}}, m_{\text{triv}})$ or $(m_{\text{triv}}, m_{\text{id}})$ is an arbitrary a priori given element of $\mathfrak{M}$.

(f) We assume that the group $\mathcal{G}_0^{\text{ad}}$ is simple and not of A_n Lie type. As $\mathcal{G}_0^{\text{ad}}$ is simple, the pair $(\mathcal{G}_0, \rho_0)$ is a non-trivial irreducible strong Mumford–Tate pair of weight $\{0, 1\}$. From this and [Pi, Table 4.2] we get that the set $\{(m_{\text{id}}, m_{\text{triv}}), (m_{\text{triv}}, m_{\text{id}})\}$ is uniquely determined by ρ_0 (i.e., it does not depend on μ_0). This implies that $\mathfrak{C}$ has only one element, cf. (e).

(g) The pair $(\mathcal{G}_0, \rho_0)$ is in the natural way a product of non-trivial irreducible strong Mumford–Tate pairs of weight $\{0, 1\}$ (cf. [Pi, Sect. 4, pp. 210–211]) which automatically involve groups whose adjoints are simple (cf. [Pi, Prop. 4.4]). If there exists $j \in J_1$ such that $\text{g.c.d.}(a_j, b_j) = 1$, then from (e) applied with $(m_{\text{id}}, m_{\text{triv}}) \in \{(a_j, b_j), (b_j, a_j)\}$ we get that this natural product does not have two or more irreducible factors and therefore the group $\mathcal{G}_0^{\text{ad}}$ is simple.

6.2.2. Lemma. *We assume that $d < 1$ and that (H_0, X_0) is not of non-special A_n type. Then for all elements $j \in J_1$ there exists a number $t_j \in \mathbb{N}$ and there exist numbers $r_{j,m} \in \mathbb{N}$, with $m \in \{1, \dots, t_j\}$, such that the following three properties hold:*

- (a) *the sum $\sum_{m=1}^{t_j} r_{j,m}$ does not depend on $j \in J_1$;*
- (b) *we have $\sum_{m=1}^{t_j} r_{j,m} < n$;*
- (c) *there exists a map $f(j): \{1, \dots, t_j\} \rightarrow \{0, 1\}$ that has the following four properties*
 - (c.i) *we have $n + 1 = \prod_{m \in \{1, \dots, t_j\}, f(j)(m)=0} (r_{j,m} + 1) \times \prod_{m \in \{1, \dots, t_j\}, f(j)(m)=1} e_{(r_{j,m}, s_{j,m})}$;*
 - (c.ii) *if $f(j)(m) = 0$, then $r_{j,m} + 1 = a_{j,m} + b_{j,m}$, with $a_{j,m}, b_{j,m} \in \mathbb{N}$ such that $c_{j,m} := \frac{a_{j,m}}{b_{j,m}} \in \mathfrak{C}$;*
 - (c.iii) *if $f(j)(m) = 1$, then there exists a pair $(r_{j,m}, s_{j,m}) \in \mathfrak{S}$ such that $2 \leq s_{j,m} \leq r_{j,m} - 2$ and $c_{j,m} := c_{(r_{j,m}, s_{j,m})} \in \mathfrak{C}$;*
 - (c.iv) *the map $q_j: \{1, \dots, t_j\} \rightarrow \mathfrak{C}$ that takes m to $c_{j,m}$ has the property that we have an inclusion $\mathfrak{C} \subseteq \text{Im}(q_j) \cup_{m \in \{1, \dots, t_j\}, f(j)(m)=0} \left\{ \frac{1}{r_{j,m}}, \frac{2}{r_{j,m}-1}, \dots, \frac{\lfloor \frac{r_{j,m}+1}{2} \rfloor}{r_{j,m}+1-\lfloor \frac{r_{j,m}+1}{2} \rfloor} \right\}$.*

Proof: See Subsection 2.2 for the totally real number field $F(H_0, X_0)$. We fix an identification of $\overline{\mathbb{Q}}$ with the algebraic closure of $\mathbb{Q}$ in $\overline{\mathbb{Q}_q}$. It allows us to identify naturally the set $\text{Hom}(F(H_0, X_0), \mathbb{R}) = \text{Hom}(F(H_0, X_0), \overline{\mathbb{Q}})$ with the set $\text{Hom}(F(H_0, X_0), \overline{\mathbb{Q}_q})$. This implies that we can identify the set J of simple factors of $H_{0,\mathbb{R}}$ with the set of simple factors of $H_{0,\overline{\mathbb{Q}_q}}$. To prove the Lemma we can use the notations of Definition 6.1 (b) and we can assume that the group $G_{1,\overline{\mathbb{Q}_q}}^{\text{ad}}$ does not have simple factors of A_n Lie type.

For $j \in J_1$ let t_j be the number of simple factors of the image $G_0(j)$ of $G_{1,\overline{\mathbb{Q}_q}}$ in the simple factor $H_0(j)$ of $H_{1,\overline{\mathbb{Q}_q}}^{\text{ad}} \xrightarrow{\sim} H_{0,\overline{\mathbb{Q}_q}}$ that corresponds to j . We will refer to these simple factors as the first, second, $\dots$, and the t_j -th simple factors of $G_0(j)$. For $m \in \{1, \dots, t_j\}$ let $r_{j,m}$ be the rank of the m -th simple factor of $G_0(j)$. Property (a) is implied by the fact that the images of $T_{1,\overline{\mathbb{Q}_q}}$ (and thus also of $G_{1,\overline{\mathbb{Q}_q}}$) in simple factors of $\tilde{H}_{1,\overline{\mathbb{Q}_q}}^{\text{ad}}$ have equal ranks. We check that (b) holds. As the group $G_{1,\overline{\mathbb{Q}_q}}^{\text{ad}}$ does not have simple factors of A_n Lie type, we have $G_0(j) \neq H_0(j)$. As $\mathcal{W}_0$ is a simple $\text{Lie}(G_0(j))$ -module and a simple

$\text{Lie}(H_0(j))$ -module, from Lemma 1.1.7 we get that the rank of $\text{Lie}(G_0(j))$ is less than the rank of $\text{Lie}(H_0(j))$ i.e., we have $\sum_{m=1}^{t_j} r_{j,m} < n$. Thus (b) holds

We define the map $f(j)$ in such a way that we have $f(j)(m) = 0$ if and only if the m -th simple factor of $G_0(j)$ is embedded in $H_0(j)$ via an SD-representation.

As $d < 1$, for each $m \in \{1, \dots, r_j\}$ there exists a number $r_{j,m} \in \mathbb{N}$ such that the m -th simple factor of $G_0(j)$ is of $A_{r_{j,m}}$ Lie type (cf. property 6.2.1 (d)). For $m \in \{1, \dots, t_j\}$ such that $f(j)(m) = 1$, let $s_{j,m} \in \mathbb{N}$ be such that $(r_{j,m}, s_{j,m}) \in \mathfrak{S}$ and the m -th simple factor of $G_0(j)$ is embedded in $H_0(j)$ via an irreducible representation associated to the minuscule weight $\varpi_{s_{j,m}}$ of the $A_{r_{j,m}}$ Lie type (cf. Corollary 1.1.9); as this representation is not an SD representation we have $2 \leq s_{j,m} \leq r_{j,m} - 2$. For $m \in \{1, \dots, t_j\}$ such that $f(j)(m) = 0$, let $a_{j,m}, b_{j,m} \in \mathbb{N}$ be numbers for which we have $a_{j,m} + b_{j,m} = r_{j,m} + 1$ and $a_{j,m} \leq \frac{r_{j,m}+1}{2}$ and for which there exists a cocharacter μ_0 as in the property 6.2.1 (c) whose image in the m -th simple factor of $G_0(j)$ has a centralizer in the m -th simple factor of $G_0(j)$ whose derived group is of $A_{a_{j,m}-1} + A_{b_{j,m}-1}$ Lie type (i.e., this image is defined by one of the cocharacters $\varpi_{a_{j,m}}^\vee$ or $\varpi_{b_{j,m}}^\vee$ of the $A_{r_{j,m}}$ Lie type); we emphasize that here we do not have a unique choice for the pair $(a_{j,m}, b_{j,m})$ (and this explains the $\cup$ sign in (c.iv)).

We check that (c) holds. Condition (c.i) is implied by the property 6.2.1 (a), cf. the fact that for $(r, s) \in \mathfrak{S}$ the irreducible representation associated to the minuscule weight ϖ_s of the A_r Lie type has dimension $e_{r,s}$. The fact that we have $c_{j,m} \in \mathfrak{C}$ for all $m \in \{1, \dots, t_j\}$ is implied by the first part of the property 6.2.1 (d), cf. the list of multiplicities in [Pi, Table 4.2]. Thus (c.ii) and (c.iii) hold. Condition (c.iv) is implied by the property 6.2.1 (e). Thus (c) also holds. $\square$

6.2.3. Remarks. (a) We assume that $c < d = 1$. If (H_0, X_0) is not of non-special A_n type, then one can list numerical conditions that are similar to the ones of Lemma 6.2.2. We do not stop to list them here. We only mention that (as in the property 6.2.2 (c)) we get maps q_j 's into $\mathfrak{C}$ which are “almost surjective” (cf. property 6.2.1 (e)) and that (as $d = 1$) the last sentence of the property 6.2.1 (d) does not a priori hold.

(b) If n is odd, then we can always identify $\tilde{f}'_1 = \tilde{f}_1$ (cf. property 4.1.1 (a), Subsubsection 4.1.2, and property 5.7 (c)).

6.2.4. Examples. We present some situations to which (the ideas of) Subsections 6.2.1 and 6.2.2 apply.

Example 1. The Shimura pair (H_0, X_0) is of non-special A_n type if there exists an element $j \in J_1$ such that $g.c.d.(a_j, b_j) = 1$ and $(a_j, b_j) \notin \{((\binom{r}{s-1}), (\binom{r}{s})) | (r, s) \in \mathfrak{S}, s \geq 2\}$. In particular, (H_0, X_0) is of non-special A_n type if there exists an element $j \in J_1$ such that $a_j = 1$ (i.e., if $\frac{1}{n} \in \mathfrak{C}$).

Example 2. We assume that $c < d$ and that there exists $j \in J_1$ such that $g.c.d.(a_j, b_j) = 1$. Then (H_0, X_0) is of non-special A_n type.

We argue Example 1 (resp. 2). As $g.c.d.(a_j, b_j) = 1$, the group $\mathcal{G}_0^{\text{ad}}$ is simple (cf. property 6.2.1 (g)). As $g.c.d.(a_j, b_j) = 1$ and as the group $\mathcal{G}_0^{\text{ad}}$ is simple, from the list of multiplicities in [Pi, Table 4.2] we get that $\mathcal{G}_0^{\text{ad}}$ is of A_m Lie type for some $m \in \{1, \dots, n\}$. But as $(a_j, b_j) \notin \{((\binom{r}{s-1}), (\binom{r}{s})) | (r, s) \in \mathfrak{S}, s \geq 2\}$ (resp. as $\mathfrak{C}$ has at least two elements), from the mentioned list (resp. from the property 6.2.1 (f)) we get that $m = n$. Thus $\mathcal{G}_0^{\text{ad}}$

has A_n Lie type and therefore it is $\mathcal{H}_0^{\text{ad}}$. We get that $G_{1, \mathbb{Q}_q}^{\text{ad}}$ has simple factors of A_n Lie type. Therefore (H_0, X_0) is of non-special A_n type.

Example 3. We assume that $c < d < 1$ and that there exists an element $j \in J_1$ such that $g.c.d.(a_j, b_j)$ does not have factors that are of the form $\tilde{a}_j + \tilde{b}_j$, with $\tilde{a}_j, \tilde{b}_j \in \mathbb{N}$ satisfying $\frac{\tilde{a}_j}{\tilde{b}_j} \in \mathfrak{C}$, or of the form $e_{(\tilde{r}_j, \tilde{s}_j)}$, with $(\tilde{r}_j, \tilde{s}_j) \in \mathfrak{S}$ satisfying $c_{(\tilde{r}_j, \tilde{s}_j)} \in \mathfrak{C}$. Then (H_0, X_0) is of non-special A_n type.

To check this, we show that the assumption that $\mathcal{G}_0^{\text{ad}}$ is not of A_n Lie type leads to a contradiction. The group $\mathcal{G}_0^{\text{ad}}$ is not simple (cf. property 6.2.1 (f)) and thus $(\mathcal{G}_0, \rho_0)$ is a tensor product of at least two non-trivial irreducible strong Mumford–Tate pairs of weight $\{0, 1\}$ (cf. property 6.2.1 (g)). From this and the property 6.2.1 (d), we get that ρ_0 is a tensor product of at least two irreducible representations associated to minuscule weights of some A_m Lie types. The dimensions of these irreducible representations are equal to one of the numbers $\tilde{a}_j + \tilde{b}_j$ or $e_{(\tilde{r}_j, \tilde{s}_j)}$ of the previous paragraph, cf. the list of multiplicities in [Pi, Table 4.2]. As at least one of these dimensions must divide $g.c.d.(a_j, b_j)$ (cf. property 6.2.1 (e)), we reached a contradiction. Thus (H_0, X_0) is of non-special A_n type.

Example 4. We assume that $c = d = 1$. If $n = 3$ or if 4 does not divide $n + 1$, then (H_0, X_0) is of non-special A_n type. To check this, we first remark that $c = d = 1$ implies that (H_0, X_0) is without involution (cf. Subsubsection 2.2.2). Thus we can appeal to Proposition 5.10 (b). With the notations of the beginning paragraph of Subsubsection 6.2.1, let $\mathcal{W}_0^d$ be a simple $H_{1, \mathbb{Q}_q}$ -submodule of $W_1 \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_q}$ which as an $H_{1, \mathbb{Q}_q}^{\text{der}}$ -module is dual to $\mathcal{W}_0$ and on which $Z^0(H_{1, \mathbb{Q}_q})$ acts in the same way as on $\mathcal{W}_0$. As $\frac{n+1}{4} \notin \mathbb{N} \setminus \{1\}$ and as $c = d = 1$, from the property 6.2.1 (c) and [Pi, Table 4.2] we get that the group $\mathcal{G}_0^{\text{ad}}$ is of $C_{\frac{n+1}{2}}$, $D_{\frac{n+1}{2}}$, $A_1 + A_1$, or some A_r Lie type. It is not of $C_{\frac{n+1}{2}}$, $D_{\frac{n+1}{2}}$, or $A_1 + A_1$ Lie type, as otherwise the $G_{1, \mathbb{Q}_q}$ -modules $\mathcal{W}_0$ and $\mathcal{W}_0^d$ would be isomorphic and thus we would have $\text{End}(W_1 \otimes_{\mathbb{Q}} \mathbb{Q}_q)^{G_1(\mathbb{Q}_q)} \neq \text{End}(W_1 \otimes_{\mathbb{Q}} \mathbb{Q}_q)^{H_{1, \mathbb{Q}_q}(\mathbb{Q}_q)}$. Thus $\mathcal{G}^{\text{ad}}$ is of some A_r Lie type. Let ϖ_s be the highest weight of the representation of $\text{Lie}(\mathcal{G}_0^{\text{ad}})$ on $\mathcal{W}_0$, cf. Corollary 1.1.9. If $r < n$, then as $c = d = 1$ we must have $c_{(r, s)} = 1$ and $n + 1 = e_{(r, s)}$. But we have $c_{(r, s)} = 1$ if and only if $r = 2s - 1$. If $r = 2s - 1 < n$, then the irreducible representation of $\text{Lie}(\mathcal{G}_0^{\text{ad}})$ of the highest weight ϖ_s is self dual (cf. [Bou2, pp. 188–189]) and this implies that the $G_{1, \mathbb{Q}_q}$ -modules $\mathcal{W}_0$ and $\mathcal{W}_0^d$ are isomorphic. Thus, in order not to get as above a contradiction with the identity $\text{End}(W_1 \otimes_{\mathbb{Q}} \mathbb{Q}_q)^{G_1(\mathbb{Q}_q)} = \text{End}(W_1 \otimes_{\mathbb{Q}} \mathbb{Q}_q)^{H_{1, \mathbb{Q}_q}(\mathbb{Q}_q)}$, we must have $r = n$. Therefore $\mathcal{G}^{\text{ad}}$ is of A_n Lie type. Thus (H_0, X_0) is of non-special A_n type.

Example 5. We assume that $c < d = 1$ and (H_0, X_0) is not of non-special A_n type. As $d = 1$, $n + 1$ is even. The pair $(\mathcal{G}_0, \rho_0)$ is a product of at least two non-trivial irreducible strong Mumford–Tate pairs of weight $\{0, 1\}$ (cf. properties 6.2.1 (f) and (g)) such that at least one of them is even dimensional (as $n + 1$ is even). If $(\mathcal{G}_0, \rho_0)$ is a product of even dimensional irreducible strong Mumford–Tate pairs of weight $\{0, 1\}$ (resp. if $(\mathcal{G}_0, \rho_0)$ is a product of non-trivial irreducible strong Mumford–Tate pairs of weight $\{0, 1\}$ such that at least one is odd dimensional and at least one is even dimensional), then from the property 6.2.1 (e) (resp. from the generation part of the property 6.2.1 (c) and from the property 6.2.1 (d)), we get that for all $j \in J_1$ such that $c_j < 1$, 2 divides both a_j and b_j (resp. we get that there exists $j \in J_1$ such that $c_j < 1$ and 2 divides both a_j and b_j). Thus if $n + 1$

is a power of 2, then for all $j \in J_1$ such that $c_j < 1$, 2 divides both a_j and b_j .

Example 6. If $n + 1$ is either 4 or a prime, then (H_0, X_0) is of non-special A_n type (regardless of what the set $\mathfrak{C}$ is). This is an elementary exercise which follows from Examples 1 and 4.

We do not stop here to restate Lemma 6.2.2 in terms of n , similar to the statements of Examples 4 and 6. Example 1.3.5 (i) (resp. 1.3.5 (iii)) is a particular case of Example 2 (resp. 5); moreover, Example 1.3.5 (ii) (resp. 1.3.5 (iv)) is only Example 1 (resp. 6).

7. The proof of the Main Theorem

Theorems 7.1 and 7.2 are the very essence of the proofs of Theorems 1.3.4 and 1.3.6 which are completed in Subsection 7.3. Corollary 7.4 is an application to the $D_4^{\mathbb{H}}$ type of the proofs of Theorems 7.1 and 7.2. Subsection 7.5 lists the “simplest” cases not settled by Theorem 1.3.4. Theorem 7.6 proves a stronger variant of Theorem 1.3.7 for direct sums of abelian motives in the sense of [DM]. We end with remarks in Subsection 7.7.

We use $+$ to denote the Lie type of products of split, absolutely simple, adjoint groups. Until the end we use the notations of Subsections 1.1 and of the paragraphs before Theorems 1.1.5 and 1.3.2. Let W_A , λ_A , ψ_A , and f_A be as in Subsection 1.5.

7.1. Theorem. *We assume that for each element $j \in J$ the adjoint Shimura pair (H_j, X_j) is of either non-inner $D_4^{\mathbb{H}}$ type or a type listed in Theorem 1.3.4. Then for each element $j \in J$, the group $G_{\overline{\mathbb{Q}_p}}$ surjects onto a simple factor of $H_{j, \overline{\mathbb{Q}_p}}$.*

Proof: To prove the Theorem we can assume that λ_A is a principal polarization of A (cf. property 3.1 (a) and [Mu2, Ch. IV, Sect. 23, Cor. 1]) and that the adjoint group H_A^{ad} is non-trivial (cf. property 3.4.1 (a)). Thus the set J that indexes the simple factors of $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is non-empty. To prove the Theorem we can also assume that the set J has precisely one element, cf. Proposition 5.4.1 and Subsubsection 5.4.3. Thus $(H_j, X_j) = (H_A^{\text{ad}}, X_A^{\text{ad}})$. We take v such that it is unramified over l .

Let the triple $(\mathcal{W}_0, \mathcal{G}_0, \mathcal{H}_0)$ be as before Subsubsection 1.4.1. The groups $\mathcal{G}_0^{\text{ad}}$ and $\mathcal{H}_0^{\text{ad}}$ are isomorphic with simple factors of $G_{\overline{\mathbb{Q}_p}}^{\text{ad}}$ and $H_{A, \overline{\mathbb{Q}_p}}^{\text{ad}}$ (respectively). Thus if $\mathcal{G}_0^{\text{ad}} = \mathcal{H}_0^{\text{ad}}$, then $G_{\overline{\mathbb{Q}_p}}^{\text{ad}}$ surjects onto a simple factor of $H_{A, \overline{\mathbb{Q}_p}}^{\text{ad}}$. We are left to check that:

(*) we can choose the simple $H_{A, \overline{\mathbb{Q}_p}}$ -module $\mathcal{W}_0$ such that we have $\mathcal{G}_0^{\text{ad}} = \mathcal{H}_0^{\text{ad}}$.

To check (*) we will consider one by one the cases (a) to (e) of Theorem 1.3.4, with the non-inner $D_4^{\mathbb{H}}$ being added to the case (d’). Only for the case (d’) we will have to make a good choice of $\mathcal{W}_0$.

(a) We assume that $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of non-special A_n type. To check the property (*) we can assume that $E^{\text{conn}} = E$. To check the property (*) we can also assume that the injective map $f_A : (H_A, X_A) \hookrightarrow (\mathbf{GSp}(W_A, \psi_A), S_A)$ of Shimura pairs factors through an injective map $\tilde{f}_A : (\tilde{H}_A, \tilde{X}_A) \hookrightarrow (\mathbf{GSp}(W_A, \psi_A), S_A)$ of Shimura pairs which is a PEL type embedding such that the monomorphism $H_{A, \overline{\mathbb{Q}}}^{\text{der}} \hookrightarrow \tilde{H}_{A, \overline{\mathbb{Q}}}^{\text{der}}$ is (cf. Proposition 5.4.1, Subsubsection 5.4.2, Proposition 4.1, and Subsubsection 4.1.2): (i) an isomorphism if

either $n = 1$ or $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is with involution, and (ii) is a diagonal embedding if $n \geq 2$ and $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is without involution. Based on Definition 6.1 (c), we can also assume that there exists a prime $q \in \mathbb{N}$ such that for each twist $\tilde{f}'_A$ of $\tilde{f}_A$ via a locally trivial class in $H^1(\mathbb{Q}, \tilde{H}_A)$ and for every MT pair (T_1, G_1) for $(f_A, \tilde{f}'_A, q)$, the group $G_{1, \overline{\mathbb{Q}}_q}^{\text{ad}}$ has simple factors of A_n Lie type.

We choose v such that we also have $k(v) = \mathbb{F}_l$, $l \notin \{p, q\}$, l is greater than the number $N(A)$ of Theorem 1.3.2, and (cf. Theorem 1.1.5) T_v has the same rank as $G_{\mathbb{Q}_p}$. The torus $T_{v, \mathbb{Q}_p}$ of $G_{\mathbb{Q}_p}$ is $\tilde{H}_A(\mathbb{Q}_p)$ -conjugate to the extension to $\mathbb{Q}_p$ of a torus T_4 of $\tilde{H}_A$ that is naturally identifiable with T_v (cf. Subsubsection 5.6.3 applied with $G_2 = G_4 = \tilde{H}_A$). Strictly speaking, we ought to substitute in the last sentence $\tilde{H}_A$ by a suitable form of it that is locally isomorphic. But from the point of view of what follows, such a substitution is irrelevant. Thus we will not complicate notations by introducing explicitly a form of $\tilde{H}_A$. A suitable $\tilde{H}_A(\overline{\mathbb{Q}}_l)$ -conjugate of $T_{4, \overline{\mathbb{Q}}_l}$ is a subtorus of $G_{\overline{\mathbb{Q}}_l}$, cf. Subsubsection 5.6.4 (applied in the context of A instead of B). Thus from Corollary 3.4.2 we get that $T_{4, \overline{\mathbb{Q}}_l}$ is generated by images of cocharacters in the set $\mathfrak{C}_l(\tilde{H}_A, \tilde{X}_A)$ of cocharacters of $\tilde{H}_{A, \overline{\mathbb{Q}}_l}$ whose extensions to $\mathbb{C}$ via an embedding $i_{\overline{\mathbb{Q}}_l} : \overline{\mathbb{Q}}_l \hookrightarrow \mathbb{C}$ that extends the embedding i_E of Subsection 1.1, belong to the set $\mathfrak{C}(\tilde{H}_A, \tilde{X}_A)$ defined in Subsection 2.2. This implies that the torus $T_1 := T_4$ is Sh -good for $(\tilde{H}_A, \tilde{X}_A)$. As the groups T_1 and $G_{\mathbb{Q}_p}$ have the same ranks, from Theorem 1.1.4 we get that $(T_1, G_{\mathbb{Q}_p})$ is an MT pair for $(f_A, \tilde{f}_A, p)$.

If p is q , then $G_{\mathbb{Q}_p}^{\text{ad}}$ has simple factors of A_n Lie type and thus the image of $T_{1, \overline{\mathbb{Q}}}$ in each simply factor of $\tilde{H}_{A, \overline{\mathbb{Q}}}^{\text{ad}}$ has rank n . This last property implies that, regardless of the fact that p is not or is q , the images of $G_{\mathbb{Q}_p}$ in the simply factors of $H_{A, \mathbb{Q}_p}^{\text{ad}}$ (equivalently the images of $T_{1, \overline{\mathbb{Q}}_p}$ in the simply factors of $\tilde{H}_{A, \mathbb{Q}_p}^{\text{ad}}$) have rank n . From this and Lemma 1.1.7 applied in the context of the Lie monomorphisms $\text{Lie}(\mathcal{G}_0^{\text{ad}}) \hookrightarrow \text{Lie}(\mathcal{H}_0^{\text{ad}}) \hookrightarrow \text{End}(\mathcal{W}_0)$, we get that $\text{Lie}(\mathcal{G}_0^{\text{ad}}) = \text{Lie}(\mathcal{H}_0^{\text{ad}})$. Thus $\mathcal{G}_0^{\text{ad}} = \mathcal{H}_0^{\text{ad}}$ (i.e., the property $(*)$ holds).

Based on Proposition 5.4.1, for the below cases (b) to (e) we can assume that f_A factors through an injective map $f_2 : (G_2, X_2) \rightarrow (\mathbf{GSp}(W_1, \psi_1), S_1) = (\mathbf{GSp}(W_A, \psi_A), S_A)$ of Shimura pairs for which the properties 5.10 (a) and (b) hold and for which the monomorphism $H_A \hookrightarrow G_2$ induces an identity $H_A^{\text{der}} = G_2^{\text{der}}$ at the level of derived groups.

As the representation ρ_0 is irreducible, from Theorem 1.1.8 we get that we have a short exact sequence $1 \rightarrow Z(\mathbf{GL}_{\mathcal{W}_0}) \rightarrow \mathcal{G}_0 \rightarrow \mathcal{G}_0^{\text{ad}} \rightarrow 1$.

(b) If $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of B_n type, then $\mathcal{H}_0$ is a $\mathbf{GSpin}_{2n+1, \overline{\mathbb{Q}}_p}$ group and $\mathcal{W}_0$ is its spin representation. As ρ_0 is irreducible, from [Pi, Prop. 4.3] we get that $\mathcal{G}_0 = \mathcal{H}_0$. Thus the property $(*)$ holds.

(c) If $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of C_n type, then $\mathcal{H}_0$ is a $\mathbf{GSp}_{2n, \overline{\mathbb{Q}}_p}$ group and $\mathcal{W}_0$ is of dimension $2n$. We assume that n is odd. Thus $2n$ can not be written as a product of two or more even numbers. As the pair $(\mathcal{G}_0, \rho_0)$ is an irreducible weak Mumford–Tate pair of weight $\{0, 1\}$ and as the representation ρ_0 is symplectic, the pair $(\mathcal{G}_0, \rho_0)$ is a product of irreducible strong Mumford–Tate pairs of weight $\{0, 1\}$ that involve (orthogonal or symplectic and thus) even dimensional irreducible representations. From the last two sentences we get

that $(\mathcal{G}_0, \rho_0)$ is a strong Mumford–Tate pair of weight $\{0, 1\}$. From this and [Pi, Prop. 4.7 or Table 4.2] we get $\mathcal{H}_0 = \mathcal{H}_0$. Thus the property $(*)$ holds.

(d) We assume that $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of $D_n^{\mathbb{H}}$ type with n odd and such that $2n \notin \{\binom{2m+1}{2m} | m \in \mathbb{N}\}$. This case is the orthogonal variant of (c). The group $\mathcal{H}_0$ is a $\mathbf{GSO}_{2n, \overline{\mathbb{Q}_p}}$ group and $\mathcal{W}_0$ is of dimension $2n$. As the representation ρ_0 is irreducible and orthogonal, it is a tensor product of (orthogonal or symplectic and thus of) even dimensional irreducible representations. Thus, as n odd, the group $\mathcal{G}_0^{\text{ad}}$ is simple. As the number $2n \in 2\mathbb{N} + 2$ is not of the form $\binom{2m+1}{2m}$, with $m \in \mathbb{N}$, it is also not of the form $\binom{4m}{2m}$, with $m \in \mathbb{N}$. From this and the fact that n is not of the form 2^m with $m \geq 2$, we get that $(\mathcal{G}_0, \rho_0) = (\mathcal{H}_0, \rho_0)$ (cf. [Pi, Table 4.2]). Thus the property $(*)$ holds.

(d') We assume that $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of non-inner $D_{2n}^{\mathbb{H}}$ type, with n a prime. The group $\mathcal{H}_0$ is a $\mathbf{GSO}_{4n, \overline{\mathbb{Q}_p}}$ group and $\mathcal{W}_0$ is of dimension $4n$.

We have $[I(H_A^{\text{ad}}, X_A^{\text{ad}}) : F(H_A^{\text{ad}}, X_A^{\text{ad}})] = 2$ (as in the proof of Lemma 4.4 one argues that for $n = 2$ the degree $[I(H_A^{\text{ad}}, X_A^{\text{ad}}) : F(H_A^{\text{ad}}, X_A^{\text{ad}})]$ is not divisible by 3). Based on the independence part of Corollary 1.3.3 and on Lemma 1.1.7, we can assume that p is such that the number fields $I(H_A^{\text{ad}}, X_A^{\text{ad}})$ and E are unramified over p , that $H_{A, \mathbb{Q}_p}^{\text{ad}}$ is unramified, and that (cf. the definition of the non-inner type and [La, p. 168]) there exists a prime w of $F(H_A^{\text{ad}}, X_A^{\text{ad}})$ which divides p and which does not split in $I(H_A^{\text{ad}}, X_A^{\text{ad}})$. We also take the prime v such that $l = p$; thus v is unramified over p . Let $F_{p,1} := F(H_A^{\text{ad}}, X_A^{\text{ad}})_w$. As w does not split in $I(H_A^{\text{ad}}, X_A^{\text{ad}})$ and as $H_{A, \mathbb{Q}_p}^{\text{ad}}$ is unramified, there exists a non-split, absolutely simple factor G_2 of $H_{A, F_{p,1}}^{\text{ad}}$. The group G_2 is unramified and splits over the unramified quadratic extension $F_{p,2} := I(H_A^{\text{ad}}, X_A^{\text{ad}}) \otimes_{F(H_A^{\text{ad}}, X_A^{\text{ad}})} F_{p,1}$ of $F_{p,1}$. We identify $\overline{F_{p,2}} = \overline{\mathbb{Q}_p}$. Let G_3 be the image of $G_{F_{p,1}}$ in G_2 .

We choose $\mathcal{W}_0$ such that $\text{Lie}(G_{2, \overline{\mathbb{Q}_p}})$ acts non-trivially on it. Thus we have natural identifications $G_{2, \overline{\mathbb{Q}_p}} = \mathcal{H}_0^{\text{ad}}$ and $G_{3, \overline{\mathbb{Q}_p}} = \mathcal{G}_0^{\text{ad}}$. Let μ_2 be a non-trivial cocharacter of $G_{2, \overline{\mathbb{Q}_p}} = \mathcal{H}_0^{\text{ad}}$ that is the image of a cocharacter of $\mathcal{H}_0$ which acts on $\mathcal{W}_0$ via the trivial and the identity characters of $\mathbb{G}_{m, \overline{\mathbb{Q}_p}}$. As $\mathcal{W}_0$ has dimension $4n$ and as $n \geq 2$, the centralizer of μ_2 in $G_{2, \overline{\mathbb{Q}_p}}$ has a derived subgroup of A_{2n-1} Lie type. Based on Theorem 1.1.8, we can choose μ_2 such that it factors through $G_{3, \overline{\mathbb{Q}_p}}$. As the centralizer of μ_2 in $G_{2, \overline{\mathbb{Q}_p}}$ has a derived subgroup of A_{2n-1} Lie type, the group $\text{Gal}(F_{p,1})$ does not fix $[\mu_2]$ (this holds even if $n = 2$, as the non-split group G_2 splits over $F_{p,2} = I(H_A^{\text{ad}}, X_A^{\text{ad}}) \otimes_{F(H_A^{\text{ad}}, X_A^{\text{ad}})} F_{p,1}$).

The representation of $\text{Lie}(\mathcal{G}_0^{\text{der}})$ on $\mathcal{W}_0$ is a tensor product of N irreducible representations that are associated to minuscule weights (cf. Corollary 1.9) and that are either symplectic or orthogonal, the number N_S of symplectic ones being even. As n is either odd or 2, $4n$ is not divisible by 16. From the last two sentences we get that $N \in \{1, 2, 3\}$. If $N = 3$, then $n = 2$ and therefore $N_S = N = 3$; this contradicts the fact that N_S is even. Thus $N \in \{1, 2\}$.

We show that the assumption $G_3 \neq G_2$ leads to a contradiction. As n is a prime, we have $4n \notin \{\binom{4m}{2m} | m \in \mathbb{N}\}$. Thus from [Pi, Table 4.2] we get that either $N \neq 1$ or $n = 2N = 2$ and $\mathcal{G}_0^{\text{ad}}$ is of B_3 Lie type. We first assume that $N = 2$. The only way to factor $4n$ as a product of two even numbers is $2 \times 2n$. Thus $2 \geq N_S \geq 1$ and therefore $N_S = 2$. Thus $\mathcal{G}_0^{\text{ad}}$ is of $C_1 + C_n$ Lie type (to be compared with (c)) and therefore

each simple factor of G_3 is absolutely simple. Each group pair over $F_{p,1}$ that involves a product of absolutely simple, adjoint groups that over $\overline{\mathbb{Q}_p}$ are of some C_m Lie types, has the property that its conjugacy class of cocharacters is fixed by $\text{Gal}(F_{p,1})$. Thus $\text{Gal}(F_{p,1})$ fixes $[\mu_2]$. Contradiction. If $n = 2N = 2$, then the same arguments lead to a contradiction: we only have to replace the $C_1 + C_m$ Lie type with the B_3 Lie type. Therefore $G_3 = G_2$ (i.e., $\mathcal{G}_0^{\text{ad}} = \mathcal{H}_0^{\text{ad}}$). Thus the property (*) holds.

(e) We assume that $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of $D_n^{\mathbb{R}}$ type. We show that the assumption $\mathcal{G}_0 \neq \mathcal{H}_0$ leads to a contradiction. The faithful representation $\mathcal{H}_0^{\text{der}} \hookrightarrow \mathbf{GL}_{\mathcal{W}_0}$ is a half spin representation. From this and [Pi, Prop. 4.3] we get that $\mathcal{G}_0^{\text{ad}}$ is of $B_{n_1} + B_{n-n_1-1}$ Lie type, where $n_1 \in \{0, \dots, n-1\}$ (the B_0 Lie type corresponds to the trivial group).

Let $\mathcal{F}_0$ be the simple factor of $H_{A, \overline{\mathbb{Q}_p}}^{\text{ad}}$ that is naturally isomorphic to $\mathcal{H}_0^{\text{ad}}$. Let $\mathcal{F}_1$ be the image of $G_{\overline{\mathbb{Q}_p}}$ in $\mathcal{F}_0$. We identify naturally $\mathcal{F}_1 = \mathcal{G}_0^{\text{ad}}$. There exists an $H_{A, \overline{\mathbb{Q}_p}}$ -submodule $\mathcal{W}_1$ of $W_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_p}$ such that the $Z^0(H_{A, \overline{\mathbb{Q}_p}})$ -modules $\mathcal{W}_0$ and $\mathcal{W}_1$ are isomorphic and the representation of $\text{Lie}(\mathcal{F}_0)$ on $\mathcal{W}_0 \oplus \mathcal{W}_1$ is the spin representation, cf. Proposition 5.10 (b). The $H_{A, \overline{\mathbb{Q}_p}}$ -modules $\mathcal{W}_0$ and $\mathcal{W}_1$ are not isomorphic. The representations of $\text{Lie}(\mathcal{F}_1)$ on $\mathcal{W}_0$ and $\mathcal{W}_1$ are isomorphic to the tensor product of the spin representations of the two simple factors of $\text{Lie}(\mathcal{F}_1)$, cf. [Pi, Prop. 4.3]. As $Z^0(G_{\overline{\mathbb{Q}_p}}) = Z^0(H_{A, \overline{\mathbb{Q}_p}})$ (cf. Theorem 1.3.1) and as the representation of $\text{Lie}(H_{A, \overline{\mathbb{Q}_p}}^{\text{ad}})$ on $\mathcal{W}_0 \oplus \mathcal{W}_1$ factors through $\text{Lie}(\mathcal{F}_0)$ (cf. Proposition 5.10 (a)), we get that the $G_{\overline{\mathbb{Q}_p}}$ -modules $\mathcal{W}_0$ and $\mathcal{W}_1$ are isomorphic. Thus the $H_{A, \overline{\mathbb{Q}_p}}$ -modules $\mathcal{W}_0$ and $\mathcal{W}_1$ are isomorphic, cf. Theorem 1.1.4. Contradiction. Therefore $\mathcal{G}_0 = \mathcal{H}_0$. Thus the property (*) holds. $\square$

7.2. Theorem. *We assume that the following two conditions hold:*

- (i) *there exists no element $j \in J$ such that (H_j, X_j) is of $D_4^{\mathbb{H}}$ type;*
- (ii) *for each element $j \in J$ the group $G_{\overline{\mathbb{Q}_p}}$ surjects onto a simple factor of $H_{j, \overline{\mathbb{Q}_p}}$.*

Then we have $G_{\overline{\mathbb{Q}_p}}^{\text{ad}} = H_{A, \overline{\mathbb{Q}_p}}^{\text{ad}}$.

Proof: We choose v such that: it is unramified over a prime $l \in \mathbb{N}$ different from p and greater than the number $N(A)$ of Theorem 1.3.2 and moreover the ranks of T_v and $G_{\overline{\mathbb{Q}_p}}$ are equal. From (ii) and Theorem 1.3.2 we get that the image of $G_{\overline{\mathbb{Q}_p}}$ in an arbitrary simple factor $\mathcal{F}_0$ of $H_{A, \overline{\mathbb{Q}_p}}^{\text{ad}}$ has the same rank as $\mathcal{F}_0$. Let $\tilde{H}_{A, \overline{\mathbb{Q}_p}}$ be the subgroup of $H_{A, \overline{\mathbb{Q}_p}}$ that contains $Z^0(H_{A, \overline{\mathbb{Q}_p}})$ and whose image in $H_{A, \overline{\mathbb{Q}_p}}^{\text{ad}}$ is the product of the images of $G_{\overline{\mathbb{Q}_p}}$ in the simple factors of $H_{A, \overline{\mathbb{Q}_p}}^{\text{ad}}$. We have $G_{\overline{\mathbb{Q}_p}} \leq \tilde{H}_{A, \overline{\mathbb{Q}_p}} \leq H_{A, \overline{\mathbb{Q}_p}}$ and $\tilde{H}_{A, \overline{\mathbb{Q}_p}}$ has the same rank as $H_{A, \overline{\mathbb{Q}_p}}$. The centralizers of $\tilde{H}_{A, \overline{\mathbb{Q}_p}}$ and $H_{A, \overline{\mathbb{Q}_p}}$ in $\mathbf{GL}_{W_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_p}}$ coincide, cf. Theorem 1.1.4. From the last two sentences we get that $\text{Lie}(\tilde{H}_{A, \overline{\mathbb{Q}_p}}) = \text{Lie}(H_{A, \overline{\mathbb{Q}_p}})$, cf. Lemma 1.1.7. Thus $\tilde{H}_{A, \overline{\mathbb{Q}_p}} = H_{A, \overline{\mathbb{Q}_p}}$ and therefore $G_{\overline{\mathbb{Q}_p}}$ surjects onto $\mathcal{F}_0$. This implies that:

- (iii) *For each simple factor $\mathcal{F}_0$ of $H_{A, \overline{\mathbb{Q}_p}}^{\text{ad}}$, there exists a unique simple factor of $G_{\overline{\mathbb{Q}_p}}^{\text{ad}}$ that surjects (isomorphically) onto $\mathcal{F}_0$.*

We now start to check that we have an identity $G_{\mathbb{Q}_p}^{\text{ad}} = H_{A, \mathbb{Q}_p}^{\text{ad}}$. Based on the property 3.1 (c) and on (iii), the identity $G_{\mathbb{Q}_p}^{\text{ad}} = H_{A, \mathbb{Q}_p}^{\text{ad}}$ is equivalent to the fact that no simple factor of $G_{\mathbb{Q}_p}^{\text{ad}}$ surjects onto two or more simple factors of $H_{A, \mathbb{Q}_p}^{\text{ad}}$. To check this last statement, we can assume that (cf. Subsubsections 5.4.1 to 5.4.3 and Subsection 5.2):

- (iv) all simple factors of either $H_{A, \mathbb{Q}}^{\text{ad}}$ or $G_{\mathbb{Q}_p}^{\text{ad}}$ have the same Lie type $\mathfrak{L}$,
- (v) and f_A factors through an injective map

$$f_A^{(2)} : (H_A^{(2)}, X_A^{(2)}) \rightarrow (\mathbf{GSp}(W_A, \psi_A), S_A)$$

of Shimura pairs for which the properties 5.10 (a) and (b) hold, with $(H_A^{(2), \text{ad}}, X_A^{(2), \text{ad}}) = (H_A^{\text{ad}}, X_A^{\text{ad}})$ and with $f_A^{(2)}$ as a Hodge quasi product $\times_{j \in J}^{\mathcal{H}} f_j^{(2)}$ of injective maps

$$f_j^{(2)} : (H_j^{(2)}, X_j^{(2)}) \hookrightarrow (\mathbf{GSp}(W_j, \psi_j), S_j)$$

of Shimura pairs; here $j \in J$ and $(H_j^{(2)}, X_j^{(2)})$ is such that we can identify $(H_j^{(2), \text{ad}}, X_j^{(2), \text{ad}}) = (H_j, X_j)$.

Let $\mathcal{F}_1$ be a simple factor of $G_{\mathbb{Q}_p}^{\text{ad}}$.

7.2.1. Example. We first treat the following particular case. We assume that all simple factors of $H_{A, \mathbb{R}}^{\text{ad}}$ are non-compact and that all simple factors of $(H_A^{\text{ad}}, X_A^{\text{ad}})$ are of $D_n^{\mathbb{R}}$ type. Thus we can also assume that $Z^0(H_A) = Z(\mathbf{GL}_W) \xrightarrow{\sim} \mathbb{G}_{m, \mathbb{Q}}$, cf. Propositions 5.10 (c) and 5.4.1. We assume that $\mathcal{F}_1$ surjects onto two distinct simple factors $\mathcal{F}'_0$ and $\mathcal{F}''_0$ of $H_{A, \mathbb{Q}_p}^{\text{ad}}$. From Proposition 5.10 (a) and (b) we get that there exist four non-isomorphic, simple $\text{Lie}(H_{A, \mathbb{Q}_p}^{\text{ad}})$ -submodules $\mathcal{W}'_0, \mathcal{W}''_0, \mathcal{W}'_1$, and $\mathcal{W}''_1$ of $W_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_p}$ with the property that for $* \in \{\iota, \iota'\}$, the $\text{Lie}(\mathcal{F}_0^*)$ -modules $\mathcal{W}_0^*$ and $\mathcal{W}_1^*$ are simple and associated to the two half spin representations of $\text{Lie}(\mathcal{F}_0^*)$. We can assume that these four $\text{Lie}(H_{A, \mathbb{Q}_p}^{\text{ad}})$ -submodules are indexed in such a way that the $\text{Lie}(\mathcal{F}_1)$ -module $\mathcal{W}'_0$ and $\mathcal{W}'_1$ are isomorphic. As $Z^0(G_{\mathbb{Q}_p}) = Z^0(H_{A, \mathbb{Q}_p}) = Z(\mathbf{GL}_{W_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_p}})$, $\mathcal{W}'_0$ and $\mathcal{W}'_1$ are also isomorphic as $Z^0(G_{\mathbb{Q}_p})$ -modules. Thus the simple $G_{\mathbb{Q}_p}$ -modules $\mathcal{W}'_0$ and $\mathcal{W}'_1$ are isomorphic. Therefore the centralizers of $G_{\mathbb{Q}_p}$ and $H_{A, \mathbb{Q}_p}$ in $\mathbf{GL}_{W_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_p}}$ are distinct and this contradicts Theorem 1.1.4. Thus $G_{\mathbb{Q}_p}^{\text{ad}} = H_{A, \mathbb{Q}_p}^{\text{ad}}$. We come back to the general case.

7.2.2. General construction. In order to show that no simple factor of $G_{\mathbb{Q}_p}^{\text{ad}}$ surjects onto two or more simple factors of $H_{A, \mathbb{Q}_p}^{\text{ad}}$, we need to have a very good understanding of the injective map $f_A^{(2)}$ and of the action of the torus $Z^0(H_A^{(2)})$ on W_A . Thus next we will list some basic properties of $f_A^{(2)}$. Based on Variant 4.9 (b) we can assume that $f_A^{(2)}$ is such that there exists a totally real number subfield F of $\overline{\mathbb{Q}}$ with the property that each $f_j^{(2)}$ factors through the composite of two injective maps

$$(H_j^{(3)}, X_j^{(3)}) \hookrightarrow (H_j^{(4)}, X_j^{(4)}) \xrightarrow{f_j^{(4)}} (\mathbf{GSp}(W_j, \psi_j), S_j)$$

of Shimura pairs for which the following three properties hold.

(i) The group $H_j^{(3),\text{ad}}$ is the $\text{Res}_{F/\mathbb{Q}}$ of an absolutely simple, adjoint group over F whose extension to $\overline{\mathbb{Q}}$ is of $\mathfrak{L}$ Lie type. Moreover the monomorphism $H_j^{(2)} \hookrightarrow H_j^{(3)}$ gives birth to an injective map $(H_j^{(2),\text{ad}}, X_j^{(2),\text{ad}}) \hookrightarrow (H_j^{(3),\text{ad}}, X_j^{(3),\text{ad}})$ of Shimura pairs which is obtained as in (1) of Subsubsection 2.2.3.

(ii) The group $H_j^{(4),\text{ad}}$ is the $\text{Res}_{F/\mathbb{Q}}$ of an absolutely simple, adjoint group over F whose extension to $\overline{\mathbb{Q}}$ is of $\mathfrak{L}_4$ Lie type. Here $\mathfrak{L}_4$ is $\mathfrak{L}$ if $\mathfrak{L} = A_n$, is A_{2n-1} if (H_j, X_j) is of C_n or $D_n^{\mathbb{H}}$ type, and is A_{2n-1} if (H_j, X_j) is of B_n or $D_n^{\mathbb{R}}$ type. If $\mathfrak{L} \neq A_n$, then the monomorphism $H_{j,\overline{\mathbb{Q}}}^{(3),\text{der}} \hookrightarrow H_{j,\overline{\mathbb{Q}}}^{(4),\text{der}}$ is a product of $[F : \mathbb{Q}]$ monomorphisms associated naturally to homomorphisms of the form $h_{\overline{\mathbb{Q}}}$, where h is as in the property 4.3.1 (c). If $\mathfrak{L} = A_n$ with $n \geq 2$ and if (H_j, X_j) is without involution (thus $(H_j^{(3),\text{ad}}, X_j^{(3),\text{ad}})$ and $(H_j^{(4),\text{ad}}, X_j^{(4),\text{ad}})$ are also without involution), then the monomorphism $H_{j,\overline{\mathbb{Q}}}^{(3),\text{der}} \hookrightarrow H_{j,\overline{\mathbb{Q}}}^{(4),\text{der}}$ is a diagonal embedding (cf. Subsubsection 4.1.2). If either $\mathfrak{L} = A_n$ and (H_j, X_j) is with involution (thus $(H_j^{(3),\text{ad}}, X_j^{(3),\text{ad}})$ and $(H_j^{(4),\text{ad}}, X_j^{(4),\text{ad}})$ are also with involution) or $\mathfrak{L} = A_1$, then the monomorphism $H_{j,\overline{\mathbb{Q}}}^{(3),\text{der}} \hookrightarrow H_{j,\overline{\mathbb{Q}}}^{(4),\text{der}}$ is an isomorphism.

(iii) The map $f_j^{(4)}$ is a PEL type embedding constructed as in Proposition 4.1.

The map $f_A^{(2)}$ factors through $f_A^{(4)} := \times_{j \in J} f_j^{(4)} : (H_A^{(4)}, X_A^{(4)}) \hookrightarrow (\mathbf{GSp}(W_A, \psi_A), S_A)$. From properties (i), (ii), and 7.2 (iii) we easily get that:

(iv) Two simple factors of either $\text{Lie}(H_{A,\overline{\mathbb{Q}_p}}^{\text{ad}}) = \text{Lie}(H_{A,\overline{\mathbb{Q}_p}}^{(2),\text{ad}})$ or $\oplus_{j \in J} \text{Lie}(H_{j,\overline{\mathbb{Q}_p}}^{(3),\text{ad}})$ do not map injectively into the same simple factor of $\text{Lie}(H_{A,\overline{\mathbb{Q}_p}}^{(4),\text{ad}})$.

(v) Two simple factors of $\text{Lie}(G_{\overline{\mathbb{Q}_p}}^{\text{ad}})$ do not map injectively into the same simple factor of $\text{Lie}(H_{A,\overline{\mathbb{Q}_p}}^{(4),\text{ad}})$. Moreover all monomorphisms from a simple factor of $\text{Lie}(G_{\overline{\mathbb{Q}_p}}^{\text{ad}})$ into a simple factor $\text{Lie}(H_{A,\overline{\mathbb{Q}_p}}^{(3),\text{ad}})$, are isomorphisms.

7.2.3. Notations. Let s and $s_{\mathbb{R}}$ be the number of simple factors of $H_A^{(4),\text{ad}}$ and $H_{A,\mathbb{R}}^{(4),\text{ad}}$ (respectively). Thus s is the number of elements of the set J and $s_{\mathbb{R}} = [F : \mathbb{Q}]s$.

Let $\mathcal{S}$ be the set of extremal vertices of the Dynkin diagram of either $\text{Lie}(H_{A,\overline{\mathbb{Q}}}^{(4),\text{ad}})$ or $\text{Lie}(H_{A,\overline{\mathbb{Q}_p}}^{(4),\text{ad}})$. The group $\text{Gal}(\mathbb{Q})$ acts on $\mathcal{S}$ (see Subsection 2.1). Until Subsubsection 7.2.6 we will assume that $\mathfrak{L} \neq A_1$; thus $\mathfrak{L}_4 \neq A_1$. As $\mathfrak{L}_4 \neq A_1$, we can identify $\mathcal{S}$ with the $\text{Gal}(\mathbb{Q})$ -set $\text{Hom}_{\mathbb{Q}}(K_{\mathcal{S}}, \overline{\mathbb{Q}})$, for $K_{\mathcal{S}}$ a suitable product indexed by $j \in J$ of totally imaginary quadratic extensions of F . By replacing F with a totally real finite field extension of it, we can assume that there exists a totally imaginary quadratic extension K of F such that we have $K_{\mathcal{S}} = K^s$ (cf. Variant 4.9 (b)). The $\mathbb{Q}$ -algebra $K_{\mathcal{S}}$ acts faithfully on W_A (see the proof of Proposition 4.1). Thus we can identify $\text{Res}_{K_{\mathcal{S}}/\mathbb{Q}} \mathbb{G}_{m,K_{\mathcal{S}}}$ with a torus of $\mathbf{GL}_{W_A}$. All monomorphisms $\mathbb{S} \hookrightarrow H_{A,\mathbb{R}}^{(4)}$ that are elements of $X_A^{(4)}$, factor through the extension

to $\mathbb{R}$ of the subgroup of $\mathbf{GL}_{W_A}$ generated by $H_A^{(4),\text{der}}$, by $Z(\mathbf{GL}_{W_A})$, and by the maximal subtorus T_c of $\text{Res}_{K_S/\mathbb{Q}}\mathbb{G}_{m,K_S}$ which over $\mathbb{R}$ is compact (cf. the proofs of Proposition 4.1 and Theorem 4.8 and the definition of Hodge quasi products).

7.2.4. Proposition. *We recall that $\mathfrak{L} \neq A_1$ and that $\mathcal{F}_1$ is a fixed simple factor of $G_{\overline{\mathbb{Q}_p}}^{\text{ad}}$.*

Let $W_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_p} = \bigoplus_{u \in \chi_A} \mathcal{V}_u$ be the minimal direct sum decomposition such that the torus $Z^0(H_A)_{\overline{\mathbb{Q}_p}}$ acts on $\mathcal{V}_u$ via the character u of it. Let $\mathcal{N}_0$ be the unique normal, semisimple subgroup of $H_{A,\overline{\mathbb{Q}_p}}^{(2),\text{der}}$ whose adjoint is the product of those simple factors of $H_{A,\overline{\mathbb{Q}_p}}^{\text{ad}} = H_{A,\overline{\mathbb{Q}_p}}^{(2),\text{ad}}$ onto which $\mathcal{F}_1$ surjects. Then there exists $u_0 \in \chi_A$ such that $\mathcal{N}_0$ acts faithfully on $\mathcal{V}_{u_0}$. If $\mathfrak{L} = A_n$, then all irreducible $\text{Lie}(\mathcal{F}_1)$ -submodules of $\mathcal{V}_{u_0}$ are isomorphic.

Proof: We recall $f_A^{(4)}$ is a PEL type embedding, cf. property 7.2.2 (iii) and Fact 2.4.1 (b). Thus the Frobenius torus T_v is naturally identified with a torus of a particular type of inner form of $H_A^{(4)}$, cf. the choice of v and Subsubsections 5.6.1 to 5.6.3 applied to (A, v) (instead of (B, v) of Subsection 5.5). We will use the torus T_v to show that the torus $Z^0(H_A)$ is small enough so that the Proposition holds; thus based on Fact 5.6.2, to ease notations we will assume that this inner form of $H_A^{(4)}$ is trivial and therefore that T_v is naturally a torus of $H_A^{(4)}$. As the arguments are long, we will group them in six parts.

Part I. The torus $T_v^{(4)}$. We check that the centralizer $T_v^{(4)}$ of T_v in $H_A^{(4)}$ is a maximal torus of $H_A^{(4)}$. It is well known that $T_v^{(4)}$ is a reductive subgroup of $H_A^{(4)}$ (cf. [Bor, Ch. IV, 13.17, Cor. 2 (a)]) of the same rank as $H_A^{(4)}$. Thus to check that $T_v^{(4)}$ is a torus, it suffices to check the following statement: the intersection of $T_{v,\overline{\mathbb{Q}_p}}^{(4)}$ with each factor of $H_{A,\overline{\mathbb{Q}_p}}^{(4),\text{der}}$ that has a simple adjoint, is a torus. All factors of $G_{\overline{\mathbb{Q}_p}}^{\text{ad}}$ are of $\mathfrak{L}$ type and an $H_A^{(4)}(\overline{\mathbb{Q}_p})$ -conjugate of $T_{v,\overline{\mathbb{Q}_p}}$ is a maximal torus of $G_{\overline{\mathbb{Q}_p}}$. Thus, if $\mathfrak{L} = A_n$, then $\mathfrak{L} = \mathfrak{L}_4$ and the statement can be easily checked over $\overline{\mathbb{Q}_p}$. If $\mathfrak{L} \neq A_n$, then based on properties 7.2.2 (i) and (ii) and on the property 7.2 (iii), the statement follows from Fact 4.3.2. Thus $T_v^{(4)}$ is a maximal torus of $H_A^{(4)}$.

Part II. The equivalence relation $\mathcal{R}$. Let $\mathcal{R}$ be the unique equivalence relation on $\mathcal{S}$ that has the following property. Two distinct elements $\mathfrak{n}_1$ and $\mathfrak{n}_2$ of $\mathcal{S}$ are in relation $\mathcal{R}$ if and only if the following two conditions hold:

(a) *they are vertices of Dynkin diagrams of the Lie algebras of two distinct simple factors $\mathcal{F}'_0$ and $\mathcal{F}''_0$ of $H_{A,\overline{\mathbb{Q}_p}}^{(4),\text{ad}}$ and there exists a simple factor $\mathcal{F}'_1$ of $G_{\overline{\mathbb{Q}_p}}^{\text{ad}}$ whose Lie algebra maps injectively into both $\text{Lie}(\mathcal{F}'_0)$ and $\text{Lie}(\mathcal{F}''_0)$;*

(b) *with respect to $\mathcal{F}'_1$ and to the image of $T_{v,\overline{\mathbb{Q}_p}}^{(4)}$ in $H_{A,\overline{\mathbb{Q}_p}}^{(4),\text{ad}}$, they are both either original vertices or ending vertices of Dynkin diagrams.*

We explain the meaning in (b) of “are both either original vertices or ending vertices”. As $T_{v,\overline{\mathbb{Q}_p}}^{(4)}$ is a maximal torus of $H_A^{(4)}$, there exist isomorphisms $\mathcal{F}'_0 \simeq \mathcal{F}''_0$ under which the images in $\mathcal{F}'_0$ and $\mathcal{F}''_0$ of an a priori fixed maximal torus of the semisimple subgroup of $G_{\overline{\mathbb{Q}_p}}$ whose adjoint is $\mathcal{F}'_1$, are canonically identified. Each such isomorphism is uniquely

determined up to an automorphism of $\mathcal{F}'_0$ that fixes a maximal torus of it and thus up to an inner automorphism of $\mathcal{F}'_0$. Thus $\mathcal{F}'_0$ and $\mathcal{F}''_0$ are naturally identified up to inner automorphisms and therefore it makes sense to speak in (b) about “are both either original vertices or ending vertices”. Based on this and the first part of the property 7.2.2 (v), properties (a) and (b) define uniquely an equivalence relation on $\mathcal{S}$. Each equivalence class of $\mathcal{R}$ contains at most one vertex of each Dynkin diagram of a simple factor of $\text{Lie}(H_{\mathbb{Q}_p}^{(4),\text{ad}})$, cf. the very definition of $\mathcal{R}$.

Part III. The Galois invariance of $\mathcal{R}$. In this Part III we check that the relation $\mathcal{R}$ is $\text{Gal}(\mathbb{Q})$ -invariant. In (a) and (b) (of Part II), the roles of $G_{\overline{\mathbb{Q}_p}}$ and $T_{v,\overline{\mathbb{Q}_p}}^{(4)}$ can be also played by an $H_A^{(4)}(\overline{\mathbb{Q}_p})$ -conjugate of $G_{\overline{\mathbb{Q}_p}}$ and by an arbitrary maximal torus of this $H_A^{(4)}(\overline{\mathbb{Q}_p})$ -conjugate. But there exists an $H_A^{(4)}(\overline{\mathbb{Q}_p})$ -conjugate of $G_{\overline{\mathbb{Q}_p}}$ that contains the extension to $\overline{\mathbb{Q}_p}$ of the subtorus of T_v of $H_A^{(4)}$. Thus not to introduce extra notations, in the next paragraph we can assume that the monomorphism $T_v \hookrightarrow H_A^{(4)}$ has the property that its extension to $\overline{\mathbb{Q}_p}$ factors as a composite monomorphism $T_{v,\overline{\mathbb{Q}_p}} \hookrightarrow G_{\overline{\mathbb{Q}_p}} \hookrightarrow H_{A,\overline{\mathbb{Q}_p}}^{(4)}$.

Let $\mathbf{n}'_1$ and $\mathbf{n}'_2$ (resp. $\mathbf{n}''_1$ and $\mathbf{n}''_2$) be the two vertices of $\mathcal{S}$ that correspond to $\mathcal{F}'_0$ (resp. to $\mathcal{F}''_0$). We can assume that $\mathbf{n}'_1$ and $\mathbf{n}''_1$ are original vertices and that $\mathbf{n}'_2$ and $\mathbf{n}''_2$ are ending vertices in the sense of Part II. The equivalence relation $\mathcal{R}$ on $\mathcal{S}$ is the smallest equivalence relation on $\mathcal{S}$ that contains all pairs of the form $(\mathbf{n}'_i, \mathbf{n}''_i)$ with $i \in \{1, 2\}$. Let $T_{v,1}$ be the unique subtorus of $T_{v,\overline{\mathbb{Q}_p}}$ with the property that $\text{Lie}(T_{v,1})$ is a Cartan Lie subalgebra of $\text{Lie}(\mathcal{F}'_1)$. Let $T_{v,1}^\perp$ be the unique subtorus of $T_{v,\overline{\mathbb{Q}_p}}$ with the property that the natural homomorphism $T_{v,1} \times_{\overline{\mathbb{Q}_p}} T_{v,1}^\perp \rightarrow T_{v,\overline{\mathbb{Q}_p}}$ is an isogeny. The torus $T_{v,1}^\perp$ has a trivial image in $\mathcal{F}'_1$ and thus the image of $\text{Lie}(T_{v,1}^\perp)$ in $\text{Lie}(\mathcal{F}'_0) \oplus \text{Lie}(\mathcal{F}''_0)$ is trivial. The Galois group $\text{Gal}(\mathbb{Q})$ acts naturally on the set of subtori of $T_{v,\overline{\mathbb{Q}_p}}$ and on the set of simple factors of $H_{A,\overline{\mathbb{Q}_p}}^{(4)}$. Let $\gamma \in \text{Gal}(\mathbb{Q})$. The torus $\gamma(T_{v,1})$ (resp. $\gamma(T_{v,1}^\perp)$) is a subtorus of $T_{v,\overline{\mathbb{Q}_p}}$ which has non-trivial images (resp. has trivial images) in each one of the two simple factors $\text{Lie}(\gamma(\mathcal{F}'_0))$ and $\text{Lie}(\gamma(\mathcal{F}''_0))$ of $\text{Lie}(H_{A,\overline{\mathbb{Q}_p}}^{(4),\text{ad}})$. Let $\mathcal{F}'_{1,1}$ and $\mathcal{F}'_{1,2}$ be the unique simple factors of $G_{\overline{\mathbb{Q}_p}}^{\text{ad}}$ whose Lie algebras map injectively into $\text{Lie}(\gamma(\mathcal{F}'_0))$ and $\text{Lie}(\gamma(\mathcal{F}''_0))$ (respectively), cf. property 7.2.2 (v). We show that the assumption that $\mathcal{F}'_{1,1} \neq \mathcal{F}'_{1,2}$ leads to a contradiction. The torus $\gamma(T_{v,1}^\perp)$ has a trivial image in $\mathcal{F}'_{1,1} \times_{\overline{\mathbb{Q}_p}} \mathcal{F}'_{1,2}$. Thus the rank of $G_{\overline{\mathbb{Q}_p}}$ (i.e., of either T_v or $T_{v,1} \times_{\overline{\mathbb{Q}_p}} T_{v,1}^\perp$) is at least equal to the rank of $T_{v,1}^\perp \times_{\overline{\mathbb{Q}_p}} \mathcal{F}'_{1,1} \times_{\overline{\mathbb{Q}_p}} \mathcal{F}'_{1,2}$. Therefore the rank of $T_{v,1}$ is at least equal to the rank of $\mathcal{F}'_{1,1} \times_{\overline{\mathbb{Q}_p}} \mathcal{F}'_{1,2}$. But as the groups $T_{v,1}$, $\mathcal{F}'_{1,1}$, and $\mathcal{F}'_{1,2}$ have the same ranks as $\mathfrak{L}$, we reached a contradiction. Thus $\mathcal{F}'_{1,1} = \mathcal{F}'_{1,2}$. Therefore we can define $\gamma(\mathcal{F}'_1) := \mathcal{F}'_{1,1} = \mathcal{F}'_{1,2}$; it is a simple factor of $G_{\overline{\mathbb{Q}_p}}^{\text{ad}}$ whose Lie algebra maps injectively into both $\text{Lie}(\gamma(\mathcal{F}'_0))$ and $\text{Lie}(\gamma(\mathcal{F}''_0))$. As the tori T_v and $T_v^{(4)}$ are defined over $\mathbb{Q}$ and as $T_{v,1}$ is defined over $\overline{\mathbb{Q}}$, it is easy to see that the vertices $\gamma(\mathbf{n}'_1)$ and $\gamma(\mathbf{n}''_1)$ are both original vertices and that the vertices $\gamma(\mathbf{n}'_2)$ and $\gamma(\mathbf{n}''_2)$ are both ending vertices of $\mathcal{S}$. Thus we have $(\gamma(\mathbf{n}'_1), \gamma(\mathbf{n}''_1)), (\gamma(\mathbf{n}'_2), \gamma(\mathbf{n}''_2)) \in \mathcal{R}$. This implies that the equivalence relation $\mathcal{R}$ is $\text{Gal}(\mathbb{Q})$ -invariant.

The Galois group $\text{Gal}(\mathbb{Q})$ acts on the quotient set $\mathcal{S}_1 := \mathcal{S}/\mathcal{R}$. Let $K_{\mathcal{S}_1}$ be the étale $\mathbb{Q}$ -algebra such that we can identify $\mathcal{S}_1$ with the $\text{Gal}(\mathbb{Q})$ -set $\text{Hom}_{\mathbb{Q}}(K_{\mathcal{S}_1}, \overline{\mathbb{Q}})$. Let $T_{c,1}$ be the maximal subtorus of $\text{Res}_{K_{\mathcal{S}_1}/\mathbb{Q}} \mathbb{G}_{m, K_{\mathcal{S}_1}}$ which over $\mathbb{R}$ is compact. We identify $K_{\mathcal{S}_1}$ with a $\mathbb{Q}$ -subalgebra of $K_{\mathcal{S}}$ and thus $K_{\mathcal{S}_1}$ is a product of at most s fields. We have $T_{c,1} \leq T_c$. Let $H_A^{(5)}$ be the reductive subgroup of $H_A^{(4)}$ generated by $H_A^{(4),\text{der}}$, $Z(\mathbf{GL}_{W_A})$, and $T_{c,1}$.

Part IV. Cocharacters. We write $H_{A,\mathbb{R}}^{(4),\text{der}} = \prod_{i=1}^{s_{\mathbb{R}}} F_i^{(4)}$ as a product of normal, semisimple subgroups that have absolutely simple adjoints. Let $W_A \otimes_{\mathbb{Q}} \mathbb{R} = \bigoplus_{i=1}^{s_{\mathbb{R}}} V_i^{(4)}$ be the direct sum decomposition into $H_{A,\mathbb{R}}^{(4)}$ -modules such that for all $i \in \{1, \dots, s_{\mathbb{R}}\}$ the group $F_i^{(4)}$ acts trivially on $V_{i_1}^{(4)}$ if $i_1 \in \{1, \dots, s_{\mathbb{R}}\} \setminus \{i\}$ and acts non-trivially on each irreducible $H_{A,\mathbb{R}}^{(4)}$ -submodule of $V_i^{(4)}$. Let $T_i^{(4)}$ be the direct factor of $T_{c,\mathbb{R}}$ that is isomorphic to $\mathbf{SO}(2)_{\mathbb{R}} = \mathbb{S}/\mathbb{G}_{m,\mathbb{R}}$ and that acts trivially on $\bigoplus_{i_1 \in \{1, \dots, s_{\mathbb{R}}\} \setminus \{i\}} V_{i_1}^{(4)}$. We identify $F_i^{(4)}$ and $T_i^{(4)}$ with their images in $\mathbf{GL}_{V_i^{(4)}}$. Let $x_4 \in X_A^{(2)} \subseteq X_A^{(4)}$. Let $J_i^{(4)}$ be the image of the restriction of $x_4 : \mathbb{S} \rightarrow H_{A,\mathbb{R}}^{(4)}$ to the compact subtorus $\mathbf{SO}(2)_{\mathbb{R}}$ of $\mathbb{S}$ in $T_i^{(4)}/(F_i^{(4)} \cap T_i^{(4)})$; it is equipped with a homomorphism $j_i : \mathbf{SO}(2)_{\mathbb{R}} \rightarrow J_i^{(4)}$. Let $\mu_A^{(3),\text{ad}} : \mathbb{G}_{m,\mathbb{C}} \rightarrow H_{A,\mathbb{C}}^{(3),\text{ad}}$ and $\mu_A^{(4),\text{ad}} : \mathbb{G}_{m,\mathbb{C}} \rightarrow H_{A,\mathbb{C}}^{(4),\text{ad}}$ be the cocharacters defined naturally by $\mu_A : \mathbb{G}_{m,\mathbb{C}} \rightarrow H_{A,\mathbb{C}}$.

For $q \in \{l, p\}$ we consider an embedding $i_{\overline{\mathbb{Q}}_q} : \overline{\mathbb{Q}}_q \hookrightarrow \mathbb{C}$ that extends the embedding i_E of Subsection 1.1. We use $i_{\overline{\mathbb{Q}}_p}$ to identify the sets of simple factors of $H_{A,\mathbb{C}}^{(4),\text{ad}}$ and $H_{A,\overline{\mathbb{Q}}_p}^{(4),\text{ad}}$. By applying the property 3.4.1 (b), we get the existence of a cocharacter $\mu_{v,\overline{\mathbb{Q}}_l}^{\text{ét}} : \mathbb{G}_{m,\overline{\mathbb{Q}}_l} \rightarrow G_{\overline{\mathbb{Q}}_l}$ whose extension via $i_{\overline{\mathbb{Q}}_l}$ is $H_A^{(4)}(\mathbb{C})$ -conjugate to μ_A . Based on Subsubsection 5.6.4, we can assume that $\mu_{v,\overline{\mathbb{Q}}_l}^{\text{ét}}$ factors through an $H_A^{(4)}(\overline{\mathbb{Q}}_l)$ -conjugate of $T_{v,\overline{\mathbb{Q}}_l}$. As an $H_A^{(4)}(\overline{\mathbb{Q}}_p)$ -conjugate of $T_{v,\overline{\mathbb{Q}}_p}$ is a maximal torus of $G_{\overline{\mathbb{Q}}_p}$, we conclude that there exists a cocharacter of $G_{\overline{\mathbb{Q}}_p}$ whose extension via $i_{\overline{\mathbb{Q}}_p}$ is $H_A^{(4)}(\mathbb{C})$ -conjugate to μ_A . From this and the second part of the property 7.2.2 (v) we get that:

(c.i) *two simple factors of the group pair $(H_{A,\mathbb{C}}^{(3),\text{ad}}, [\mu_A^{(3),\text{ad}}])$ that correspond to two simple factors of $\text{Lie}(H_{A,\overline{\mathbb{Q}}_p}^{(3),\text{ad}})$ into which the same simple factor of $\text{Lie}(G_{\overline{\mathbb{Q}}_p}^{\text{ad}})$ injects, are isomorphic.*

From the property (c.i) and the description of the monomorphisms $H_{j,\overline{\mathbb{Q}}}^{(3),\text{der}} \hookrightarrow H_{j,\overline{\mathbb{Q}}}^{(4),\text{der}}$ in the property 7.2.2 (ii), we get that:

(c.ii) *two simple factors of the group pair $(H_{A,\mathbb{C}}^{(4),\text{ad}}, [\mu_A^{(4),\text{ad}}])$ that correspond to two simple factors of $\text{Lie}(H_{A,\overline{\mathbb{Q}}_p}^{(4),\text{ad}})$ into which the same simple factor of $\text{Lie}(G_{\overline{\mathbb{Q}}_p}^{\text{ad}})$ injects, are isomorphic.*

In properties (c.i) and (c.ii) we can replace $(H_{A,\mathbb{C}}^{(\star),\text{ad}}, [\mu_A^{(\star),\text{ad}}])$ by $(H_{A,\mathbb{R}}^{(\star),\text{ad}}, [\mu_A^{(\star),\text{ad}}])$ (cf. [De2, Prop. 1.2.2]), where $\star \in \{3, 4\}$.

Part V. Downsizing the torus $Z^0(H_A)$. We show that we can modify x_4 (and thus implicitly $X_A^{(2)}$ and $X_A^{(4)}$) such that its images in $X_A^{\text{ad}} = X_A^{(2),\text{ad}}$ and $X_A^{(4),\text{ad}}$ are the same and the composite homomorphism

$$(j_1, \dots, j_{s_{\mathbb{R}}}) : \mathbf{SO}(2)_{\mathbb{R}} \rightarrow \prod_{i=1}^{s_{\mathbb{R}}} J_i^{(4)} \hookrightarrow \prod_{i=1}^{s_{\mathbb{R}}} T_i^{(4)} / (F_i^{(4)} \cap T_i^{(4)}) = T_{c,\mathbb{R}} / (T_{c,\mathbb{R}}[2])$$

factors through $T_{c,1,\mathbb{R}} / (T_{c,1,\mathbb{R}}[2])$. Such a modification of x_4 is allowed (cf. Proposition 5.4.1 and the constructions of Subsection 5.2) and it is defined naturally by a replacement of the $s_{\mathbb{R}}$ -tuple $(j_1, \dots, j_{s_{\mathbb{R}}})$ with another $s_{\mathbb{R}}$ -tuple $(j_1^{\varepsilon_1}, \dots, j_{s_{\mathbb{R}}}^{\varepsilon_{s_{\mathbb{R}}}})$, where $\varepsilon_1, \dots, \varepsilon_{s_{\mathbb{R}}} \in \{-1, 1\}$ and where we can have $\varepsilon_i = -1$ only if $F_i^{(4)}$ is compact (cf. the paragraph after properties (a) and (b) of the proof of Proposition 4.1). The homomorphism $(j_1, \dots, j_{s_{\mathbb{R}}})$ factors through $T_{c,1,\mathbb{R}} / (T_{c,1,\mathbb{R}}[2])$ if and only if the following property holds:

(d) for arbitrary distinct elements $i', i'' \in \{1, \dots, s_{\mathbb{R}}\}$ that correspond to factors $\mathcal{F}'_0$ and $\mathcal{F}''_0$ as in (a) of Part II, the representations of $\mathbf{SO}(2)_{\mathbb{R}}$ on $V_{i'}^{(4)}$ and $V_{i''}^{(4)}$ defined by $j_{i'}$ and $j_{i''}$ (respectively) are such that we have identifications

$$V_{i'}^{(4)} \otimes_{\mathbb{R}} \mathbb{C} = V_{i'}^{(4)+} \oplus V_{i'}^{(4)-} = V_{i''}^{(4)+} \oplus V_{i''}^{(4)-} = V_{i''}^{(4)} \otimes_{\mathbb{R}} \mathbb{C}$$

of $\mathbf{SO}(2)_{\mathbb{C}} = \mathbb{G}_{m,\mathbb{C}}$ -modules, that are defined naturally by (2) and that have the property that the representations of $F_{i',\mathbb{C}}^{(4)}$ and $F_{i'',\mathbb{C}}^{(4)}$ on $V_{i'}^{(4)+}$ and $V_{i''}^{(4)+}$ (respectively) are both via original vertices of the Dynkin diagrams of the Lie algebras of $F_{i',\mathbb{C}}^{(4)}$ and $F_{i'',\mathbb{C}}^{(4)}$ (here original vertices are in the sense of (b) of Part II, the role of $\overline{\mathbb{Q}_p}$ being replaced by the one of $\mathbb{C}$).

Based on the property (c.ii) the groups $F_{i'}^{(4)}$ and $F_{i''}^{(4)}$ are either both compact or both non-compact. It is easy to see that we can choose the ε_i 's such that (d) holds for all pairs (i', i'') such that both groups $F_{i'}^{(4)}$ and $F_{i''}^{(4)}$ are compact. If $F_{i'}^{(4)}$ and $F_{i''}^{(4)}$ are both non-compact, then from the property (c.ii) and the uniqueness of $\varepsilon_{i'}$ and $\varepsilon_{i''}$ (they are both 1) we get that (d) automatically holds for the pair (i', i'') . Therefore we can assume that (d) holds and thus that the homomorphism $(j_1, \dots, j_{s_{\mathbb{R}}})$ factors through $T_{c,1,\mathbb{R}} / (T_{c,1,\mathbb{R}}[2])$. We get that μ_A factors through the reductive group $H_{A,\mathbb{C}}^{(5)}$ defined at the end of Part III. As H_A is a Mumford–Tate group, we get that $Z^0(H_A) \leq Z^0(H_A^{(5)})$.

Part VI. End of the proof of 7.2.4. Let $W_A \otimes_{\mathbb{Q}} \overline{\mathbb{Q}_p} = \bigoplus_{t \in \chi_A^{(5)}} \mathcal{V}_t$ be the minimal direct sum decomposition such that the torus $Z^0(H_{A,\overline{\mathbb{Q}_p}}^{(5)})$ acts on each $\mathcal{V}_t$ via its character t . See proof of Proposition 4.1 for how the torus $\text{Res}_{K_S/\mathbb{Q}} \mathbb{G}_{m,K_S}$ (and thus also how its subtorus $T_{c,1}$) acts on W_A . The direct sum decompositions $V_i^{(4)} \otimes_{\mathbb{R}} \mathbb{C} = V_i^{(4)+} \oplus V_i^{(4)-}$ (see (d)) are such that $\text{Res}_{K_S/\mathbb{Q}} \mathbb{G}_{m,K_S,\mathbb{C}}$ acts on each $V_i^{(4),u}$ via a unique character, where $i \in \{1, \dots, s_{\mathbb{R}}\}$ and $u \in \{-, +\}$ (see (2) of the proof of Proposition 4.1). The upper indices $+$ and $-$ are such that $T_{c,1,\mathbb{C}}$ acts on $V_{i'}^{(4),u'}$ and $V_{i''}^{(4),u''}$ via the same character if and only if $u' = u'' \in \{-, +\}$ and the elements $i', i'' \in \{1, \dots, s_{\mathbb{R}}\}$ are as in (d). We get the existence of two distinct characters $t_1, t_2 \in \chi_A^{(5)}$ such that $\mathcal{N}_0$ acts faithfully on both $\mathcal{V}_{t_1}$

and $\mathcal{V}_{t_2}$ and acts trivially on $\mathcal{V}_t$ if $t \in \chi_A^{(5)} \setminus \{t_1, t_2\}$. As $Z^0(H_A)$ is a subtorus of $Z^0(H_A^{(5)})$, each $\overline{\mathbb{Q}_p}$ -vector space $\mathcal{V}_u$ (of Proposition 7.2.4) is a direct sum of some of the $\mathcal{V}_t$'s. Thus we can take the character $u_0 \in \chi_A$ such that we have either $\mathcal{V}_{t_1} \subseteq \mathcal{V}_{u_0}$ or $\mathcal{V}_{t_2} \subseteq \mathcal{V}_{u_0}$.

We assume that $\mathfrak{L} = A_n$. Due to the vertices parts of (b) and (d) and to the identity $\mathfrak{L} = \mathfrak{L}_4$, the irreducible $\text{Lie}(\mathcal{F}_1)$ -submodules of $\mathcal{V}_{u_0}$ are all isomorphic. $\square$

7.2.5. End of the proof for $\mathfrak{L} \neq A_1$. From properties 7.2.2 (iii) and (iv) we get that the representation of $\mathcal{N}_0$ on $\mathcal{V}_{u_0}$ is a direct sum of irreducible representations of its factors that have absolutely simple adjoints (to be compared with the property 5.7 (a)). Let $\text{Lie}(\mathcal{F}_{1,0})$ be a simple factor of $\text{Lie}(\mathcal{N}_0)$. If $\mathfrak{L}$ is B_n or C_n with $n \geq 2$, then from Proposition 5.10 (a) we get that the simple $\text{Lie}(\mathcal{F}_{1,0})$ -submodules of $\mathcal{V}_{u_0}$ are all isomorphic. If $\mathfrak{L} = D_4$, then each (H_j, X_j) is of $D_4^{\mathbb{R}}$ type and thus from Proposition 5.10 (b) we get that both half spin representations of $\text{Lie}(\mathcal{F}_{1,0})$ are among the simple $\text{Lie}(\mathcal{N}_0)$ -submodules of $\mathcal{V}_{u_0}$. Let now $\mathfrak{L}$ be D_n , with $n \geq 5$. If $\mathcal{N}_0$ has a normal factor that is a $\mathbf{Spin}_{2n, \overline{\mathbb{Q}_p}}$ group, then from the property 7.2.4 (c.ii) we get that each simple, adjoint Shimura pair (H_j, X_j) such that the intersection $\mathcal{N}_0^{\text{ad}} \cap H_{j, \overline{\mathbb{Q}_p}}$ (taken inside $H_{A, \overline{\mathbb{Q}_p}}^{\text{ad}}$) is non-trivial, is of $D_n^{\mathbb{R}}$ type. Thus $\mathcal{N}_0$ is a product of $\mathbf{Spin}_{2n, \overline{\mathbb{Q}_p}}$ groups (cf. property (iv) of Variant 4.9 (a)) and therefore again from Proposition 5.10 (b) we get that both half spin representations of $\text{Lie}(\mathcal{F}_{1,0})$ are among the simple $\text{Lie}(\mathcal{N}_0)$ -submodules of $\mathcal{V}_{u_0}$. If $\mathcal{N}_0$ has a normal factor that is an $\mathbf{SO}_{2n, \overline{\mathbb{Q}_p}}$ group, then a similar argument shows that each simple, adjoint Shimura pair (H_j, X_j) such that the intersection $\mathcal{N}_0^{\text{ad}} \cap H_{j, \overline{\mathbb{Q}_p}}$ (taken inside $H_{A, \overline{\mathbb{Q}_p}}^{\text{ad}}$) is non-trivial, is of $D_n^{\mathbb{H}}$ type, and that $\mathcal{N}_0$ is a product of $\mathbf{SO}_{2n, \overline{\mathbb{Q}_p}}$ groups. Thus the simple $\text{Lie}(\mathcal{F}_{1,0})$ -submodules of $\mathcal{V}_{u_0}$ are all isomorphic to the standard $2n$ dimensional simple $\text{Lie}(\mathcal{F}_1)$ -module.

Regardless of what $\mathfrak{L} \neq A_1$ is, based on the previous paragraph we conclude that the $\text{Lie}(\mathcal{F}_1)$ -module $\mathcal{V}_{u_0}$ is either isotypic or a direct sum of two non-isomorphic, isotypic $\text{Lie}(\mathcal{F}_1)$ -modules. The last situation happens if and only if $\mathcal{N}_0$ is a product of $\mathbf{Spin}_{2n, \overline{\mathbb{Q}_p}}$ groups.

Let s_0 be the number of simple factors of $\mathcal{N}_0^{\text{ad}}$. The Lie monomorphism $\text{Lie}(\mathcal{F}_1) \hookrightarrow \text{Lie}(\mathcal{N}_0^{\text{ad}})$ is isomorphic to the diagonal embedding $\text{Lie}(\mathcal{F}_{1,0}) \hookrightarrow \text{Lie}(\mathcal{F}_{1,0})^{s_0}$. If $\mathcal{N}_0$ is not (resp. is) a product of $\mathbf{Spin}_{2n, \overline{\mathbb{Q}_p}}$ groups, then the $\text{Lie}(\mathcal{N}_0)$ -module $\mathcal{V}_{u_0}$ is a direct sum of $2s_0$ (resp. $4s_0$) non-isomorphic, isotypic $\text{Lie}(\mathcal{N}_0)$ -modules; thus the $H_{A, \overline{\mathbb{Q}_p}}$ -module $\mathcal{V}_{u_0}$ is also a direct sum of $2s_0$ (resp. $4s_0$) non-isomorphic, isotypic $H_{A, \overline{\mathbb{Q}_p}}$ -modules. But the $G_{\overline{\mathbb{Q}_p}}$ -module $\mathcal{V}_{u_0}$ is a direct sum of 2 (resp. 4) non-isomorphic, isotypic $G_{\overline{\mathbb{Q}_p}}$ -modules, cf. the very definition of $\mathcal{N}_0$ in Proposition 7.2.4. Based on this and Theorem 1.1.4, we get that $s_0 = 1$. Thus the simple factor $\mathcal{F}_1$ of $G_{\overline{\mathbb{Q}_p}}^{\text{ad}}$ does not surject onto two or more simple factors of $H_{A, \overline{\mathbb{Q}_p}}^{\text{ad}}$. This proves the Theorem 7.2 for $\mathfrak{L} \neq A_1$.

7.2.6. End of the proof for $\mathfrak{L} = A_1$. If $\mathfrak{L} = A_1$, then the only modifications required to be made in Subsubsections 7.2.3 to 7.2.5 are as follows. For $j \in J$ we have $H_j^{(3)} = H_j^{(4)}$ and we have to replace the factor of $K_{\mathfrak{s}}$ that corresponds to j by a totally imaginary quadratic extension K_j of F obtained as in the case $n = 1$ of the proof of Proposition 4.1; by replacing F with a totally real finite field extension of it (cf. Variant 4.9 (b)), we can assume that there exists a totally imaginary quadratic extension K_0 of $\mathbb{Q}$ such that we have

$K_j = K := F \otimes_{\mathbb{Q}} K_0$ for all $j \in J$. We can identify $\mathcal{S}$ with the $\text{Gal}(\mathbb{Q})$ -set $\text{Hom}_{\mathbb{Q}}(F^s, \overline{\mathbb{Q}})$. We define $K_{\mathcal{S}} := K^s$. The equivalence class $\mathcal{R}$ on $\mathcal{S}$ is defined this time by the property 7.2.4 (a) alone. If $\mathcal{S}_1 := \mathcal{S}/\mathcal{R}$, let $F_{\mathcal{S}_1}$ be the $\mathbb{Q}$ -subalgebra of F^s such that we can identify $\mathcal{S}_1$ with the $\text{Gal}(\mathbb{Q})$ -set $\text{Hom}_{\mathbb{Q}}(F_{\mathcal{S}_1}, \overline{\mathbb{Q}})$. Let $K_{\mathcal{S}_1} := F_{\mathcal{S}_1} \otimes_{\mathbb{Q}} K_0$. Let $T_{c,1}$ be as in the proof of Proposition 7.2.4. The rest of the arguments do not have to be modified. This ends the proof of Theorem 7.2 for $\mathfrak{L} = A_1$ and thus also for all $\mathfrak{L}$. $\square$

7.3. Proofs of 1.3.4 and 1.3.6. If for all elements $j \in J$ the type of (H_j, X_j) is among those listed in Theorem 1.3.4, then the property 7.2 (i) obviously holds and Theorem 7.1 implies that the property 7.2 (ii) also holds. Thus $G_{\mathbb{Q}_p}^{\text{ad}} = H_{A, \mathbb{Q}_p}^{\text{ad}}$, cf. Theorem 7.2. Therefore $G_{\mathbb{Q}_p} = H_{A, \mathbb{Q}_p}$, cf. Theorem 1.3.1. This ends the proof of Theorem 1.3.4.

To prove Theorem 1.3.6, we can assume that the condition 7.2 (ii) holds (cf. Proposition 5.4.1 and Subsubsection 5.4.3). The hypothesis of Theorem 1.3.6 implies that the conditions 7.2 (i) also holds. Thus Theorem 1.3.6 follows from Theorem 7.2. $\square$

7.4. Corollary. *Let $J_4 := \{j \in J \mid (H_j, X_j) \text{ is of either } D_4^{\mathbb{H}} \text{ or } D_4^{\mathbb{R}} \text{ type}\}$. If $j \in J_4$ (resp. if $j \in J \setminus J_4$) we assume that (H_j, X_j) is of non-inner $D_4^{\mathbb{H}}$ type (resp. is of one of the types listed in Theorem 1.3.4). We also assume that there exists a prime $q \in \mathbb{N}$ such that the following two properties hold:*

- (i) *the simple factors of $\prod_{j \in J_4} H_{j, \mathbb{Q}_q}$ are absolutely simple and pairwise non-isomorphic;*
- (ii) *for all elements $j \in J_4$ the Shimura pair (H_j, X_j) is with $\mathbb{Q}_q$ -involution.*

Then Conjecture 1.1.1 holds for A .

Proof: The proof of the Corollary is very much the same as the (d') part of the proof of Theorem 7.1 and the proof of Theorem 7.2. We can assume that the set J_4 is non-empty, cf. Theorem 1.3.4. Based on Theorem 1.1.6 we can assume that $p = q$. As in the (d') part of the proof of Theorem 7.1, based on (ii) and the absolutely simple part of (i) we get that for each element $j \in J_4$ the group $G_{\mathbb{Q}_p}$ surjects onto a simple factor of $H_{j, \mathbb{Q}_p}$.

Similar to the arguments that checked the property 7.2 (iii), based on Theorem 1.1.4 and Lemma 1.1.7 we argue that $G_{\overline{\mathbb{Q}_p}}$ surjects onto each simple factor of $\prod_{j \in J_4} H_{j, \overline{\mathbb{Q}_p}}$. Thus $G_{\mathbb{Q}_p}$ surjects onto $\prod_{j \in J_4} H_{j, \mathbb{Q}_p}$, cf. the pairwise non-isomorphic part of (i). Therefore in the proof of Theorem 7.2 we can work only with a Lie type $\mathfrak{L} \neq D_4$ and thus Theorem 7.2 applies (i.e., we have $G_{\mathbb{Q}_p}^{\text{ad}} = H_{A, \mathbb{Q}_p}^{\text{ad}}$). Therefore $G_{\mathbb{Q}_p} = H_{A, \mathbb{Q}_p}$, cf. Theorem 1.3.1. $\square$

7.5. The simplest cases not settled by 1.3.4. For the sake of future references we list below the “simplest” cases not settled by Theorem 1.3.4 (they involve adjoint groups whose ranks are small and of fixed parity). We assume that H_A^{ad} is a simple group over $\mathbb{Q}$ but we do not impose any restriction on either p or $G_{\mathbb{Q}_p}$. The image $\mathfrak{I}$ of $G_{\overline{\mathbb{Q}_p}}$ in a fixed simple factor of $H_{A, \mathbb{Q}_p}^{\text{ad}}$ is an adjoint group, cf. property 3.1 (c); let $\mathfrak{L}$ be its Lie type. In A5 to A9 below, the other possible values for $\mathfrak{L}$ allowed by [Pi, Table 4.2] are excluded as in Example 4 of Subsubsection 6.2.4. If $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of A_n type, then let $(H_0, X_0) := (H_A^{\text{ad}}, X_A^{\text{ad}})$ and let the set $\mathfrak{C}$ be as in Subsection 6.2.

A5 and A7. The Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of A_5 (resp. A_7) type and $\mathfrak{C} = \{1, \frac{1}{2}\}$ (resp. $\mathfrak{C}$ is $\{1\}$ or $\{1, \frac{1}{3}\}$). We could theoretically have $\mathfrak{L} = A_1 + A_2$ (resp. $\mathfrak{L} = A_1 + A_3$).

A8. The Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of A_8 type and $\mathfrak{C} = \{\frac{1}{2}\}$. We could theoretically have $\mathfrak{L} = A_2 + A_2$.

A9. The Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of A_9 type. We could theoretically have either $\mathfrak{L} = A_4$ and $\mathfrak{C} = \{\frac{2}{3}\}$ or $\mathfrak{L} = A_1 + A_4$ and $\mathfrak{C} \in \{\{1, \frac{1}{4}\}, \{1, \frac{2}{3}\}\}$.

A11. The Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of A_{11} type. We could theoretically have $\mathfrak{L} \in \{A_1 + A_5, A_1 + A_1 + A_2, A_2 + A_3, A_2 + C_2\}$. The Lie types $A_1 + A_1 + A_2$ and $A_2 + C_2$ have the same rank; if $\mathfrak{L}$ is one of these two types, then $\mathfrak{C} = \{1, \frac{1}{2}\}$.

C4 and C6. The Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of C_4 (resp. C_6) type. We could theoretically have $\mathfrak{L} = A_1 + A_1 + A_1$ (resp. $\mathfrak{L} = A_1 + D_3$).

D4 and D6. The Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of $D_4^{\mathbb{H}}$ type and Corollary 7.4 does not apply (resp. is of inner $D_6^{\mathbb{H}}$ type). We could theoretically have $\mathfrak{L} = B_3$ or $\mathfrak{L} = B_1 + B_2$ (resp. $\mathfrak{L} = C_1 + C_3$).

D35. The Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of $D_{35}^{\mathbb{H}}$ type. We could theoretically have $\mathfrak{L} = A_7$.

The A_n types with $n \in \{1, 2, 3, 4, 6, 10, 12\}$ are settled by Example 6 of Subsubsection 6.2.4.

7.5.1. Remarks. (a) The numerical tests of Subsection 6.2 “work” regardless of what q is. However, due to the fact that in Definition 6.1 (c) we are allowed to choose the prime q , there exist large classes of simple, adjoint Shimura pairs of non-special A_n type for which the numerical tests of Subsection 6.2 fail. We include one example related to A8 of Subsection 7.5. We assume that the group H_A^{ad} is simple and $H_{A, \mathbb{Q}_p}^{\text{ad}}$ has a simple factor $\mathcal{C}$ that is the adjoint group of the group of invertible elements of a central division algebra over $\mathbb{Q}_p$ of dimension 81. If $\mathfrak{I}$ is a subgroup of $\mathcal{C}_{\overline{\mathbb{Q}_p}}$, then we have $\mathfrak{L} \neq A_2 + A_2$ as otherwise $\mathfrak{I}$ (and therefore also $\mathcal{C}$) splits over a finite field extension of $\mathbb{Q}_p$ of degree a divisor of 12 and thus not divisible by 9. Thus $\mathfrak{L} = A_8$ and therefore $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of non-special A_8 type. Similar such examples can be constructed in connection to A7 of Subsection 7.5. This remark and Subsections 6.2 and 7.5 motivate the terminology “non-special” used in Definition 6.1 (c).

(b) Let (H_0, X_0) be a simple, adjoint, unitary Shimura pair. If by chance $F(H_0, X_0) \otimes_{\mathbb{Q}} \mathbb{Q}_q$ is an unramified field extension of $\mathbb{Q}_q$, then one can re-obtain easily most of the numerical properties of Subsection 6.2 without being “bothered” to mention T_1 in Definition 6.1 (b) or to appeal to Proposition 5.10. But this does not suffice to get them all (like Example 4 of Subsubsection 6.2.4) or to get Theorem 1.3.4 (a). Thus even if we have a good understanding of the (assumed to be very simple) arithmetics of the field $F(H_0, X_0)$, Theorem 7.2 (and thus implicitly Proposition 5.10) looks to us unavoidable i.e., the proof of Theorem 7.2 can not be simplified. However, in practice it is often easy to check that the numbers t_j of Lemma 6.2.2 do not depend on $j \in J_1$, and accordingly, that the pairs $(r_{j,m}, s_{j,m})$ (counted with multiplicities) and defined when $f(j)(m) = 1$, do not depend on $j \in J_1$. As in A9 of Subsection 7.5, we might not have a unique possibility for some of t_j ’s.

In what follows the abelian motives over a number subfield of $\mathbb{C}$ are defined using absolute Hodge cycles, see [DM, Sect. 6]. Using realizations in the Betti and étale co-

homologies, as for abelian varieties we speak about the Mumford–Tate groups and the attached (adjoint) Shimura pairs of their extensions to $\mathbb{C}$ and about the Mumford–Tate conjecture for them. Theorem 1.3.7 is a particular case of the following more general result.

7.6. Theorem (the independence property for abelian motives). *We recall that in Subsection 1.1 we fixed an embedding $i_E : E \hookrightarrow \mathbb{C}$. Let $(\mathcal{D}_i)_{i \in I}$ be a finite set of abelian motives over the number field E . We assume that for each $i \in I$, the Mumford–Tate conjecture is true for $\mathcal{D}_i$ and no simple factor of the adjoint Shimura pair attached to the extension of $\mathcal{D}_i$ to $\mathbb{C}$ is of $D_4^{\mathbb{H}}$ type. Then the Mumford–Tate conjecture is true for $\mathcal{D} := \bigoplus_{i \in I} \mathcal{D}_i$.*

Proof: We take the abelian variety A over E such that $\mathcal{D}$ is an object of the Tannakian category of abelian motives over E generated by the H_1 -motive $W_A = L_A \otimes_{\mathbb{Z}} \mathbb{Q}$ of A . By passing to a finite field extension of E , we can choose A such that it has a principal polarization λ_A (cf. [Mu2, Ch. IV, Sect. 23, Cor. 1]). Let $h_A : \mathbb{S} \rightarrow H_{A, \mathbb{R}}$ be as in Subsection 1.3. Let $f_A : (H_A, X_A) \hookrightarrow (\mathbf{GSp}(W_A, \psi_A), S_A)$ be as in Subsection 1.5. Let $H_{\mathcal{D}}$ be the reductive group over $\mathbb{Q}$ that is the image of H_A in $\mathbf{GL}_{W_{\mathcal{D}}}$, where $W_{\mathcal{D}}$ is the rational vector space that is the Betti realization of $\mathcal{D}$. We have to show that the identity component $G_{\mathcal{D}, \mathbb{Q}_p}$ of the algebraic envelope of the image of the p -adic Galois representation attached to $\mathcal{D}$ is $H_{\mathcal{D}, \mathbb{Q}_p}$. Always $G_{\mathcal{D}, \mathbb{Q}_p}$ is a subgroup of $H_{\mathcal{D}, \mathbb{Q}_p}$, cf. Theorem 1.1.2. Based on this and Theorem 1.3.1, we only have to show that $H_{\mathcal{D}, \mathbb{Q}_p}^{\text{ad}} = G_{\mathcal{D}, \mathbb{Q}_p}^{\text{ad}}$.

Let H_B^{der} and H_C^{der} be the commuting normal subgroups of H_A^{der} such that their adjoints are naturally identified with $H_{\mathcal{D}}^{\text{ad}}$ and $H_A^{\text{ad}}/H_{\mathcal{D}}^{\text{ad}}$ (respectively). Let H'_B be a reductive subgroup of H_A of the same rank as H_A such that (cf. [Ha, Lem. 5.5.3]) the $\mathbb{R}$ -ranks of $H_B^{\prime, \text{ab}}$ and H_A^{ab} are both 1 and we have $H_B^{\prime, \text{der}} = H_B^{\text{der}}$. Two maximal compact tori of H_C^{der} are $H_C^{\text{der}}(\mathbb{R})$ -conjugate. This implies that there exists a homomorphism $h_B : \mathbb{S} \rightarrow H'_{B, \mathbb{R}}$ such that the following two properties hold: (i) its composite with the inclusion $i_A^B : H'_{B, \mathbb{R}} \hookrightarrow H_{A, \mathbb{R}}$ is $H_A(\mathbb{R})$ -conjugate to h_A , and (ii) the homomorphisms $\mathbb{S} \rightarrow H_{B, \mathbb{R}}^{\text{ad}} = H_{\mathcal{D}, \mathbb{R}}^{\text{ad}}$ defined naturally by h_B and h_A coincide. We choose h_B such that the smallest subgroup H_B of H'_B with the property that h_B factors through $H_{B, \mathbb{R}}$, is a reductive group that has H_B^{der} as its derived group. By replacing E with a finite field extension of it, we can choose h_B such that moreover the polarized abelian variety over $\mathbb{C}$ defined (see Riemann's theorem of [De1, Thm. 4.7]) by the triple $(L_A, \varepsilon \psi_A, h_B)$ is the pull back of a polarized abelian variety (B, λ_B) over E ; here $\varepsilon \in \{-1, 1\}$ depends on the connected component of S_A to which $f_{A, \mathbb{R}} \circ i_{A, \mathbb{R}}^B \circ h_B$ belongs.

Let C be the abelian variety over E obtained similarly to B but starting from H_C^{der} (here we might have to replace once more E by a finite field extension of it). The abelian varieties A and $B \times_E C$ are adjoint-isogenous. For each simple factor H_j of $H_B^{\text{ad}} = H_B^{\prime, \text{ad}} = H_{\mathcal{D}}^{\text{ad}}$, there is $i \in I$ such that H_j is isomorphic to a simple factor of the adjoint group $H_{\mathcal{D}_i}^{\text{ad}}$ defined similarly to $H_{\mathcal{D}}^{\text{ad}}$ but for $\mathcal{D}_i$ instead of $\mathcal{D}$. As the Mumford–Tate conjecture holds for $\mathcal{D}_i$, the identity component of the algebraic envelope of the image of the p -adic Galois representation attached to B surjects naturally onto $H_{j, \mathbb{Q}_p}$. Thus this identity component surjects also onto $H_{B, \mathbb{Q}_p}^{\text{ad}}$, cf. Theorem 7.2. From this and Theorem 1.3.1 we get that the Mumford–Tate conjecture holds for B . Thus we have $H_{B, \mathbb{Q}_p}^{\text{ad}} = H_{\mathcal{D}, \mathbb{Q}_p}^{\text{ad}} = G_{\mathcal{D}, \mathbb{Q}_p}^{\text{ad}}$. $\square$

7.6.1. Example. This example is only meant to justify the name “independence property” for Theorems 1.3.7 and 7.6. We assume that $\text{End}(A_{\overline{\mathbb{Q}}}) = \mathbb{Z}$; thus the adjoint Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is simple (see [Pi, Prop. 5.12]). We also assume that $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is of $D_{2n+2}^{\mathbb{R}}$ type and that the group $H_{A,\mathbb{R}}^{\text{ad}}$ is not absolutely simple. Let $f_2 : (G_2, X_2) \hookrightarrow (\mathbf{GSp}(W_1, \psi_1), S_1)$ be an injective map as in Subsections 5.2 and 5.3 (thus $(G_2^{\text{ad}}, X_2^{\text{ad}}) = (H_A^{\text{ad}}, X_A^{\text{ad}})$). We assume that there exists an abelian variety B over E that is the pull back of an abelian scheme $\mathcal{A}_2$ as in Subsubsection 5.4.2. Let $C := A \times_E B$. We also assume that B is such that the adjoint group of the Mumford–Tate group H_C of C_C is $H_A^{\text{ad}} \times_{\mathbb{Q}} G_2^{\text{ad}} = H_A^{\text{ad}} \times_{\mathbb{Q}} H_A^{\text{ad}}$. The identity component of the algebraic envelope of the image of the p -adic Galois representation attached to C could theoretically be (without contradicting the results of Subsections 1.1.2 to 1.1.9 and of Section 3) a reductive group over $\mathbb{Q}_p$ whose adjoint is isomorphic to $H_{A,\mathbb{Q}_p}^{\text{ad}}$ (and thus it is different from $H_{C,\mathbb{Q}_p}^{\text{ad}}$). But Theorem 1.3.4 (e) implies that this identity component is $H_{C,\mathbb{Q}_p}$.

7.7. Concluding remarks. (a) Theorems 1.3.4 and 7.6 extend (this is standard) to abelian varieties over finitely generated fields of characteristic 0.

(b) We assume that $\text{End}(A_{\overline{E}}) = \mathbb{Z}$ and that the adjoint Shimura pair $(H_A^{\text{ad}}, X_A^{\text{ad}})$ is simple of $D_{2n+2}^{\mathbb{R}}$ type. Property 4.8 (vi) and the shifting process of Subsection 5.4 allow another proof of [Pi, Sect. 7, pp. 230–236] i.e., allow us to get directly the lifting property of l -adic Galois representations mentioned in [Pi, Sect. 7, top of p. 230] and due to Wintenberger.

(c) One can not get a general form of [Pi, Thm. 5.14] in the pattern of Theorem 1.3.4 (c) by using only the ideas of Sections 1 to 7. In a future paper we will present the general principle behind loc. cit. that extends significantly loc. cit. and this paper in the contexts when for each $j \in J$ the group $H_{j,\mathbb{R}}$ has only one simple, non-compact factor.

(d) For the left cases of the Mumford–Tate conjecture, one can assume that all simple factors of $H_{j,\overline{\mathbb{Q}}}$ have the same Lie type $\mathfrak{L}$ and (if it helps) that properties 5.10 (a), (b), or (c) holds (cf. Proposition 5.4.1, Subsubsection 5.4.3, and Theorem 7.2). If $\mathfrak{L} \neq D_4$, then we can also assume that the group H_A^{ad} is $\mathbb{Q}$ -simple (cf. Theorem 7.2).

(e) Corollary 7.4 can be used to strengthen both Theorem 7.6 and (d) for the case when $\mathfrak{L} = D_4$.

(f) Many ideas of this paper can be used for other classes of motives, provided (some of) the analogues of [De2, Subsubsects. 2.3.9 to 2.3.13], Theorem 1.1.4, [Sa1], and [Zi, Thm. 4.4] are known to be true in some form. In connection to Conjecture 1.1.1, the results [De2, Subsubsects. 2.3.9 to 2.3.13] and [Zi, Thm. 4.4] were neither used nor quoted prior to (earlier versions of) this work.

(g) By combining [An, Thm. 1.5.1 and Cor. 1.5.2] with Theorem 1.3.4, one gets a different proof of the result [An, Thm. 1.6.1 3)] that pertains to the Mumford–Tate conjecture for hyperkähler varieties over number fields.

(h) We have a variant of the notion of non-special A_n type (see Definition 6.1 (c)) in which the order of quantifiers is changed: there exists an f_1 such for all $\tilde{f}'_1$ and T_1 as in Definition 6.1 (b), there exist two distinct primes $q_1, q_2 \in \mathbb{N}$ with the property that for all

MT pairs (T_1, G_1) and (T_2, G_2) for $(f_1, \tilde{f}'_1, q_1)$ and $(f_1, \tilde{f}'_1, q_2)$ (respectively), both groups $G_{1, \mathbb{Q}_{q_1}}^{\text{ad}}$ and $G_{2, \mathbb{Q}_{q_2}}^{\text{ad}}$ have simple factors of A_n Lie type. Presently we do not know if such a variant could lead to significant improvements to Theorem 1.3.4 (a).

References

- [An] Y. André, *On the Shafarevich and Tate conjectures for hyperkähler varieties*, Math. Ann. **305** (1996), no. 2, pp. 205–248.
- [Bl] D. Blasius, *A p -adic property of Hodge cycles on abelian varieties*, Motives (Seattle, WA, 1991), pp. 293–308, Proc. Symp. Pure Math., Vol. **55**, Part 2, Amer. Math. Soc., Providence, RI, 1994.
- [Bog] F. A. Bogomolov, *Sur l'algébricité des représentations l -adiques*, C. R. Acad. Sci. Paris Sér. A-B Math. **290** (1980), no. 15, pp. 701–703.
- [Bor] A. Borel, *Linear algebraic groups. Second enlarged edition*, Grad. Texts in Math., Vol. **126**, Springer-Verlag, New York, 1991.
- [Bou1] N. Bourbaki, *Groupes et algèbres de Lie*, Chapitre **4–6**, Actualités Scientifiques et Industrielles, No. **1337**, Hermann, Paris, 1968.
- [Bou2] N. Bourbaki, *Groupes et algèbres de Lie*, Chapitre **7–8**, Actualités Scientifiques et Industrielles, No. **1364**, Hermann, Paris, 1975.
- [BHC] A. Borel and Harish-Chandra, *Arithmetic subgroups of algebraic groups*, Ann. of Math. (2) **75** (1962), no. 3, pp. 485–535.
- [BLR] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Néron models*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), Vol. **21**, Springer-Verlag, Berlin, 1990.
- [BO] P. Berthelot and A. Ogus, *F-crystals and de Rham cohomology. I*, Invent. Math. **72** (1983), no. 2, pp. 159–199.
- [Ch] W. C. Chi, *l -adic and λ -adic representations associated to abelian varieties defined over number fields*, Amer. J. Math. **114** (1992), no. 2, pp. 315–353.
- [De1] P. Deligne, *Travaux de Shimura*, Séminaire Bourbaki, 23ème année (1970/71), Exp. No. 389, pp. 123–163, Lecture Notes in Math., Vol. **244**, Springer-Verlag, Berlin, 1971.
- [De2] P. Deligne, *Variétés de Shimura: interprétation modulaire, et techniques de construction de modèles canoniques*, Automorphic forms, representations and L -functions (Oregon State Univ., Corvallis, OR, 1977), Part 2, pp. 247–289, Proc. Sympos. Pure Math., Vol. **33**, Amer. Math. Soc., Providence, RI, 1979.
- [De3] P. Deligne, *Hodge cycles on abelian varieties*, Hodge cycles, motives, and Shimura varieties, pp. 9–100, Lecture Notes in Math., Vol. **900**, Springer-Verlag, Berlin-New York, 1982.
- [DM] P. Deligne and J. Milne, *Tannakian categories*, Hodge cycles, motives, and Shimura varieties, pp. 101–128, Lecture Notes in Math., Vol. **900**, Springer-Verlag, Berlin-New York, 1982.
- [Fa] G. Faltings, *Endlichkeitssätze für abelsche Varietäten über Zahlkörpern*, Invent. Math. **73** (1983), no. 3, pp. 349–366.
- [Fo] J.-M. Fontaine, *Le corps des périodes p -adiques*, Périodes p -adiques (Bures-sur-Yvette, 1988), pp. 59–111, J. Astérisque **223**, Soc. Math. de France, Paris, 1994.

- [FH] W. Fulton and J. Harris, *Representation theory. A first course*, Grad. Texts in Math., Vol. **129**, Springer-Verlag, New York, 1991.
- [Ha] G. Harder, *Über die Galoiskohomologie halbeinfacher Matrizengruppen. II*, Math. Z. **92** (1966), pp. 396–415.
- [He] S. Helgason, *Differential geometry, Lie groups, and symmetric spaces*, Pure and Applied Mathematics, Vol. **80**, Academic Press, Inc. [Harcourt Brace Jovanovich, Publishers], New York-London, 1978.
- [Ko1] R. E. Kottwitz, *Stable trace formula: elliptic singular terms*, Math. Ann. **275** (1986), no. 3, pp. 365–399.
- [Ko2] R. E. Kottwitz, *Points on some Shimura Varieties over finite fields*, J. of Amer. Math. Soc. **5** (1992), no. 2, pp. 373–444.
- [La] S. Lang, *Algebraic number theory. Second edition*, Grad. Texts in Math., Vol. **110**, Springer-Verlag, New York, 1994.
- [LP] M. Larsen and R. Pink, *Abelian varieties, l -adic representations and l -independence*, Math. Ann. **302** (1995), no. 3, pp. 561–579.
- [Mi1] J. S. Milne, *The action of an automorphism of $\mathbb{C}$ on a Shimura variety and its special points*, Arithmetic and geometry, Vol. I, pp. 239–265, Progr. in Math., Vol. **35**, Birkhäuser, Boston, MA, 1983.
- [Mi2] J. S. Milne, *The points on a Shimura variety modulo a prime of good reduction, The Zeta functions of Picard modular surfaces*, pp. 151–253, Univ. Montréal, Montreal, Quebec, 1992.
- [Mi3] J. S. Milne, *Shimura varieties and motives*, Motives (Seattle, WA, 1991), pp. 447–523, Proc. Symp. Pure Math., Vol. **55**, Part 2, Amer. Math. Soc., Providence, RI, 1994.
- [Mi4] J. S. Milne, *Descent for Shimura varieties*, Mich. Math. J. **46** (1999), no. 1, pp. 203–208.
- [Mu1] D. Mumford, *Families of abelian varieties*, Algebraic Groups and Discontinuous Subgroups (Boulder, CO, 1965), pp. 347–351, Proc. Sympos. Pure Math., Vol. **9**, Amer. Math. Soc., Providence, RI, 1966.
- [Mu2] D. Mumford, *Abelian varieties*, Tata Inst. of Fund. Research Studies in Math., No. **5**, Published for the Tata Institute of Fundamental Research, Bombay; Oxford Univ. Press, London, 1970 (reprinted 1988).
- [No] R. Noot, *Classe de conjugaison du Frobenius des variétés abéliennes à réduction ordinaire*, Ann. Inst. Fourier (Grenoble) **45** (1995), no. 5, pp. 1239–1248.
- [Pi] R. Pink, *l -adic algebraic monodromy groups, cocharacters, and the Mumford–Tate conjecture*, J. reine angew. Math. **495** (1998), pp. 187–237.
- [Sa1] I. Satake, *Holomorphic imbeddings of symmetric domains into a Siegel space*, Amer. J. Math. **87** (1965), pp. 425–461.
- [Sa2] I. Satake, *Symplectic representations of algebraic groups satisfying a certain analyticity condition*, Acta Math. **117** (1967), pp. 215–279.
- [Se1] J.-P. Serre, *Groupes algébriques associés aux modules de Hodge–Tate*, Journées de Géom. Alg. de Rennes. (Rennes, 1978), Vol. III, pp. 155–188, J. Astérisque **65**, Soc. Math. de France, Paris, 1979.
- [Se2] J.-P. Serre, *Propriétés conjecturales des groupes de Galois motiviques et des*

- représentations l -adiques*, Motives (Seattle, WA, 1991), pp. 377–400, Proc. Sympos. Pure Math., Vol. **55**, Part 1, Amer. Math. Soc., Providence, RI, 1994.
- [Se3] J.-P. Serre, *Collected papers*, Vol. IV, 1985–1998, Springer-Verlag, Berlin, 2000.
- [Sh] G. Shimura, *Moduli of abelian varieties and number theory*, Algebraic Groups and Discontinuous Subgroups (Boulder, CO, 1965), pp. 312–332, Proc. Sympos. Pure Math., Vol. **9**, Amer. Math. Soc., Providence, RI, 1966.
- [Ta1] S. G. Tankeev, *Cycles on abelian varieties and exceptional numbers*, Izv. Ross. Akad. Nauk. Ser. Mat. **60** (1996), no. 2, pp. 159–194, translation in Izv. Math. **60** (1996), no. 2, pp. 391–424.
- [Ta2] S. G. Tankeev, *On the weights of an l -adic representation and the arithmetic of Frobenius eigenvalues*, (Russian) Izv. Ross. Akad. Nauk Ser. Mat. **63** (1999), no. 1, pp. 185–224, translation in Izv. Math. **63** (1999), no. 1, pp. 181–218.
- [Tat] J. Tate, *Endomorphisms of abelian varieties over finite fields*, Invent. Math. **2** (1966), pp. 134–144.
- [Ts] T. Tsuji, *p -adic étale cohomology and crystalline cohomology in the semi-stable reduction case*, Invent. Math. **137** (1999), no. 2, pp. 233–411.
- [Va1] A. Vasiu, *Integral canonical models of Shimura varieties of preabelian type*, Asian J. Math. **3** (1999), no. 2, pp. 401–518.
- [Va2] A. Vasiu, *Surjectivity criteria for p -adic representations, Part I*, Manuscripta Math. **112** (2003), no. 3, pp. 325–355.
- [Za1] Y. G. Zarhin, *Abelian varieties having a reduction of K3 type*, Duke Math. J. **65** (1992), no. 3, pp. 511–527.
- [Za2] Y. Zarhin, *Very simple 2-adic representations and hyperelliptic Jacobians*, Moscow Math. J. **2** (2002), no. 2, pp. 403–431.
- [Zi] T. Zink, *Isogenieklassen von Punkten von Shimuramannigfaltigkeiten mit Werten in einem endlichen Körper*, Math. Nachr. **112** (1983), pp. 103–124.
- [Wi] J.-P. Wintenberger, *Un scindage de la filtration de Hodge pour certaines variétés algebriques sur les corps locaux*, Ann. of Math. (2) **119** (1984), no. 3, pp. 511–548.

Adrian Vasiu,
 Department of Mathematical Sciences, Binghamton University,
 Binghamton, New York 13902-6000, U.S.A.
 e-mail: adrian@math.binghamton.edu